



Timing-dependent valence reversal: a principle of reinforcement processing and its possible implications

Bertram Gerber^{1,2,3}, Christian König^{1,4}, Markus Fendt^{3,5},
Marta Andreatta⁶, Marcel Romanos⁸, Paul Pauli^{6,7} and
Ayse Yarali^{3,4}

Punishment feels bad, but relief upon its termination feels good. As a consequence of such timing-dependent valence reversal, memories of opposite valence can result from associating stimulus A with, for example, the occurrence of punishment (A–) versus punishment termination (–A): A– training results in aversive memory, but –A training in appetitive memory (corresponding effects exist for reward occurrence and termination). Whereas learning through the occurrence of punishment is well studied, much less is known about learning through its termination. Current research investigates how dopaminergic system function contributes to these processes in *Drosophila*, rats and humans. We argue that dopamine-related psychopathology may entail distortions in learning through punishment termination, and that this may contribute, for example, to non-suicidal self-injury or post-traumatic stress disorder.

Addresses

¹ Leibniz Institute for Neurobiology, Genetics of Learning and Memory, Magdeburg, Germany

² Otto von Guericke University, Institute for Biology, Behavioral Genetics, Magdeburg, Germany

³ Center for Behavioral Brain Sciences, Magdeburg, Germany

⁴ Leibniz Institute for Neurobiology, Molecular Systems Biology of Learning, Magdeburg, Germany

⁵ Otto-von-Guericke University, Institute for Pharmacology and Toxicology, Magdeburg, Germany

⁶ University of Würzburg, Department of Psychology (Biological Psychology, Clinical Psychology, and Psychotherapy), Würzburg, Germany

⁷ University of Würzburg, Center of Mental Health, Würzburg, Germany

⁸ University Hospital Würzburg, Center of Mental Health, Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, Würzburg, Germany

Corresponding author:

Gerber, Bertram (bertram.gerber@lin-magdeburg.de)

Current Opinion in Behavioral Sciences 2019, 26:114–120

This review comes from a themed issue on **Pain and aversive motivation**

Edited by **Ben Seymour** and **Joshua Johansen**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 11th January 2019

<https://doi.org/10.1016/j.cobeha.2018.12.001>

2352-1546/© 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).


Introduction

Avoiding punishment and obtaining reward are evolutionarily deeply rooted behavioral goals. However, half of the processes through which punishment and reward affect behavior are notably understudied. That is, past research has provided detailed insight into how we learn about predictors of the occurrence of punishment and reward [1–4], but much less is known about how we learn about their termination [5]. Since the occurrence of punishment feels bad, but its termination feels good (Figure 1), associating stimuli with these experiences results in memories of opposite valence, namely in aversive and appetitive memory, respectively. Conversely, the occurrence versus the termination of reward supports appetitive versus aversive learning. There are thus four, not just two, predictive relations between stimuli and the reinforcement paired with them (Figure 1) [5]. This reversal in memory valence through a switch in event timing is called timing-dependent valence reversal. The present paper argues that timing-dependent valence reversal reflects a general principle of how reinforcement is processed; it details what has recently been revealed about the role of dopaminergic neurons in timing-dependent valence reversal; and it explores the possible impact of distortions in timing-dependent valence reversal on mental health.

Generality of timing-dependent valence reversal

Emotionally significant events, in addition to the initial, primary affect that their occurrence induces, also induce a

Figure 1



Timing-dependent valence reversal.

Left: The occurrence of punishment induces negative affect (Pain), while its termination induces positive affect (Relief). Likewise, the occurrence versus termination of reward feels good and bad, respectively (not shown). All four of these affective states can enter into association with stimuli paired with them. **Right:** Of the four predictive relations of stimuli (A) and reward (+) or punishment (-), two are notably understudied (open boxes). Learning through unpaired presentations of stimuli and reinforcement is not covered in this paper (discussion in Ref. [5]).

secondary and oppositely-valenced after-affect upon their termination. This goes both for events of a primary-aversive and of a primary-appetitive nature (Figure 1) [6]. These four types of affect can induce memories for stimuli associated with them ([5,7,8] and references therein). Typically, stimuli associated with the occurrence of punishment (A-) acquire negative learned valence, whereas stimuli associated with the termination of punishment (-A) acquire positive learned valence. Using punishment, this has been observed, for example, in startle-modulation by Pavlovian learning in humans and rats, eyelid conditioning in rabbits, Sidman avoidance learning in rats, and in olfactory as well as visual Pavlovian conditioning in the fly *Drosophila*. This called for an extension of the threat imminence model [9,10] to include the effects produced upon the termination of punishment ('post-strike' phase: [5]). Likewise, stimuli presented with reward occurrence (A+) and termination (+A) acquire positive and negative learned valence, respectively, as shown for odor-sugar associative learning in honeybees. These findings of timing-dependent valence reversal across species, paradigms, stimulus modalities and valence domains prompted major revisions of the prediction-error learning rule [11]. It remains unresolved how these behavioral effects, which are observed across notably different time scales in the various species and paradigms, relate to molecular coincidence detection and/or to spike-timing-dependent synaptic plasticity observed in the second to millisecond range (discussion in Refs. [5,12]).


Timing-dependent valence reversal in *Drosophila*

In the most widely used Pavlovian conditioning paradigm for *Drosophila*, the flies are presented with an odor A

followed by electric shock punishment (A-), and a second odor B unpaired from punishment. Such A-/B training results in a relative avoidance of A in a subsequent test of choice between the two odors. By contrast, presenting odor A upon the termination of punishment (-A/B training) leads to a relative approach toward A. Thus, the paradigm is bivalent: the balance in the choice between A and B can be tipped either way, making it possible to reveal learned avoidance or approach, respectively, within the same paradigm. Follow-up experiments have shown that memory after -A training is weaker than after A- training, requires a longer inter-stimulus-interval (i.e. a longer time interval between punishment and A), is best for mild punishment, requires more training trials (cf. Ref. [5]), and increases with punishment duration [13^{**}]; the latter conforms to the classical proposal of Ref. [6]. Memory scores established through A- and through -A training show non-correlating variation across inbred fly strains derived from the wild, suggesting considerable non-overlap of their genetic determinants [14]; whether the same is the case for memories after -A and A+ training remains to be tested. It was further revealed that odor-specific associative memory shortly after A- training is composed of two components, one that is dependent on Synapsin [15], an evolutionarily conserved presynaptic protein known to regulate synaptic strength, and is susceptible to cold-amnesia (cf. Ref. [5]), and one that is Synapsin-independent and resistant to cold-amnesia. Memory after -A training, by contrast, is only of the former kind, that is Synapsin-dependent and susceptible to cold-amnesia. What is known about the circuit-level localization of the engrams after A- and -A training?

The concerted efforts of the *Drosophila* community suggest that the engram underlying odor-specific associative short-term memory after A- training can be localized to a single layer of synapses in third-order olfactory neurons of the fly brain (reviewed in Ref. [16]). Specifically, the engram is apparently local to the presynaptic terminals of those mushroom body Kenyon cells that are activated by odor A (Figures 2, S1), which we will call A = KCs. Learned avoidance of A is thought to come about through an odor-specific depression in the excitatory, cholinergic connection of the A = KCs to approach-promoting mushroom body output neurons (MBONs) [17-20,21^{*},22,23]. To the limited extent tested, this same synaptic layer also appears to harbor the engram after -A training [13^{**},15]. Thus, A- training can evidently establish depression whereas -A training might possibly establish potentiation of these A = KC-MBON synapses (indeed, these synapses have been shown to be capable of potentiation upon associative training: [24-26; also see Refs. 20,27-29]). But how is punishment signaled to the KCs? Answering this question has been facilitated by a breakthrough in understanding

Figure 2



Circuit and microcircuit for associative learning in flies.

Left: Simplified working hypothesis of olfactory associative learning in flies. Odor A (orange cloud) activates a subset of KCs (orange fill). In these KCs, a coincidence can be detected with modulatory signaling from DANs that intersect the mushroom body and that convey either punishment or reward information. Such coincidence can lead to a depression of the synapses from the respective KCs to the MBONs, indicated by the irregular star. In the case of A– learning, reduced activation of an approach-mediating MBON would lead to learned avoidance because avoidance tendencies through non-depressed MBONs in other mushroom body compartments would prevail (for more detail see Figure S1). A+ learning is thought to come about in an analogous way. According to this scenario, A– and –A learning would take place by depression and potentiation, respectively, within the same compartment, and the same would be the case, in a separate compartment, for A+ and +A learning. Grey connections between ascending punishment and reward signaling imply the possibility of mutual inhibition between them, and thus of post-inhibitory rebound activation. This mutual inhibition could also be indirect, via for example cross-compartmental feedback from MBONs. It would imply as an alternative scenario that A– and +A learning as well as –A and A+ learning take place in the same compartment, respectively, and through depression in all cases. The organization of innate olfactory, punishment- and reward-related behavior largely bypasses the mushroom body. Please note that for simplicity KC-KC, KC-DAN, DAN-MBON, and MBON-MBON synapses are omitted from this figure; how they contribute to the learning processes discussed in this paper is as yet unknown. **Right:** Schematic of a mushroom body compartment with the chemical synapses between KCs, DANs and MBONs as determined from electron microscopy (data from Refs. [30,31]). The stippled box indicates that MBON-MBON connections are located largely outside the mushroom body. The fly image is taken from Ref. [51], copyright Elsevier.

the cellular and synaptic architecture of the mushroom body in notable detail and completeness [18,30,31]. This architecture is reviewed here in simplified form.

The KCs receive excitatory, cholinergic input from second-order neurons of multiple sensory modalities, including olfactory projection neurons, establishing a sparse representation of the flies' sensory environment. The KC axons form a parallel bundle intersected by the terminals of ascending modulatory neurons. Typically, these modulatory neurons are either octopaminergic or dopaminergic (OANs, DANs). The modulatory effects exerted on the KCs may be complex, given that multiple types of octopamine and dopamine receptor can be expressed in KCs, and given that the presence of typically more than one morphological type of synaptic vesicle

suggests that further signaling molecules might be released as well. In any event, the DANs can be broadly classified as mediating either punishment or reward (a similar situation may be emerging in mammals as well: [32–35]). For a given DAN the DAN-KC synapses are restricted to only a small region along the KCs axons, non-overlapping with neighboring DANs. These regions are also respected by the dendritic branches of the MBONs. This results in a peculiar and valenced compartmental structure of the mushroom body (Figure 2).

It has been found that DANs mediating punishment information target compartments whose MBONs are approach-promoting [19]. This is led to the mentioned working hypothesis that learned avoidance after A– training comes about by a depression of the A=KC-MBON


synapses, and learned approach after $-A$ training by their potentiation. In turn, DANs mediating reward information target compartments with avoidance-promoting MBONs, suggesting that the same logic applies for $A+$ and $+A$ learning. If this were so, and in line with both spike-timing-dependent plasticity (STDP) at the KC-MBON synapse [26] and an earlier theoretical approach [36], the onset and the offset of the activation of a given DAN should confer timing-dependent valence reversal. Is this indeed the case?

Individual DANs can mediate memories of opposing valence, dependent on timing

In *Drosophila*, transgenic effectors can be expressed permitting non-invasive manipulation of neuronal activity in awake, freely behaving animals. This allows high-temporal-resolution control of activity in hemispherically single, identified DANs with known and bilaterally highly symmetrical synaptic connectivity. This has revealed that the activation of DANs called PPL1-01 can produce timing-dependent valence reversal [13^{**},37^{**}] (synonyms for PPL1-01 are: PPL1- γ 1pedc, MB-MP1, and MP). That is, similar to what has been found for $A-$ and $-A$ training, presenting odor A with the onset of PPL1-01 activation resulted in learned avoidance of A , whereas presenting odor A with the offset of PPL1-01 activation resulted in learned approach (Figure 3). Thus, the same dopaminergic neurons can mediate memories of opposing valence, depending on the timing of their activation (a similar result was obtained, in larval *Drosophila*, in the appetitive domain for the DAN called DAN-i1 [38^{*}]).

These results are consistent with a scenario whereby the above-mentioned dopaminergic and STDP-like mechanism at the KC-MBON synapse underlies timing-dependent valence reversal. They imply that a given DAN can induce memories through both $A-$ and $-A$ training. However, the observation that different compartments with DANs processing punishment and reward information, respectively, show alternating cycles of activation and inhibition [20] has prompted an alternative scenario [13^{**}]. What if PPL1-01 inhibits, possibly via a non-dopaminergic mechanism, a DAN mediating reward signals? Once PPL1-01 activation is terminated that reward-mediating DAN may not only be released from inhibition but may show post-inhibitory rebound activation, which is a frequently observed physiological phenomenon (see following section). Such post-inhibitory rebound activation of a rewarding DAN could produce a depression of the $A = KC$ -MBON synapses—in ‘its own’ compartment. In other words, the alternative scenario is that $A-$ and $-A$ learning come about *through different DANs*, and that both do involve the depression of $A = KC$ -MBON synapses yet *in different compartments*. In this case, and in contrast to the STDP-scenario detailed above, a given DAN would mediate the effects of both $A+$ and $-A$ training. The proposed inhibition between oppositely-

Figure 3




Timing-dependent valence reversal in flies.

Top: Summary of timing-dependent valence reversal in flies across studies. Different sets of flies received either $A-$ training at the respectively indicated negative inter-stimulus-intervals (ISIs) or $-A$ training (positive ISIs). The lightning bolt indicates the period of electric shock delivery. The memory scores quantify the extent to which, relative to flies receiving unpaired presentations of odor A and shock, $A-$ training leads to learned avoidance of A (negative memory scores), whereas $-A$ training leads to learned approach (positive memory scores). Each dot reflects the median memory score of on average $N=22$ sets of $n=150$ flies each (total $N=919$, total n approx. 140 000). Dots sharing the same color come from the same study. The red and green areas under the curve are meant to facilitate comparison to Figure 1. The raw data and references to their original publication can be found in Table S1. **Bottom:** Flies expressing the blue-light-gated cation channel ChR2-XXL in the DAN called PPL1-01 were trained as in the top panel, except that instead of electric shock blue light was turned on (blue star and vertical line). This reveals timing-dependent valence reversal through PPL1-01 activation. The box plots represent the median, 25/75, and 10/90% quantiles as middle line, box boundaries, and whiskers, respectively, of on average $N=23$ sets of flies with $n=150$ animals each (data from Ref. [13^{**}]; Table S1). Coloring of the plots indicates significant difference from chance. The inset sketch shows how PPL1-01 innervates the mushroom body (after [18]) (data from Ref. [13^{**}]; Table S1) (the fly image is taken from Ref. [51], copyright Elsevier).

valenced DANs is unlikely to come about by direct chemical synapses, however, as no such connections have yet been revealed [30,31^{*}].

Figure 4



Timing-dependent valence reversal in rats and humans.

Top: Different groups of rats were submitted either to A– training (negative ISIs) or –A training (positive ISIs). A 5-s light was used as stimulus A and combined with a mild electric shock punishment of 0.5 s duration, indicated by the lightning bolt. In two different control groups (open plots), either only stimulus A was presented, or A and punishment were presented in a random temporal relationship. One day later, the effect of stimulus A on the magnitude of the auditory startle response was measured. Plotted is the mean percent difference between startle magnitude in the presence versus in the absence of A. Startle attenuation, that is positive valence, is plotted toward the top of the Y-axis, whereas startle potentiation, that is negative valence, is plotted toward the bottom of the Y-axis. Coloring of the plots indicates significant differences from the median of the random-control (stippled line) (data from Refs. [43,44]; Table S1). Other details as in Figure 3, bottom. **Bottom:** Summary of timing-dependent valence reversal in humans across studies. A– training (negative ISIs) and –A training (positive ISIs) were implemented in either within-subject or between-subject protocols, as detailed in Ref. [52], using 8-s-duration visual geometric shapes displayed on a computer screen as A and 0.2-s-duration mild electric shock as punishment, indicated by the lightning bolt. The magnitude of the startle responses was transformed into standard scores (z-scores). The colored dots indicate the mean results of the respective treatment condition, with startle attenuation, that is implicit positive valence, plotted toward the top of the Y-axis, whereas startle potentiation, that is implicit negative valence, is plotted toward the bottom of the Y-axis. Triangles reflect startle modulation by a within-subject *Control* stimulus that was presented at a very long ISI (15–25 s, dependent on study; see Ref. [52]). The red and green areas under the curve are meant to facilitate comparison to Figure 1. Data are taken from Ref. [52] (Table S1), and reflect the performance of N=214 subjects in total.

Timing-dependent valence reversal in rats and humans

A workhorse paradigm for current studies on timing-dependent valence reversal in mammals is the modulation of the evolutionarily conserved auditory startle reflex [39–42]. Rats and humans exhibit potentiated startle in the presence of a visual stimulus A previously paired with mild electric shock punishment (A–), whereas they show attenuated startle after appetitive training (A+). Intense research into the mechanisms of such A– and A+ learning has revealed a conserved role of the amygdala for A– learning and of the dopaminergic inputs to the nucleus accumbens of the striatum for A+ learning.

As startle modulation is a bivalent measure, it lent itself to showing timing-dependent valence reversal in mammals. Indeed, an attenuation of startle and thus positive valence was revealed after –A training (Figure 4), as was a double dissociation of the brain regions for –A and A– learning [43]: in rats, –A learning was impaired by acute inactivation of the ventral striatum but not of the amygdala, whereas A– learning was impaired by inactivation of the amygdala but not of the ventral striatum. Fittingly, functional imaging in humans showed that A activates the ventral striatum upon –A training but not upon A– training, whereas the amygdala was activated by A upon A– but not upon –A training. Recent data from the rat show that it is specifically dopaminergic transmission from the posterior-medial ventral tegmental area to the nucleus accumbens shell that is required for –A learning but dispensable for A– learning [44,45**]. Whether dopaminergic transmission can also bring about timing-dependent valence reversal, whether corresponding dissociations can be observed in the appetitive domain (A+ versus +A), what the role of other dopaminergic projections is, and whether the same or different dopaminergic neurons mediate these effects, are questions that remain to be investigated. We note that rebound activation of dopamine neurons upon the termination of aversive stimulation is well documented [35,46–48].

Possible implications

The central basic-research question underlying the above discussion is whether different subsets of dopamine neurons and/or receptors are involved in different aspects of timing-dependent valence reversal and across valence domains. We believe that answering this question will have implications for understanding, and treating, distortions in motivated behavior in humans and its concomitant psychopathologies. For example, ‘excessively strong’ relief may help in understanding pathological risk-taking or non-suicidal self-cutting as behaviors seeking to bring about relief [49]. But does this indeed come about by exaggerated –A learning, which under normal conditions is a rather modest effect (Figure 4), or rather by blunted A– learning? Or both? Is it related to dopaminergic hyperfunction or hypo-function, or maybe to both—in different

sets of dopamine neurons and/or employing different dopamine receptors? What does this mean for the treatment of dopaminergic dysfunction? Corresponding questions arise concerning ‘excessively weak’ relief. That is, maybe the risk of post-traumatic stress disorder is increased when during the post-encounter phase subjects mnemonically focus on the adverse emotions experienced during trauma, rather than on the relief of having survived it [50]? Might a distorted mnemonic focus also contribute to the pathological avoidance behavior characteristic of anxiety disorders? Again, it would have opposing implications depending on whether in either case this comes about by blunted –A learning, by exaggerated A– learning, or both, and whether the affected transmitter systems are hypo-functional, hyper-functional, or, in different sets of cells or in relation to different dopamine receptors, both. In all the above-mentioned cases, the aim for treatment, and possibly the aim for prevention as well, would thus need to be to restore a proper balance between the four types of predictive learning of stimuli and reinforcement (Figure 1).

Conclusions

The available evidence suggests us that timing-dependent valence reversal is a general principle of how punishment and probably also reward are processed. Given that avoiding punishment and obtaining reward are important goals of behavior, we believe that distortions in timing-dependent valence reversal can have a significant, and hitherto insufficiently appreciated, impact on motivated behavior and related psychopathology. Because dopaminergic system function emerges as a determinant for timing-dependent valence reversal across species, we are optimistic that the understanding, treatment, and prevention of these pathologies will benefit from analyses that focus specifically upon dopamine system function in learning from the occurrence and the termination of punishment and reward.

Conflict of interest statement

Nothing declared.

Acknowledgements

Institutional support: Otto von Guericke Universität Magdeburg, Wissenschaftsgemeinschaft Gottfried Wilhelm Leibniz (WGL), Leibniz Institut für Neurobiologie (LIN). Grant support: Deutsche Forschungsgemeinschaft (DFG) (GE 1091/4-1, CRC 779-B11 and FOR 2705-TP2 to BG; CRC 779-B15 to AY; FE 483/7-1 and CRC779-B13 to MF; CRC-TRR 58 B01 and B08 to PP and MA, respectively, and Z02 to MR and PP), and Schram-Stiftung (T287/25458/2013 to AY).

Comments from Philippe Tobler (U Zürich), Stephanie Wegener (HHMI Janelia, LIN Magdeburg), Thomas Niewalda and Michael Schleyer (LIN Magdeburg) on earlier versions of this article are gratefully acknowledged. We thank V. Ruta (Rockefeller U) and Y. Aso (HHMI Janelia) for sharing thoughts and unpublished data, and R.D.V. Glasgow (Zaragoza, Spain) for language editing.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.cobeha.2018.12.001>.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Heisenberg M: **Mushroom body memoir: from maps to models.** *Nat Rev Neurosci* 2003, **4**:266–275.
 2. LeDoux J: **Rethinking the emotional brain.** *Neuron* 2012, **73**:653–676.
 3. Hayes DJ, Duncan NW, Xu J, Northoff G: **A comparison of neural responses to appetitive and aversive stimuli in humans and other mammals.** *Neurosci Biobehav Rev* 2014, **45**:350–368.
 4. Schultz W: **Neuronal reward and decision signals: from theories to data.** *Physiol Rev* 2015, **95**:853–951.
 5. Gerber B, Yarali A, Diegelmann S, Wotjak CT, Pauli P, Fendt M: **Pain-relief learning in flies, rats, and man: basic research and applied perspectives.** *Learn Mem* 2014, **21**:232–252.
 6. Solomon RL, Corbit JD: **An opponent-process theory of motivation. I. Temporal dynamics of affect.** *Psychol Rev* 1974, **81**:119–145.
 7. Hellstern F, Malaka R, Hammer M: **Backward inhibitory learning in honeybees: a behavioral analysis of reinforcement processing.** *Learn Mem* 1998, **4**:429–444.
 8. Vogt K, Yarali A, Tanimoto H: **Reversing stimulus timing in visual conditioning leads to memories with opposite valence in *Drosophila*.** *PLoS One* 2015, **10**:e0139797.
 9. Blanchard RJ, Blanchard DC: **Antipredator defensive behaviors in a visible burrow system.** *J Comp Psychol* 1989, **103**:70–82.
 10. Fanselow MS: **Neural organization of the defensive behavior system responsible for fear.** *Psychonom Bull Rev* 1994, **1**:429–438.
 11. Malaka R: **Models of classical conditioning.** *Bull Math Biol* 1999, **61**:33–83.
 12. Suvrathan A: **Beyond STDP-towards diverse and functionally relevant plasticity rules.** *Curr Opin Neurobiol* 2018, **54**:12–19.
 13. König C, Khalili A, Ganesan M, Nishu AP, Garza AP, Niewalda T, •• Gerber B, Aso Y, Yarali A: **Reinforcement signaling of punishment versus relief in fruit flies.** *Learn Mem* 2018, **25**:247–257.
- Including the proper genetic controls, the study demonstrates that optogenetic activation of the PPL1-01 dopamine neurons can produce timing-dependent valence reversal in *Drosophila* (see also Ref. [37]). Further, an RNAi knockdown of the enzyme needed for dopamine biosynthesis in the very neurons that are being activated reduces aversive memory upon A-then-PPL1-01 training, but leaves appetitive memory after training with the reverse timing unaffected. Concerns of a possibly incomplete knockdown by RNAi aside, this may suggest that a non-dopaminergic mechanism contributes to memory established by PPL1-01-then-A training.
14. Appel M, Scholz CJ, Kocabay S, Savage S, König C, Yarali A: **Independent natural genetic variation of punishment-versus relief-memory.** *Biol Lett* 2016, **12**:20160657.
 15. Niewalda T, Michels B, Jungnickel R, Diegelmann S, Kleber J, Kähne T, Gerber B: **Synapsin determines memory strength after punishment- and relief-learning.** *J Neurosci* 2015, **35**:7487–7502.
 16. Gerber B, Aso Y: **Localization, diversity and behavioral expression of associative engrams in *Drosophila*.** In *Learning Theory and Behavior, R Menzel (Ed.). Vol 1 of Learning and Memory: A Comprehensive Reference*. Edited by Byrne J. Oxford: Elsevier; 2017:463–473.

17. Séjourné J, Plaçais PY, Aso Y, Siwanowicz I, Trannoy S, Thoma V, Tedjakumala SR, Rubin GM, Tchénio P, Ito K *et al.*: **Mushroom body efferent neurons responsible for aversive olfactory memory retrieval in *Drosophila***. *Nat Neurosci* 2011, **14**:903-910.
18. Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo TT, Dionne H, Abbott LF, Axel R, Tanimoto H, Rubin GM: **The neuronal architecture of the mushroom body provides a logic for associative learning**. *Elife* 2014, **3**:e04577.
19. Aso Y, Sitaraman D, Ichinose T, Kaun KR, Vogt K, Belliard-Guerin G, Plaçais PY, Robie AA, Yamagata N, Schnaitmann C *et al.*: **Mushroom body output neurons encode valence and guide memory-based action selection in *Drosophila***. *Elife* 2014, **3**:e04580.
20. Cohn R, Morantte I, Ruta V: **Coordinated and compartmentalized neuromodulation shapes sensory processing in *Drosophila***. *Cell* 2015, **163**:1742-1755.
21. Hige T, Aso Y, Modi MN, Rubin GM, Turner GC: **Heterosynaptic plasticity underlies aversive olfactory learning in *Drosophila***. *Neuron* 2015, **88**:985-998.
- Demonstrates that pairings of odor with optogenetic activation of the PPL1-01 neurons establish both aversive associative short-term memory, and synaptic depression of the KC-MBON synapses. This depression is independent of MBON spiking, suggesting a presynaptic mechanism.
22. Barnstedt O, Oswald D, Felsenberg J, Brain R, Moszynski JP, Talbot CB, Perrat PN, Waddell S: **Memory-relevant mushroom body output synapses are cholinergic**. *Neuron* 2016, **89**:1237-1247.
23. Perisse E, Oswald D, Barnstedt O, Talbot CB, Huetteroth W, Waddell S: **Aversive learning and appetitive motivation toggle feed-forward inhibition in the *Drosophila* mushroom body**. *Neuron* 2016, **90**:1086-1099.
24. Maelshagen J: **Neural correlates of olfactory learning paradigms in an identified neuron in the honeybee brain**. *J Neurophysiol* 1993, **69**:609-625.
25. Menzel R, Man G: **Neural plasticity of mushroom body-extrinsic neurons in the honeybee brain**. *J Exp Biol* 2005, **208**:4317-4332.
26. Cassenaer S, Laurent G: **Conditional modulation of spike-timing-dependent plasticity for olfactory learning**. *Nature* 2012, **482**:47-52.
27. Bouzaiane E, Trannoy S, Scheunemann L, Plaçais PY, Preat T: **Two independent mushroom body output circuits retrieve the six discrete components of *Drosophila* aversive memory**. *Cell Rep* 2015, **11**:1280-1292.
28. Oswald D, Felsenberg J, Talbot CB, Das G, Perisse E, Huetteroth W, Waddell S: **Activity of defined mushroom body output neurons underlies learned olfactory behavior in *Drosophila***. *Neuron* 2015, **86**:417-427.
29. Felsenberg J, Barnstedt O, Cognigni P, Lin S, Waddell S: **Re-evaluation of learned information in *Drosophila***. *Nature* 2017, **544**:240-244.
30. Eichler K, Li F, Litwin-Kumar A, Park Y, Andrade I, Schneider-Mizell CM, Saumweber T, Huser A, Eschbach C, Gerber B *et al.*: **The complete connectome of a learning and memory centre in an insect brain**. *Nature* 2017, **548**:175-182.
31. Takemura SY, Aso Y, Hige T, Wong A, Lu Z, Xu CS, Rivlin PK, Hess H, Zhao T, Parag T *et al.*: **A connectome of a learning and memory center in the adult *Drosophila* brain**. *Elife* 2017, **6**:e26975.
- Chemical-synapse connectome of the alpha lobe of the adult *Drosophila* mushroom body, confirming the circuit motifs described in the larva [30]. These include the previously known DAN-KC and KC-MBON synapses, plus the hitherto uncharacterized KC-DAN, DAN-MBON, and KC-KC synapses.
32. Matsumoto M, Takada M: **Distinct representations of cognitive and motivational signals in midbrain dopamine neurons**. *Neuron* 2013, **79**:1011-1024.
33. Bromberg-Martin ES, Matsumoto M, Hikosaka O: **Dopamine in motivational control: rewarding, aversive, and alerting**. *Neuron* 2010, **68**:815-834.
34. Menegas W, Akiti K, Amo R, Uchida N, Watabe-Uchida M: **Dopamine neurons projecting to the posterior striatum reinforce avoidance of threatening stimuli**. *Nat Neurosci* 2018, **21**:1421-1430.
35. Brischoux F, Chakraborty S, Brierley DI, Ungless MA: **Phasic excitation of dopamine neurons in ventral VTA by noxious stimuli**. *Proc Natl Acad Sci U S A* 2009, **106**:4894-4899.
36. Yarali A, Nehr Korn J, Tanimoto H, Herz AV: **Event timing in associative learning: from biochemical reaction dynamics to behavioural observations**. *PLoS One* 2012, **7**:e32885.
37. Aso Y, Rubin GM: **Dopaminergic neurons write and update memories with cell-type-specific rules**. *Elife* 2016, **5**:e16135.
- The study demonstrates that optogenetic activation of the PPL1-01 dopamine neurons can produce timing-dependent valence reversal in *Drosophila* (see also Ref. [13]). Both studies further test for similar effects in additional MBONs.
38. Saumweber T, Rohwedder A, Schleyer M, Eichler K, Chen YC, Aso Y, Cardona A, Eschbach C, Kobler O, Voigt A *et al.*: **Functional architecture of reward learning in mushroom body extrinsic neurons of larval *Drosophila***. *Nat Commun* 2018, **9**:1104.
- Together with Ref. [30], this study provides a full atlas and chemical-synapse connectome of the KCs, DANs and MBONs of larval *Drosophila*. Further, it describes timing-dependent valence reversal in the appetitive domain (A+ and +A) through the dopaminergic DAN-i1 neuron.
39. Grillon C, Baas J: **A review of the modulation of the startle reflex by affective states and its application in psychiatry**. *Clin Neurophysiol* 2003, **114**:1557-1579.
40. Davis M, Antoniadis EA, Amaral DG, Winslow JT: **Acoustic startle reflex in rhesus monkeys: a review**. *Rev Neurosci* 2008, **19**:171-185.
41. Fendt M, Koch M: **Translational value of startle modulations**. *Cell Tissue Res* 2013, **354**:287-295.
42. Andreatta M, Pauli P: **Appetitive vs. aversive conditioning in humans**. *Front Behav Neurosci* 2015, **9**:128.
43. Andreatta M, Fendt M, Mühlberger A, Wieser MJ, Imobersteg S, Yarali A, Gerber B, Pauli P: **Onset and offset of aversive events establish distinct memories requiring fear and reward networks**. *Learn Mem* 2012, **19**:518-526.
44. Bergado Acosta JR, Kahl E, Kogias G, Uzuneser TC, Fendt M: **Relief learning requires a coincident activation of dopamine D1 and NMDA receptors within the nucleus accumbens**. *Neuropharmacology* 2017, **114**:58-66.
45. Mayer D, Kahl E, Uzuneser TC, Fendt M: **Role of the mesolimbic dopamine system in relief learning**. *Neuropsychopharmacology* 2018, **43**:1651-1659.
- The study establishes that a dopaminergic pathway from the posterior-medial ventral tegmental area to the nucleus accumbens shell is activated by aversive stimuli, and that this pathway is required for -A but not for A-learning in rats.
46. Wang DV, Tsien JZ: **Convergent processing of both positive and negative motivational signals by the VTA dopamine neuronal populations**. *PLoS One* 2011, **6**:e17047.
47. Fiorillo CD, Song MR, Yun SR: **Multiphasic temporal dynamics in responses of midbrain dopamine neurons to appetitive and aversive stimuli**. *J Neurosci* 2013, **33**:4710-4725.
48. Fiorillo CD, Yun SR, Song MR: **Diversity and homogeneity in responses of midbrain dopamine neurons**. *J Neurosci* 2013, **33**:4693-4709.
49. Hooley JM, Franklin JC: **Why do people hurt themselves? A new conceptual model of nonsuicidal self-injury**. *Clin Psychol Sci* 2018, **6**:428-451.
50. Lee JC, Wang LP, Tsien JZ: **Dopamine rebound-excitation theory: putting brakes on PTSD**. *Front Psychiatry* 2016, **7**:163.
51. Chyb S, Gompel N: *Atlas of *Drosophila* Morphology*. Academic Press; 2013.
52. Andreatta M, Mühlberger A, Pauli P: **When does pleasure start after the end of pain? The time course of relief**. *J Comp Neurol* 2016, **524**:1653-1667.