

Review

Structural and mechanistic basis of RNA processing by protein-only ribonuclease P enzymes

Arjun Bhatta ^{1,2} and Hauke S. Hillen ^{1,2,3,*,@}

Ribonuclease P (RNase P) enzymes are responsible for the 5' processing of tRNA precursors. In addition to the well-characterised ribozyme-based RNase P enzymes, an evolutionarily distinct group of protein-only RNase Ps exists. These proteinaceous RNase Ps (PRORPs) can be found in all three domains of life and can be divided into two structurally different types: eukaryotic and prokaryotic. Recent structural studies on members of both families reveal a surprising diversity of molecular architectures, but also highlight conceptual and mechanistic similarities. Here, we provide a comparison between the different types of PRORP enzymes and review how the combination of structural, biochemical, and biophysical studies has led to a molecular picture of protein-mediated tRNA processing.

Protein-based RNase Ps in all domains of life

tRNAs mediate the translation of genetic information from nucleotide sequences into protein molecules, and thus play a fundamental role in gene expression [1–3]. They are transcribed as precursors from their respective genes, and must undergo several maturation steps before they are functionally competent [4–6]. In almost all organisms, this entails the endonucleolytic removal of a leader sequence at the 5' end of the tRNA. This processing step is catalysed by a class of enzymes called RNase P [6–8].

The first RNase P enzymes discovered were ribozymes, consisting of a catalytic RNA and a varying number of accessory proteins [9–13]. Such **ribonucleoprotein**-based RNase Ps (**RNP** RNase Ps; see [Glossary](#)) are found in all three domains of life and were long presumed to be the only form of RNase P [12,14]. This was challenged in 1988, when Peter Gegenheimer and colleagues showed that plant chloroplast RNase P exhibits protein-like behaviour and is insensitive to micrococcal nuclease, thus suggesting a protein-based RNase P enzyme devoid of any RNA component [15,16]. Subsequent characterisations of RNase P preparations from mammalian and trypanosomatid mitochondria suggested similar protein-catalysed RNase P activity in these organelles [17–19].

These observations were initially met with scepticism [20], and it was not until 2008 that the existence of an entirely PRORP was unequivocally demonstrated by Walter Rossmann and colleagues. His group showed that human mitochondrial RNase P (mtRNase P) is a multisubunit protein complex, identified its constituent subunits, and demonstrated protein-catalysed RNase P activity of this complex both *in vitro* and *in vivo* [21]. Homologues of the **endonuclease** subunit of mtRNase P were subsequently identified in four out of five eukaryotic supergroups [22]. In contrast to mtRNase P, these homologues outside the metazoan lineage appear to be single-subunit enzymes that are catalytically active without the requirement of additional proteins [23,24]. In addition to the eukaryotic PRORP enzymes, a second form of PRORP was recently discovered in *Aquifex aeolicus*

Highlights

Ribonuclease P (RNase P) enzymes catalyse endonucleolytic processing of tRNA 5' ends.

Long thought to be universally conserved as ribozymes, recent studies have identified two distinct types of protein-only RNase P enzymes across all domains of life.

Structural studies on prototypic members of all types of protein-only RNase P reveal a surprising diversity of molecular architectures.

Structures of tRNA-bound protein-only RNase Ps reveal substantial differences in their substrate recognition mechanisms.

Despite their differences, comparisons suggest that all protein-only RNase P enzymes use a similar catalytic mechanism for tRNA 5' processing.

¹Department of Cellular Biochemistry, University Medical Center Goettingen, Humboldtallee 23, D-37073 Goettingen, Germany

²Research Group Structure and Function of Molecular Machines, Max Planck Institute for Multidisciplinary Sciences, Am Fassberg 11, D-37077 Goettingen, Germany

³Cluster of Excellence Multiscale Bioimaging: from Molecular Machines to Networks of Excitable Cells (MBExC), University of Goettingen, D-37075 Goettingen, Germany

*Correspondence: hauke.hillen@med.uni-goettingen.de (H.S. Hillen).
©Twitter: @hauke_hillen



(*aq*RNase P), a bacterium that possesses neither an RNP RNase P nor a eukaryotic-type protein-only RNase P [25]. Instead, it uses a minimal prokaryotic form of protein-only RNase P that is comprised only of a single **metallonuclease** domain. Homologues of *A. aeolicus* RNase P (HARPs) were also identified in some bacteria and many archaea [25], indicating that this type of protein-only RNase P is widespread among the prokaryotic and archaeal lineages. Taken together, these discoveries establish the existence of two distinct archetypes of protein-only RNase Ps, eukaryotic PRORPs and prokaryotic HARPs, spanning across all domains of life.

The mechanism of ribozyme-catalysed tRNA 5'-processing has been elucidated in detail through structural and functional studies [26–38]. By contrast, the molecular basis of RNA processing by protein-only RNase P enzymes long remained speculative. Over the past two decades, a large body of biochemical and biophysical studies have provided insights into the mechanisms of substrate recognition and catalysis by protein-only RNase P enzymes. Recently, high-resolution structures of members of each of the different types of protein-only RNase Ps have been reported (Table 1). From the combination of functional and structural studies, a molecular picture of tRNA processing by PRORPs has begun to emerge. In this review, we summarise our current understanding of protein-catalysed tRNA 5'-processing and highlight both similarities and differences between the different types of protein-only RNase P enzymes.

Molecular architectures of protein-only RNase P enzymes

Eukaryotic protein-only RNase Ps

Eukaryotic PRORPs all share a characteristic tripartite domain architecture comprised of an N-terminal **pentatricopeptide repeat (PPR)** domain, a split zinc-binding domain (ZBD), and a C-terminal PiIT N terminus (PIN)-like nuclease domain [22,39,40]. While most PRORPs are single-subunit enzymes (ssPRORPs), the PRORP enzymes in metazoan mitochondria are multisubunit protein complexes (msPRORPs).

Single-subunit protein-only RNase Ps

The most well-studied ssPRORPs are those of *Arabidopsis thaliana* and *Trypanosoma brucei*. Both organisms do not encode an RNP RNase P and instead rely exclusively on ssPRORP enzymes for 5'-processing [22–24,41]. Nuclear and organellar tRNA processing are carried out

Table 1. Structures of protein-only RNase P enzymes

Protein or complex	RNA	PDB code	Refs
<i>Arabidopsis thaliana</i> PRORP1	–	4G24	[39]
<i>A. thaliana</i> PRORP2	–	5DIZ	[42]
<i>A. thaliana</i> PRORP2	–	5FT9	[43]
<i>Homo sapiens</i> PRORP - free	–	4ROU	[57]
<i>H. sapiens</i> PRORP - free	–	4XGL	[53]
<i>A. thaliana</i> PRORP1 PPR domain in complex with tRNA	Yeast tRNA ^{phe}	6LVR	[49]
<i>Aquifex aeolicus</i> RNase P	–	7F3E	[61]
<i>Halorhodospira halophila</i> RNase P	–	7OG5	[62]
<i>Thermococcus celer</i> RNase P	–	7E8J	[65]
<i>Planctomycetes bacterium</i> RNase P	–	7E8K	[65]
<i>Planctomycetes bacterium</i> RNase P in complex with tRNA	<i>Escherichia coli</i> tRNA ^{his}	7E8O	[65]
<i>H. sapiens</i> mtRNase P in complex with tRNA	<i>H. sapiens</i> mt-tRNA ^{tyr}	7ONU	[59]

Glossary

Acceptor arm: a structural element of tRNA formed by base pairing of nucleotides at positions 1–7 and 66–72 in a tRNA structure.

Anticodon arm: a stem-loop structure formed by the nucleotides at positions 27–43 in a tRNA structure.

Anticodon loop: a 7-nt long non-base-paired loop between nucleotide positions 32 and 38 in a canonical tRNA structure. Positions 34–36 in the anticodon loop correspond to the anticodon triplet.

D arm: a stem-loop structure formed by the nucleotides at positions 10–25 in a tRNA structure.

Dehydrogenase: an oxidoreductase enzyme catalysing the removal of hydrogen from its substrate.

Dodecameric assembly: a macromolecular complex comprising of 12 subunits (monomers).

Endonuclease: a hydrolase enzyme catalysing hydrolysis of a phosphodiester bond within a polynucleotide chain (i.e., not involving a terminal nucleotide).

Exonuclease: a hydrolase enzyme catalysing the hydrolysis of a terminal phosphodiester bond in a polynucleotide chain.

Metallonuclease: a hydrolase enzyme catalysing exo- or endonucleolytic cleavage of polynucleotides, which require binding of metal ions in the active site for catalytic activity.

Methyltransferase: a transferase enzyme catalysing the transfer of a methyl (-CH₃) group from a cofactor to a substrate.

Pentatricopeptide repeat (PPR): a 35-amino-acid long motif, adopting a helix–turn–helix structure. Typically, multiple PPR repeats are present in tandem and are involved in RNA binding.

PPR code: molecular principle that has been proposed for the modular, sequence-specific recognition of RNA bases by plant PPR proteins [67,68].

Ribonucleoprotein (RNP): a macromolecular complex comprised of RNA and protein subunits.

T arm: a stem-loop structure formed by the nucleotides at positions 49–65 in a tRNA structure.

tRNA elbow: a tRNA tertiary structural element formed by stacking and hydrogen bonding interactions between T and D loops of tRNA.

by distinct isoforms of ssPRORP, which share a conserved domain organisation and up to 30% sequence similarity between the two organisms and with human PRORP [23,24,41]. *A. thaliana* and *T. brucei* PRORPs are catalytically active without the requirement of any additional subunits *in vitro* and *in vivo*, and are therefore *bona fide* ssPRORPs [23,24,41].

The molecular architecture of ssPRORPs has been elucidated through crystal structures of two *A. thaliana* PRORP isoforms, PRORP1 and PRORP2 (*atPRORP1* and *atPRORP2*) (Figure 1A,B) [39,42,43]. These structures reveal a V-shaped arrangement of the three domains, in which the PPR and nuclease domains form the two arms of the 'V', respectively, and the ZBD forms the central vertex. The nuclease domain adopts a PIN/NYN domain-like fold with structural resemblance to the nuclease domains of DNA polymerase I and FLAP nucleases [39,40,44,45]. Its active site contains four catalytic aspartate residues that are highly conserved among all PRORPs [22,39,42]. The ZBD is comprised of a four-stranded antiparallel β sheet, which contains three invariant cysteines and one histidine involved in coordination of a Zn^{2+} ion [39]. The PPR domain is composed of 11 α helices, which together constitute five complete PPR repeats as well as one incomplete repeat. These repeats form a curved structure with an inner concave surface that faces the active site of the nuclease domain. Since many PPR proteins are RNA-binding proteins, the PPR domain was proposed to be involved in substrate binding [39,43,46–48]. This was recently confirmed by a structure of the *atPRORP1* PPR domain in complex with tRNA, which shows that the PPR domain forms specific contacts with the RNA [49]. Comparison with the previous substrate-free structures indicates that the V-shaped arrangement of PPR and nuclease domains would need to widen to




Figure 1. Structures of eukaryotic protein-only ribonuclease (RNase) Ps. (A) Structure of *Arabidopsis thaliana* proteinaceous RNase P PRORP1 (*atPRORP1*) (PDB: 4G24) [39]. A schematic domain representation is shown above the structure with individual domains coloured in shades of green. Protein regions modelled in the structure are indicated by a black line above the schematic. (B) Structure of *A. thaliana* PRORP2 (*atPRORP2*) (PDB: 5DIZ) [42]. Domain representation and colouring as in (A). (C) Structure of substrate-bound human mitochondrial RNase P complex (PDB: 7ONU) [59]. A schematic domain representation of the nuclease subunit *Homo sapiens* proteinaceous RNase P (*hsPRORP*) is shown above the structure with the same colouring as in (A). The additional subunits TRMT10C and SDR5C1 are coloured in shades of blue and grey, respectively. The tRNA precursor is shown in red. The longer extensions in *atPRORP1* and *hsPRORP1* in (A) and (C) correspond to their respective organellar targeting sequences. Abbreviation: PPR, pentatricopeptide repeat.

accommodate the tRNA between the two domains [49]. This is supported by small-angle X-ray scattering (SAXS) studies of full-length *at*PRORP in complex with tRNA and *in silico* normal mode analysis, which suggest that *at*PRORPs possess conformational flexibility [42,43,48]. Thus, ssPRORPs may undergo rearrangements during substrate binding.

In summary, the structures of *at*PRORP enzymes reveal the conserved molecular architecture of ssPRORPs and provide insights into their mechanism of substrate binding.

Metazoan multisubunit protein-only RNase P

In contrast to the widespread ssPRORPs, msPRORPs have thus far only been found in mitochondria of metazoans [21,50]. In humans, the mtRNase P is a trimeric complex comprised of an endonuclease, a **methyltransferase**, and a fatty acid/sterol **dehydrogenase** [21,51,52]. Its nucleolytic subunit *Homo sapiens* PRORP (*hs*PRORP; also known as MRPP3) is an orthologue of ssPRORPs and exhibits a high degree of sequence similarity to the ssPRORPs from *A. thaliana* and *T. brucei* [23,53]. In contrast to these, however, *hs*PRORP requires the two additional mtRNase P subunits, TRMT10C and SDR5C1, for tRNA processing [21]. TRMT10C (tRNA methyltransferase 10 type C; also known as MRPP1) is a SPOUT-fold methyltransferase which is responsible for methylating mitochondrial tRNAs, while SDR5C1 (short-chain dehydrogenase/reductase family 5C, member 1; also known as MRPP2 or HSD17B10) is a dehydrogenase involved in β oxidation of a diverse array of substrates [51,52,54]. TRMT10C and SDR5C1 interact with each other independent of *hs*PRORP to form a complex that can bind and methylate the pre-tRNA [52,55,56].

The first structural insights into human mtRNase P came from two crystal structures of *hs*PRORP, which reveal a V-shaped organisation resembling ssPRORPs (Figure 2A) [53,57]. In both structures, the active site of the nuclease domain is partially disordered, and catalytic residues are engaged in intra-molecular interactions. While these interactions differ between the two structures, the authors of both studies concluded that *hs*PRORP adopts an autoinhibited conformation in isolation. However, both studies used truncated variants of *hs*PRORP lacking parts of the PPR domain that are inactive in biochemical assays even in the presence of TRMT10C and SDR5C1 [53,57], which has raised concerns about the physiological relevance of the observed states




Figure 2. Structural comparison between free and complex-bound human proteinaceous ribonuclease P (PRORP). Structures of substrate-free *Homo sapiens* PRORP (*hs*PRORP) (A) (PDB: 4XGL) [53] and the pre-tRNA-bound mitochondrial ribonuclease P (mtRNase P) complex (B) (PDB: 7ONU) [59], coloured as in Figure 1C. The nuclease domain of substrate-free *hs*PRORP needs to rotate outwards by $\sim 25^\circ$ (pivot axis, perpendicular to the image plane, is shown as black circle) in order to accommodate pre-tRNA between its pentatricopeptide repeat (PPR) and nuclease domains and correctly position the substrate in its active site, as seen in the mtRNase P complex.

[8,58]. Whether free full-length *hs*PRORP also adopts an autoinhibited conformation remains to be determined experimentally. Taken together, these structures confirmed *hs*PRORPs' similarity to plant *ss*PRORPs, and suggested that it may adopt a catalytically incompetent conformation prior to formation of the substrate-engaged mtRNase P complex.


Recently, the cryoelectron microscopy (cryo-EM) structure of the complete human mtRNase P complex bound to pre-tRNA was reported (Figure 1C) [59]. This structure reveals how *hs*PRORP, TRMT10C, and SDR5C1 interact with each other and with the pre-tRNA substrate. SDR5C1 forms a homo-tetramer, which interacts with TRMT10C to form a platform to which the tRNA substrate binds through multiple interactions. This positions the methyl acceptor, a purine base in position 9 of the tRNA, in the methyltransferase domain of TRMT10C. *Hs*PRORP binds atop this complex, with the scissile phosphodiester bond at the 5' end of the tRNA positioned in the nuclease active site. Compared with the previous structures, *hs*PRORP adopts a more open conformation in which the autoinhibitory interactions are abolished and the active site organisation resembles that observed in *at*PRORP1 [59]. This conformation appears to be stabilised by interactions with TRMT10C and the tRNA, suggesting that *hs*PRORP may become activated upon recruitment to the complex (Figure 2B) [59]. The structure of mtRNase P thus reveals the active conformation of *hs*PRORP and shows how this multifunctional complex integrates two enzymatic activities into one molecular machinery.

Prokaryotic minimal protein-only RNase Ps

The most recent type of PRORP enzymes discovered are the prokaryotic protein-only RNase Ps, also known as HARPs. While HARPs appear to have functionally replaced RNP RNase Ps in some bacteria like *A. aeolicus*, they coexist alongside the RNP RNase Ps in many archaea [25,60]. HARPs represent the most minimal PRORP enzymes known to date, as they are comprised solely of a metallo-nuclease domain that belongs to the same PIN domain-like superfamily as the nuclease domains of eukaryotic PRORPs [25,40,45].

Recent cryo-EM studies of HARPs from *A. aeolicus* and *Halorhodospira halophila* demonstrate that they adopt a different architecture than eukaryotic PRORPs [61,62]. They assemble into large **dodecameric assemblies** comprised of hexamers of dimers, wherein two monomers dimerise via their spike helix or protruding helix (SH/PrH) microdomains, and the resulting dimers interact side by side to form a superhelical assembly (Figure 3) [61,62]. The overall fold of the individual monomers is distinct from the nuclease domain of eukaryotic PRORPs, and instead shows similarities to different members of the PIN family, for example, the VapC4 toxin from *Pyrococcus horikoshii* [40,60,61,63,64]. This suggests that HARPs may have an independent evolutionary origin from eukaryotic PRORPs [61,62]. Biochemical experiments show that the oligomerisation of HARPs is necessary for their RNase P activity [62], and it has been proposed that two adjacent dimers may cooperate to mediate processing of a single tRNA molecule [61,62]. In this case, a tetrameric assembly formed by the association of two dimers would represent the minimal catalytic unit of HARPs [61,62]. This hypothesis is supported by a recently reported crystal structure of *Planctomycetes bacterium* HARP (*pb*HARP) in complex with pre-tRNA [65]. As dodecameric assemblies, HARPs could contain up to ten substrate-binding sites, which may allow simultaneous processing of multiple tRNAs [61,62,65]. However, this model remains speculative, as a structure of substrate-engaged dodecameric HARP assembly has not yet been determined.

In summary, these structural studies have elucidated the molecular architecture of prototypic members of the different forms of protein-only RNase Ps. In conjunction with a large body of previous biochemical and biophysical data, this now enables a comparison of their molecular mechanisms of substrate recognition, specificity, and cleavage.



Trends in Biochemical Sciences

Figure 3. Structures of prokaryotic minimal protein-only ribonuclease (RNase) Ps. (A) Structure of a dimeric assembly of protein-only RNase P from *Aquifex aeolicus* (aaRNase P) (PDB: 7F3E) [61]. The domain composition of aaRNase P is shown above the structure and coloured to represent monomer 1. Modelled regions are indicated with a black line above the domain representation. (B) Structure of a dimeric assembly of protein-only RNase P from *Halorhodospira halophila* (hhRNase P) (PDB: 7OG5) [62]. Domain representation as in (A). (C,D) Structures of dodecameric assemblies of protein-only RNase Ps from *A. aeolicus* (C) (PDB: 7F3E) and *H. halophila* (D) (PDB: 7OG5). The individual dimeric assemblies, as shown in (A) and (B), are indicated.

Substrate recognition and specificity

The PRORP enzymes generally interact with the same elements in tRNAs as RNP RNase Ps, but utilise distinct molecular binding principles [48,49,59,61,62,65]. Structures of RNP RNase Ps show that these enzymes recognise their substrates through specific interactions with the

acceptor arm and the **elbow** of the tRNA, as well as the 5' and 3' extensions [29,32,34]. In particular, they recognise a conserved base pair (G/U19-C56) between the **T arm** and the **D arm**, which together form the elbow, through RNA–RNA stacking interactions (Figure 4A,B).

Substrate binding by single-subunit protein-only RNase Ps

The mechanism of substrate binding by ssPRORPs was elucidated through a combination of biochemical, biophysical, and structural studies. This revealed that the PPR domain interacts with the tRNA elbow in a conceptually similar manner as in RNP RNase Ps, mediated through two mechanisms (Figure 4C,D) [43,48,49]. First, it forms a positively charged binding pocket which accommodates the tRNA elbow and stabilises the phosphate backbone. Second, it forms base-specific interactions with the tRNA elbow. The structure of the *at*PRORP1 PPR domain in complex with yeast tRNA^{Phe} shows that residues Y133, Y140, and R210 interact with D/U17, the G18- Ψ /U55 base pair, and the G/U19-C56 base pair of the tRNA, respectively [49]. This is consistent with previous SAXS data and biochemical data, which show that substitution of these protein residues or bases leads to decreased substrate affinity and cleavage efficiency [43,47,48]. While no substrate-bound structures are available for other ssPRORPs, biochemical




Figure 4. Substrate recognition in ribonuclease (RNase) P enzymes. (A,C,E,G) Schematic representations of tRNAs from different organisms and cellular compartments, indicating the conservation of their secondary structure elements: (A) human nucleus, (C) *Arabidopsis thaliana* mitochondria, (E) *Planctomycetes bacterium*, and (G) human mitochondria [92,93]. Transparent grey circles represent individual nucleotides. Blue or purple circles indicate conserved nucleotides. G/U19-C56 interactions are coloured in purple. Acceptor arm, D arm, anticodon arm, variable region, and T arm are labelled Acc, D, AC, V, and T, respectively. Strong overlap and low transparency of circles indicates conservation, while weak overlap and high transparency indicates variability. Regions interacting with human nuclear ribonucleoprotein (RNP) RNase P, *A. thaliana* proteinaceous RNase P (*at*PRORP1), *Homo sapiens* PRORP (*hs*PRORP), and *hs*TRMT10C are marked by brown, green, green, and blue arcs, respectively. (B) Interaction of human nuclear RNP RNase P with tRNA (PDB: 6AHU) [32]. The catalytic RNA component is coloured in brown, and protein subunits are coloured in grey. Interactions with the tRNA elbow are shown in inset. Acceptor arm, D arm, anticodon arm, variable region, and T arm are coloured red, orange, maroon, light-brown, and magenta, respectively. Hydrogen bonds are shown as broken lines throughout. (D) Interaction of *at*PRORP1 pentatricopeptide repeat (PPR) domain with the tRNA elbow (PDB: 6LVR) [49]. The PPR domain is coloured in green. Interactions with the tRNA elbow are shown in inset. (F) Interaction of *Planctomycetes bacterium* RNase P with tRNA (PDB: 7E8O) [65]. The nucleotide at position 56 is a G in the tRNA substrate used for this study, and G/U19-C56 interactions are not present. (H) Interaction of human mitochondrial RNase P (mtRNase P) complex with pre-tRNA (PDB: 7ONU) [59]. Colouring as in Figure 1C. Interactions with the tRNA elbow and the anticodon loop are shown in inset.

data indicate that *at*PRORP2 and *at*PRORP3 also form base-specific interactions between the tRNA elbow and the PPR domain [47,66]. Thus, although the precise interactions may differ, specific recognition of tRNA elbow bases appears to be a conserved feature among ssPRORPs. Notably, these interactions differ from the previously proposed **PPR code** derived from plant organellar PPR proteins, which bind RNAs in a one-base-per-repeat fashion [46,47,49,67–69]. Thus, PPR proteins appear to have diverged substantially with respect to their RNA binding mechanisms. Taken together, these observations suggest that ssPRORPs have adapted their PPR domain to recognise similar structural elements in tRNAs as the RNP RNase Ps.

Substrate binding by mitochondrial multisubunit protein-only RNase Ps

By contrast, the mechanism of substrate recognition by the msPRORPs in metazoan mitochondria appears to differ. The structure of human mtRNase P shows that the PPR domain of *hs*PRORP forms a similar charge-complementary binding groove for the tRNA elbow as in ssPRORPs, but it does not form base-specific interactions (Figure 4G,H) [59]. Instead, the additional subunits TRMT10C and SDR5C1 interact with the pre-tRNA by two different mechanisms. First, the TRMT10C/SDR5C1 complex forms a shape- and charge-complementary binding platform that interacts with all four arms of the tRNA via nonspecific backbone stabilisation. Second, TRMT10C additionally interacts with the **anticodon loop** through a specific interaction between its residue R181 and a conserved base in position 33 of the tRNA (Figure 4G,H) [59]. Thus, the additional subunits in metazoan mtRNase Ps enable a unique substrate recognition mechanism that involves interactions with tRNA elements that are not bound by any other known RNase P.

A possible explanation for the recognition of distinct elements by the mitochondrial msPRORPs may lie in the structural properties of their substrates. Metazoan mitochondrial tRNAs exhibit a large degree of variability in otherwise conserved tRNA elements [70–72] (Figure 4A,C,E,G). In particular, the sequences, lengths, and structures of their T and D arms are highly degenerate (Figure 4G). The canonical substrate recognition mechanism used by RNP RNase P and ssPRORP enzymes, which involves specific interactions with the elbow region, may thus not be feasible or sufficient for the binding of all metazoan mitochondrial tRNAs. By contrast, the base in the anticodon loop that interacts with TRMT10C is conserved as pyrimidine in mitochondrial tRNAs, and thus offers an alternative specific recognition element. This suggests that the additional subunits in metazoan mitochondrial msPRORPs may have been adopted to compensate for the structural variability in their tRNA substrates [8,21,59].

Substrate binding by prokaryotic minimal protein-only RNase Ps

Based on structural modelling, it was suggested that HARPs may recognise similar tRNA elements as ssPRORPs [61,62]. In particular, it was proposed that the SH/PrH microdomain may interact with the elbow. This was recently confirmed by the crystal structure of *pb*HARP in complex with pre-tRNA, which shows that a positively charged patch formed by conserved arginine residues (R116, R123, R138, and R142 in *pb*HARP) interacts with the tRNA elbow via backbone interactions [65]. Mutation of these residues to alanine results in decreased activity of both *pb*HARP and *hh*HARP [62,65]. In contrast to ssPRORPs, however, no base-specific interactions between the tRNA elbow and HARPs were observed (Figure 4E,F) [65]. It has been proposed that the binding of the elbow to the SH/PrH microdomain of one monomer of a dimer may position the 5' end of the tRNA in the active site of a monomer in the adjacent dimer [61,62]. The distance between these two binding sites could then act as a molecular ruler to define the 5' cleavage site, as has been analogously proposed for RNP RNase Ps and ssPRORPs [29,34,47]. However, this model remains speculative to date, as the 5' cleavage site was not positioned in an active site in the crystal structure of *pb*HARP in complex with pre-tRNA [65]. Taken together, the available

data suggest that HARPs interact with their substrates in a conceptually similar fashion as RNP RNase Ps and ssPRORPs, but may rely less on base-specific interactions with the tRNA elbow. However, further work is required to fully elucidate their molecular mechanism.

Substrate specificity of protein-only RNase P enzymes


The observed differences in substrate recognition strategies by the different forms of RNase Ps may also partially explain their substrate specificities. The multisubunit mtRNase P, which makes interactions with the all four tRNA arms, cannot process mt-tRNA^{Ser(AGY)}, a mitochondrial tRNA lacking the entire D arm [73,74]. By contrast, many other RNase P enzymes can cleave non-tRNA substrates with structural similarities to tRNAs. For example, structured RNAs such as the 4.5S RNA and tmRNAs in *Escherichia coli*, or MALAT1 and MEN-β/NEAT1 in humans, have been described as substrates for RNP RNase P [75–77]. Several observations indicate that ssPRORPs and HARPs may also have a broad substrate spectrum. First, both are able to rescue the otherwise lethal knockout of endogenous RNP RNase P in yeast or *E. coli* [23,25,78,79]. Second, the ssPRORP isoform that localises to plant mitochondria has been shown to process t elements, noncanonical substrates encoded in the mitochondrial genome, both *in vivo* and *in vitro* [41]. Third, artificial substrates resembling tRNAs with truncated D arm or **anticodon arm** can be processed by *A. thaliana* PRORPs *in vitro* [47,80]. While processing of non-tRNA substrates by HARPs has not been demonstrated, their conceptually similar substrate binding mechanism suggests that they may have a similarly broad substrate spectrum as ssPRORPs.

Taken together, these structural and functional studies show that all types of protein-only RNase Ps interact with the tRNA elbow and the 5' cleavage site. However, their detailed interaction strategies differ, and metazoan mtRNase Ps additionally interact with several other structural elements of tRNAs.

Mechanism of catalysis by protein-only RNase Ps

Despite their distinct structures and substrate recognition modes, all protein-based RNase Ps appear to use a common catalytic mechanism. In particular, they all share a conserved active site architecture. In *at*PRORP1 and *hs*PRORP, the active sites are comprised of four highly conserved aspartate residues, which coordinate catalytic metal ions near the cleavage site (Figure 5A,B) [39,59]. Three out of these four aspartates – D399, D475, and D493 in *at*PRORP1; D409, D479, and D499 in *hs*PRORP – are also strictly conserved among HARPs and adopt a similar organisation to coordinate two catalytic metal ions (Figure 5C) [61,62,65]. The fourth aspartate (D474 in *at*PRORP1, D478 in *hs*PRORP) is replaced by a glutamate residue in *hh*HARP and *aq*RNase P, but is not conserved in HARPs from other organisms, including *pb*HARP, and may thus not be essential for catalysis. Instead, HARP active sites contain a different conserved aspartate residue at position 138 in *aq*RNase P (D139 in *hh*HARP and D151 in *pb*HARP), which is absent in PRORPs. While these differences may reflect the distinct evolutionary origins of HARPs and PRORPs, the otherwise remarkable similarity in the active site organisation indicates conservation of the catalytic mechanism [25,61,62,65].

From biochemical and structural data on both single-subunit and multisubunit PRORPs, a mechanistic model for catalysis by protein-only RNase Ps has emerged [39,59,81–83]. In this model, the four active site aspartates coordinate two Mg²⁺ ions, which in turn coordinate the pro-S_p oxygen and the 3' proximal phosphoester-bonded oxygen of the scissile phosphate in the RNA substrate. This is supported by the structure of substrate-bound human mtRNase P, which reveals the architecture of the *hs*PRORP active site in a pre-catalytic state with the RNA positioned for cleavage (Figure 5D) [59]. Its arrangement is reminiscent of the active site configurations of



Trends in Biochemical Sciences

Figure 5. Catalytic mechanism of protein-only ribonuclease (RNase) P. (A) Active site structure of *Arabidopsis thaliana* proteinaceous RNase P (atPRORP1) (PDB: 4G24) [39]. AtPRORP1 is coloured green. Two metal (Mn^{2+}) ions bound in the active site are shown as purple spheres. Active sites shown in (B–E) are superimposed on (A). Conserved catalytic aspartates and residues within 5 Å of the two metal ions are shown as sticks throughout. (B) Active site structure of human PRORP in the mitochondrial RNase P (mtRNase P) complex [59]. *Homo sapiens* proteinaceous RNase P (hsPRORP) is coloured green, and one metal ion observed in the active site is marked in purple. The RNA is omitted for clarity. (C) Active site structure of prokaryotic protein-only RNase Ps from *Planctomycetes bacterium* (pbPRORP) [65]. *PbRNase P* is coloured in dark and light magenta, respectively. The two metal (Ca^{2+}) ions bound in the active site are shown as purple spheres. (D) Substrate-engaged active site of human PRORP in the mtRNase P complex (PDB: 7ONU) [59]. HsPRORP is coloured as in (B), while pre-tRNA is coloured dark red. Coordination interactions between metal 2 and catalytic aspartates or the scissile phosphate are shown as broken yellow lines. (E) Substrate-engaged active site structure of human exonuclease 1 (hsExo1) (PDB: 5V08) [85]. HsExo1 is coloured blue, while substrate DNA is coloured orange. Ordered water molecules in the active site are shown as red spheres. Coordinating interactions are shown as broken yellow lines, and hydrogen bonds are shown as thin broken grey lines. The active site organisation and substrate conformation in hsExo1 is similar to hsPRORP in the mtRNase P complex (D). Abbreviation: HARP, homologue of *Aquifex aeolicus* RNase P.

human **exonuclease 1** (hsExo1) (Figure 5E) and the Klenow fragment, suggesting that protein-only RNase Ps employ a similar two-ion mechanism as these exonucleases, as previously proposed [39,83–87]. The metal ion at site 2 (M2), together with a proximal aspartate (D499 in hsPRORP), may position an activated water molecule opposite of the scissile P–O bond for nucleophilic attack. The metal ion at site 1 (M1) may stabilise the transition state by coordinating the oxygen atom of the scissile P–O bond. The recent crystal structure pbHARP reveals a similar configuration of catalytic residues and coordination of two metal ions in the active site, suggesting that HARPs likely share the two-ion catalytic mechanism [65]. Notably, RNP RNase Ps have also been proposed to use a two-metal catalytic mechanism, which conceptually resembles that described above [29,35,88–91]. Taken together, the structures of PRORPs and HARPs indicate

that protein-only RNase P enzymes use an ancient catalytic mechanism employed by many functionally disparate nucleases.

Concluding remarks

PRORP enzymes represent a striking example of convergent evolution, as they carry out an identical reaction as their ribozyme-based counterparts but are evolutionarily unrelated. Since their discovery several decades ago, a combination of structural and functional studies has resulted in a basic molecular understanding of protein-only RNase P enzymes. These studies have revealed a surprising diversity of molecular architectures among these enzymes, as well as a conceptually unique substrate recognition and binding mode used by msPRORPs. Despite these differences, however, their catalytic mechanism appears to be highly conserved. Nevertheless, a high-resolution structure with substrate bound in the active site is thus far only available for the eukaryotic mitochondrial msPRORP [59]. As a consequence, many mechanistic assumptions for ssPRORPs and HARPs are based on extensive biochemical data as well as structural and evolutionary comparison. Therefore, the next important step will be to obtain molecular snapshots of ssPRORPs and HARPs in complex with their respective substrates in the active site in order to verify these hypotheses. Furthermore, the biological functions of individual PRORP enzymes will need to be further investigated. The now available structural and mechanistic data will provide a framework for future studies aimed at answering the many open questions (see [Outstanding questions](#)) about protein-only RNase Ps.

Acknowledgments

We apologise to those colleagues whose work we could not cite due to the limited length of this review. We wish to thank the members of the Hillen laboratory for valuable comments and discussions. H.S.H. was supported by the Deutsche Forschungsgemeinschaft (FOR2848, SFB1190, EXC 2067/1- 390729940). The study was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC 2067/1-390729940.

Declaration of interests

The authors declare no conflicts of interest.

References

- Crick, F.H.C. (1968) The origin of the genetic code. *J. Mol. Biol.* 38, 367–379
- Hoagland, M. (1996) Biochemistry or molecular biology? The discovery of 'soluble RNA'. *Trends Biochem. Sci.* 21, 77–80
- Hoagland, M.B. *et al.* (1958) A soluble ribonucleic acid intermediate in protein synthesis. *J. Biol. Chem.* 231, 241–257
- Wolin, S.L. and Matera, A.G. (1999) The trials and travels of tRNA. *Genes Dev.* 13, 1–10
- Hopper, A.K. and Phizicky, E.M. (2003) tRNA transfers to the limelight. *Genes Dev.* 17, 162–180
- Phizicky, E.M. and Hopper, A.K. (2010) tRNA biology charges to the front. *Genes Dev.* 24, 1832–1860
- Kazantsev, A.V. and Pace, N.R. (2006) Bacterial RNase P: a new view of an ancient enzyme. *Nat. Rev. Microbiol.* 4, 729–740
- Schencking, I. *et al.* (2020) Diversity and evolution of RNase P. In *Evolutionary Biology—A Transdisciplinary Approach* (Pontarotti, P. *et al.*, eds), pp. 255–299. Springer
- Robertson, H.D. *et al.* (1972) Purification and properties of a specific *Escherichia coli* ribonuclease which cleaves a tyrosine transfer ribonucleic acid precursor. *J. Biol. Chem.* 247, 5243–5251
- Esakova, O. and Krasilnikov, A.S. (2010) Of proteins and RNA: the RNase P/MRP family. *RNA* 16, 1725–1747
- Frank, D.N. and Pace, N.R. (1998) Ribonuclease P: unity and diversity in a tRNA processing ribozyme. *Annu. Rev. Biochem.* 67, 153–180
- Walker, S.C. and Engelke, D.R. (2008) Ribonuclease P: the evolution of an ancient RNA enzyme. *Crit. Rev. Biochem. Mol. Biol.* 41, 77–102
- Phan, H.D. *et al.* (2021) The many faces of RNA-based RNase P, an RNA-world relic. *Trends Biochem. Sci.* 46, 976–991
- Altman, S. (2007) A view of RNase P. *Mol. Biosyst.* 3, 604–607
- Wang, M.J. *et al.* (1988) Novel mechanisms for maturation of chloroplast transfer RNA precursors. *EMBO J.* 7, 1567–1574
- Thomas, B.C. *et al.* (1995) Spinach chloroplast RNase P: a putative protein enzyme. *Nucleic Acids Symp. Ser.* 33, 95–98
- Rossmannith, W. *et al.* (1995) Human mitochondrial tRNA processing. *J. Biol. Chem.* 270, 12885–12891
- Rossmannith, W. and Karwan, R.M. (1998) Characterization of human mitochondrial RNase P: novel aspects in tRNA processing. *Biochem. Biophys. Res. Commun.* 247, 234–241
- Salavati, R. *et al.* (2001) Mitochondrial ribonuclease P activity of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* 115, 109–117
- Pinker, F. *et al.* (2013) PPR proteins shed a new light on RNase P biology. *RNA Biol.* 10, 1457–1468
- Holzmann, J. *et al.* (2008) RNase P without RNA: identification and functional reconstitution of the human mitochondrial tRNA processing enzyme. *Cell* 135, 462–474
- Lechner, M. *et al.* (2015) Distribution of ribonucleoprotein and protein-only RNase P in Eukarya. *Mol. Biol. Evol.* 32, 3186–3193
- Gobert, A. *et al.* (2010) A single *Arabidopsis* organellar protein has RNase P activity. *Nat. Struct. Mol. Biol.* 17, 740–744
- Taschner, A. *et al.* (2012) Nuclear RNase P of *Trypanosoma brucei*: a single protein in place of the multicomponent RNA-protein complex. *Cell Rep.* 2, 19–25
- Nickel, A.I. *et al.* (2017) Minimal and RNA-free RNase P in *Aquifex aeolicus*. *Proc. Natl. Acad. Sci. U. S. A.* 114, 11121–11126

Outstanding questions

What is the atomic structure of full-length substrate-bound ssPRORPs?

What is the architecture of a dodecameric, substrate-bound HARP complex?

What are the *in vivo* substrates of protein-only RNase Ps, especially in organisms that contain both RNP- and protein-only RNase Ps?

Do additional, yet undiscovered types of protein-only RNase P enzymes exist?

How are protein-only RNase Ps regulated?

26. Steitz, T.A. and Steitz, J.A. (1993) A general two-metal-ion mechanism for catalytic RNA. *Proc. Natl. Acad. Sci. U. S. A.* 90, 6498–6502
27. Kazantsev, A. *et al.* (2003) High-resolution structure of RNase P protein from *Thermotoga maritima*. *Proc. Natl. Acad. Sci. U. S. A.* 100, 7497–7502
28. Kazantsev, A. *et al.* (2011) Solution structure of RNase P RNA. *RNA* 17, 1159–1171
29. Reiter, N.J. *et al.* (2010) Structure of a bacterial ribonuclease P holoenzyme in complex with tRNA. *Nature* 468, 784–789
30. Kirsebom, L.A. (2007) RNase P RNA mediated cleavage: substrate recognition and catalysis. *Biochimie* 89, 1183–1194
31. Jarrous, N. and Gopalan, V. (2010) Archaeal/eukaryal RNase P: subunits, functions and RNA diversification. *Nucleic Acids Res.* 38, 7885–7894
32. Wu, J. *et al.* (2018) Cryo-EM structure of the human ribonuclease P holoenzyme. *Cell* 175, 1393–1404.e11
33. Wan, F. *et al.* (2019) Cryo-electron microscopy structure of an archaeal ribonuclease P holoenzyme. *Nat. Commun.* 10, 1–13
34. Lan, P. *et al.* (2018) Structural insight into precursor tRNA processing by yeast ribonuclease P. *Science* 362, eaat6678
35. Wamecke, J.M. *et al.* (1996) Ribonuclease P (RNase P) RNA is converted to a Cd²⁺-ribozyme by a single Rp-phosphorothioate modification in the precursor tRNA at the RNase P cleavage site. *Proc. Natl. Acad. Sci. U. S. A.* 93, 8924–8928
36. Burgin, A.B. and Pace, N.R. (1990) Mapping the active site of ribonuclease P RNA using a substrate containing a photoaffinity agent. *EMBO J.* 9, 4111–4118
37. Kazakov, S. and Altman, S. (1991) Site-specific cleavage by metal ion cofactors and inhibitors of M1 RNA, the catalytic subunit of RNase P from *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* 88, 9193–9197
38. Guerrier-Takada, C. *et al.* (1989) Specific interactions in RNA enzyme-substrate complexes. *Science* 246, 1578–1584
39. Howard, M.J. *et al.* (2012) Mitochondrial ribonuclease P structure provides insight into the evolution of catalytic strategies for precursor-tRNA 5' processing. *Proc. Natl. Acad. Sci. U. S. A.* 109, 16149–16154
40. Gobert, A. *et al.* (2019) Involvement of PIN-like domain nucleases in tRNA processing and translation regulation. *IUBMB Life* 71, 1117–1125
41. Gutmann, B. *et al.* (2012) PRORP proteins support RNase P activity in both organelles and the nucleus in *Arabidopsis*. *Genes Dev.* 26, 1022–1027
42. Karasik, A. *et al.* (2016) Nuclear protein-only ribonuclease P2 structure and biochemical characterization provide insight into the conserved properties of tRNA 5' end processing enzymes. *J. Mol. Biol.* 428, 26
43. Pinker, F. *et al.* (2017) Biophysical analysis of *Arabidopsis* protein-only RNase P alone and in complex with tRNA provides a refined model of tRNA binding. *J. Biol. Chem.* 292, 13904–13913
44. Anantharaman, V. and Aravind, L. (2006) The NYN domains: novel predicted RNases with a PIN domain-like fold. *RNA Biol.* 3, 18–27
45. Matelska, D. *et al.* (2017) Comprehensive classification of the PIN domain-like superfamily. *Nucleic Acids Res.* 45, 6995–7020
46. Imai, T. *et al.* (2014) Pentatricopeptide repeat motifs in the processing enzyme PRORP1 in *Arabidopsis thaliana* play a crucial role in recognition of nucleotide bases at TψC loop in precursor tRNAs. *Biochem. Biophys. Res. Commun.* 450, 1541–1546
47. Brillante, N. *et al.* (2016) Substrate recognition and cleavage-site selection by a single-subunit protein-only RNase P. *Nucleic Acids Res.* 44, 2323–2336
48. Gobert, A. *et al.* (2013) Structural insights into protein-only RNase P complexed with tRNA. *Nat. Commun.* 4, 1353
49. Teramoto, T. *et al.* (2020) Pentatricopeptide repeats of protein-only RNase P use a distinct mode to recognize conserved bases and structural elements of pre-tRNA. *Nucleic Acids Res.* 48, 11815–11826
50. Sen, A. *et al.* (2016) Loss of the mitochondrial protein-only ribonuclease P complex causes aberrant tRNA processing and lethality in *Drosophila*. *Nucleic Acids Res.* 44, 6409–6422
51. Luo, M.J. *et al.* (1995) Short-chain 3-hydroxy-2-methylacyl-CoA dehydrogenase from rat liver: purification and characterization of a novel enzyme of isoleucine metabolism. *Arch. Biochem. Biophys.* 321, 214–220
52. Vilardo, E. *et al.* (2012) A subcomplex of human mitochondrial RNase P is a bifunctional methyltransferase—extensive moonlighting in mitochondrial tRNA biogenesis. *Nucleic Acids Res.* 40, 11583–11593
53. Reinhard, L. *et al.* (2015) Structure of the nuclease subunit of human mitochondrial RNase P. *Nucleic Acids Res.* 43, 5664–5672
54. Yang, S.Y. *et al.* (2005) Multiple functions of type 10 17β-hydroxysteroid dehydrogenase. *Trends Endocrinol. Metab.* 16, 167–175
55. Oerum, S. *et al.* (2018) Structural insight into the human mitochondrial tRNA purine N1-methyltransferase and ribonuclease P complexes. *J. Biol. Chem.* 293, 12862–12876
56. Vilardo, E. and Rossmannith, W. (2015) Molecular insights into HSD10 disease: impact of SDR5C1 mutations on the human mitochondrial RNase P complex. *Nucleic Acids Res.* 43, 5112–5119
57. Li, F. *et al.* (2015) Auto-inhibitory mechanism of the human mitochondrial RNase P protein complex. *Sci. Rep.* 5, 9878
58. Klemm, B.P. *et al.* (2016) The diversity of ribonuclease P: protein and RNA catalysts with analogous biological functions. *Biomolecules* 6, 27
59. Bhatta, A. *et al.* (2021) Structural basis of RNA processing by human mitochondrial RNase P. *Nat. Struct. Mol. Biol.* 28, 713–723
60. Schwarz, T.S. *et al.* (2019) Homologs of *Aquifex aeolicus* protein-only RNase P are not the major RNase P activities in the archaea *Haloflex volcanii* and *Methanosarcina mazei*. *IUBMB Life* 71, 1109–1116
61. Teramoto, T. *et al.* (2021) Minimal protein-only RNase P structure reveals insights into tRNA precursor recognition and catalysis. *J. Biol. Chem.* 297, 101028
62. Feyh, R. *et al.* (2021) Structure and mechanistic features of the prokaryotic minimal RNase P. *eLife* 10, e70160
63. Hatti, K. *et al.* (2017) Structure determination of contaminant proteins using the MarathonMR procedure. *J. Struct. Biol.* 197, 372–378
64. Daniels, C.J. *et al.* (2019) Both kinds of RNase P in all domains of life: surprises galore. *RNA* 25, 286–291
65. Li, Y. *et al.* (2022) Crystal structures and insights into precursor tRNA 5'-end processing by prokaryotic minimal protein-only RNase P. *Nat. Commun.* 13, 1–13
66. Klemm, B.P. *et al.* (2017) Molecular recognition of pre-tRNA by *Arabidopsis* protein-only ribonuclease P. *RNA* 23, 1860
67. Barkan, A. *et al.* (2012) A combinatorial amino acid code for RNA recognition by pentatricopeptide repeat proteins. *PLoS Genet.* 8, e1002910
68. Yagi, Y. *et al.* (2013) Elucidation of the RNA recognition code for pentatricopeptide repeat proteins involved in organelle RNA editing in plants. *PLoS One* 8, e57286
69. Yagi, Y. *et al.* (2013) Pentatricopeptide repeat proteins involved in plant organellar RNA editing. *RNA Biol.* 10, 1419–1425
70. Suzuki, T. *et al.* (2011) Human mitochondrial tRNAs: biogenesis, function, structural aspects, and diseases. *Annu. Rev. Genet.* 45, 299–329
71. Giegé, R. *et al.* (2012) Structure of transfer RNAs: similarity and variability. *Wiley Interdiscip. Rev. RNA* 3, 37–61
72. Holzmann, J. and Rossmannith, W. (2009) tRNA recognition, processing, and disease: hypotheses around an unorthodox type of RNase P in human mitochondria. *Mitochondrion* 9, 284–288
73. Lopez Sanchez, M.I.G. *et al.* (2011) RNA processing in human mitochondria. *Cell Cycle* 10, 2904–2916
74. Reinhard, L. *et al.* (2017) The MRPP1/MRPP2 complex is a tRNA-maturation platform in human mitochondria. *Nucleic Acids Res.* 45, 12469–12480
75. Peck-Miller, K.A. and Altman, S. (1991) Kinetics of the processing of the precursor to 4.5 S RNA, a naturally occurring substrate for RNase P from *Escherichia coli*. *J. Mol. Biol.* 221, 1–5
76. Wilusz, J.E. *et al.* (2008) 3' end processing of a long nuclear-retained noncoding RNA yields a tRNA-like cytoplasmic RNA. *Cell* 135, 919–932
77. Himeno, H. *et al.* (2014) tmRNA-mediated trans-translation as the major ribosome rescue system in a bacterial cell. *Front. Genet.* 5, 66
78. Weber, C. *et al.* (2014) Playing RNase P evolution: swapping the RNA catalyst for a protein reveals functional uniformity of highly divergent enzyme forms. *PLoS Genet.* 10, e1004506

79. Gößringer, M. *et al.* (2017) Protein-only RNase P function in *Escherichia coli*: viability, processing defects and differences between PRORP isoenzymes. *Nucleic Acids Res.* 45, 7441–7454
80. Howard, M.J. *et al.* (2016) Differential substrate recognition by isozymes of plant protein-only ribonuclease P. *RNA* 22, 782–792
81. Walczyk, D. *et al.* (2016) Analysis of the cleavage mechanism by protein-only RNase P using precursor tRNA substrates with modifications at the cleavage site. *J. Mol. Biol.* 428, 4917–4928
82. Pavlova, L. *et al.* (2012) tRNA processing by protein-only versus RNA-based RNase P: kinetic analysis reveals mechanistic differences. *ChemBioChem* 13, 2270–2276
83. Howard, M.J. *et al.* (2015) Mechanistic studies reveal similar catalytic strategies for phosphodiester bond hydrolysis by protein-only and RNA-dependent ribonuclease P. *J. Biol. Chem.* 290, 13454–13464
84. Orans, J. *et al.* (2011) Structures of human exonuclease 1 DNA complexes suggest a unified mechanism for nuclease family. *Cell* 145, 212–223
85. Shi, Y. *et al.* (2017) Interplay of catalysis, fidelity, threading, and processivity in the exo- and endonucleolytic reactions of human exonuclease I. *Proc. Natl. Acad. Sci. U. S. A.* 114, 6010–6015
86. Beese, L.S. and Steitz, T.A. (1991) Structural basis for the 3'-5' exonuclease activity of *Escherichia coli* DNA polymerase I: a two metal ion mechanism. *EMBO J.* 10, 25–33
87. Freemont, P.S. *et al.* (1988) Cocystal structure of an editing complex of Klenow fragment with DNA. *Proc. Natl. Acad. Sci. U. S. A.* 85, 8924–8928
88. Pfeiffer, T. *et al.* (2000) Effects of phosphorothioate modifications on precursor tRNA processing by eukaryotic RNase P enzymes. *J. Mol. Biol.* 298, 559–565
89. Warnecke, J.M. *et al.* (1999) Role of metal ions in the hydrolysis reaction catalyzed by RNase P RNA from *Bacillus*. *J. Mol. Biol.* 290, 433–445
90. Thomas, B.C. *et al.* (2000) Evidence for an RNA-based catalytic mechanism in eukaryotic nuclear ribonuclease P. *RNA* 6, 554–562
91. Li, Xinqiang and Gegenheimer, P. (1997) Ribonuclease P catalysis requires Mg^{2+} coordinated to the pro-RP oxygen of the scissile bond. *Biochemistry* 36, 2425–2438
92. Cognat, V. *et al.* (2013) PlantRNA, a database for tRNAs of photosynthetic eukaryotes. *Nucleic Acids Res.* 41, D273–D279
93. Jühling, F. *et al.* (2009) tRNAdb 2009: compilation of tRNA sequences and tRNA genes. *Nucleic Acids Res.* 37, D159–D162