

## ORIGINAL ARTICLE

# The complete genome sequence of the algal symbiont *Dinoroseobacter shibae*: a hitchhiker's guide to life in the sea

Irene Wagner-Döbler<sup>1</sup>, Britta Ballhausen<sup>1</sup>, Martine Berger<sup>2</sup>, Thorsten Brinkhoff<sup>2</sup>, Ina Buchholz<sup>1</sup>, Boyke Bunk<sup>3</sup>, Heribert Cypionka<sup>2</sup>, Rolf Daniel<sup>4</sup>, Thomas Drepper<sup>5</sup>, Gunnar Gerdts<sup>6</sup>, Sarah Hahnke<sup>2</sup>, Cliff Han<sup>7</sup>, Dieter Jahn<sup>3</sup>, Daniela Kalhoefer<sup>2</sup>, Hajnalka Kiss<sup>7</sup>, Hans-Peter Klenk<sup>8</sup>, Nikos Kyrpides<sup>7</sup>, Wolfgang Liebl<sup>9</sup>, Heiko Liesegang<sup>4</sup>, Linda Meincke<sup>7</sup>, Amrita Pati<sup>7</sup>, Jörn Petersen<sup>8</sup>, Tanja Piekarski<sup>3</sup>, Claudia Pommerenke<sup>1</sup>, Silke Pradella<sup>8</sup>, Rüdiger Pukall<sup>8</sup>, Ralf Rabus<sup>2</sup>, Erko Stackebrandt<sup>8</sup>, Sebastian Thole<sup>2</sup>, Linda Thompson<sup>7</sup>, Petra Tielen<sup>3</sup>, Jürgen Tomasch<sup>1</sup>, Mathias von Jan<sup>8</sup>, Nittaya Wanphrut<sup>3</sup>, Antje Wichels<sup>6</sup>, Hajo Zech<sup>2</sup> and Meinhard Simon<sup>2</sup>

<sup>1</sup>Helmholtz-Centre for Infection Research, Braunschweig, Germany; <sup>2</sup>Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, Oldenburg, Germany; <sup>3</sup>Institute of Microbiology, Technical University of Braunschweig, Braunschweig, Germany; <sup>4</sup>Institute for Microbiology and Genetics, University of Göttingen, Göttingen, Germany; <sup>5</sup>Forschungszentrum Jülich, Jülich, Germany; <sup>6</sup>Biologische Anstalt Helgoland, Alfred-Wegener Institute for Polar and Marine Research, Bremerhaven, Germany; <sup>7</sup>Joint Genome Institute, San Francisco, CA, USA; <sup>8</sup>German Collection for Microorganisms and Cell Cultures, Braunschweig, Germany and <sup>9</sup>Technical University of München, München, Germany

***Dinoroseobacter shibae* DFL12<sup>T</sup>**, a member of the globally important marine *Roseobacter* clade, comprises symbionts of cosmopolitan marine microalgae, including toxic dinoflagellates. Its annotated 4417868 bp genome sequence revealed a possible advantage of this symbiosis for the algal host. *D. shibae* DFL12<sup>T</sup> is able to synthesize the vitamins B<sub>1</sub> and B<sub>12</sub> for which its host is auxotrophic. Two pathways for the *de novo* synthesis of vitamin B<sub>12</sub> are present, one requiring oxygen and the other an oxygen-independent pathway. The *de novo* synthesis of vitamin B<sub>12</sub> was confirmed to be functional, and *D. shibae* DFL12<sup>T</sup> was shown to provide the growth-limiting vitamins B<sub>1</sub> and B<sub>12</sub> to its dinoflagellate host. The *Roseobacter* clade has been considered to comprise obligate aerobic bacteria. However, *D. shibae* DFL12<sup>T</sup> is able to grow anaerobically using the alternative electron acceptors nitrate and dimethylsulfoxide; it has the arginine deiminase survival fermentation pathway and a complex oxygen-dependent Fnr (fumarate and nitrate reduction) regulon. Many of these traits are shared with other members of the *Roseobacter* clade. *D. shibae* DFL12<sup>T</sup> has five plasmids, showing examples for vertical recruitment of chromosomal genes (*thiC*) and horizontal gene transfer (*cox* genes, gene cluster of 47 kb) possibly by conjugation (*vir* gene cluster). The long-range (80%) synteny between two sister plasmids provides insights into the emergence of novel plasmids. *D. shibae* DFL12<sup>T</sup> shows the most complex viral defense system of all *Rhodobacterales* sequenced to date.

The ISME Journal (2010) 4, 61–77; doi:10.1038/ismej.2009.94; published online 10 September 2009

**Subject Category:** integrated genomics and post-genomics approaches in microbial ecology

**Keywords:** symbiosis; vitamin; cobalamin; thiamine; *Roseobacter*; dinoflagellate

## Introduction

One of the most abundant and metabolically versatile groups of bacteria in the world's oceans is

the *Roseobacter* clade. Members of this alphaproteobacterial lineage are prominently involved in the global marine carbon and sulfur cycles (Moran *et al.*, 2004; Selje *et al.*, 2004; Buchan *et al.*, 2005; Wagner-Döbler and Biebl, 2006; Howard *et al.*, 2006; Moran and Miller, 2007; Brinkhoff *et al.*, 2008). Many *Roseobacter* species live as epibionts on marine algae where they reach high abundance during phytoplankton blooms (reviewed by Buchan *et al.* (2005)). Some phytoplankton blooms, the so-called

Correspondence: I Wagner-Döbler, Helmholtz-Centre for Infection Research, Inhoffenstr. 7, Braunschweig, Germany.

E-mail: iwd@helmholtz-hzi.de

Received 27 April 2009; revised 29 June 2009; accepted 14 July 2009; published online 10 September 2009

red tides, are caused by toxic dinoflagellates, and *Roseobacter* species were frequently isolated from their laboratory cultures (Lafay *et al.*, 1995; Prokic *et al.*, 1998; Miller and Belas, 2004), and also detected on dinoflagellates in nature (Hasegawa *et al.*, 2007). A direct interaction between these algae and their associated bacteria is suggested by the fact that *Roseobacter* cells have been found attached to the surface of toxic *Pfiesteria* species using fluorescent *in situ* hybridization (FISH) (Alavi *et al.*, 2001).

The *Roseobacter* isolates from dinoflagellate cultures must be able to grow on metabolites excreted by the algae because they are heterotrophs (Moran and Miller, 2007) and there are no carbon sources present in the mineral media used for cultivating phototrophic algae. Many *Roseobacter* bacteria can degrade dimethylsulfoniopropionate (Moran *et al.*, 2007), an osmoprotectant released in large quantities during algal blooms, and they show positive chemotaxis toward this compound (Yoch 2002; Miller *et al.*, 2004; Miller and Belas, 2006). A strain of the abundant RCA (*Roseobacter* clade-affiliated) cluster could be cultivated in coculture with an axenic dinoflagellate (Mayali *et al.*, 2008). These studies show that *Roseobacter* bacteria can thrive on algal metabolites. However, it is presently not known whether this relationship provides an advantage to the dinoflagellate host.

In this study, we report the complete genome sequence of *Dinoroseobacter shibae* DFL12<sup>T</sup>, the type strain of a species from the *Roseobacter* clade. It was isolated from *Prorocentrum lima*, a benthic dinoflagellate, by picking a single dinoflagellate cell from a culture, washing it several times and placing it on an agar surface (Allgaier *et al.*, 2003; Biebl *et al.*, 2005). *P. lima* produces okadaic acid, which can cause diarrhetic shellfish poisoning during red tides (Pan *et al.*, 1999). *D. shibae* DFL12<sup>T</sup> was chosen for sequencing because of its symbiosis with dinoflagellates, its ability to perform light-driven ATP synthesis using bacteriochlorophyll *a* in the presence of oxygen (Allgaier *et al.*, 2003; Biebl and Wagner-Döbler, 2006; Wagner-Döbler and Biebl, 2006), the large number of extrachromosomal replicons (Pradella *et al.*, 2004) and its novel acylated homoserine lactone (AHL) compounds (Wagner-Döbler *et al.*, 2005). The complete genome sequence of *D. shibae* DFL12<sup>T</sup> revealed traits that presumably are highly adaptive in the habitat of bloom-forming algae and that may be characteristic for the whole *Roseobacter* clade. Moreover, we discovered an essential gift provided to the dinoflagellate by the bacteria.

## Materials and methods

### Sequencing, assembly and finishing

The genome of *D. shibae* DFL12<sup>T</sup> was sequenced at the Joint Genome Institute (JGI) Production Geno-

mic Facility using a combination of 3 and 8 kb (plasmid), and 40 kb (fosmid) DNA libraries. All general aspects of library construction and sequencing performed at the JGI can be found at <http://www.jgi.doe.gov>. The Phred/Phrap/Consed software package (<http://www.phrap.com>) was used for sequence assembly and quality assessment (Ewing and Green, 1998; Ewing *et al.*, 1998; Gordon *et al.*, 1998). After the shotgun stage, reads were assembled with parallel phrap (High Performance Software, LLC). Mis-assemblies were corrected with Dupfinisher (Han and Chain, 2006) or transposon bombing of bridging clones (Epicentre Biotechnologies, Madison, WI, USA). Gaps between contigs were closed by editing in Consed, custom primer walk or PCR amplification. A total of 742 additional reactions were necessary to close the gaps and to raise the quality of the finished sequence. The completed genome sequences of *D. shibae* DFL12<sup>T</sup> contain 67 596 reads, achieving an average of a 11-fold sequence coverage per base with an error rate less than 1 in 100 000. The sequences of *D. shibae* DFL12<sup>T</sup>, comprising a chromosome and five plasmids, can be accessed using the GenBank accession numbers NC\_009952, NC\_009955, NC\_009956, NC\_009957, NC\_009958 and NC\_009959. Manual curation and reannotation of the genome was carried out using The Integrated Microbial Genomes Expert Review System (img/er <http://imgweb.jgi-psf.org>) (Markowitz *et al.*, 2008) and the Artemis software package (<http://www.sanger.ac.uk/Software/Artemis/v9>).

### Phylogenetic analysis

Phylogenetic analyses of 16S rRNA gene sequences were performed with the ARB software package (<http://www.arb-home.de>) (Ludwig *et al.*, 2004). A phylogenetic tree was constructed using neighbor-joining and maximum-likelihood analyses. Only sequences with more than 1200 bp were considered in these calculations. Sequences <1200 bp were added afterward using parsimony.

### Design of *D. shibae*-specific oligonucleotides for PCR and FISH

Specific primers and probes were developed using the ProbeDesign function of the ARB software package (<http://www.arb-home.de>). Sequences of specific primers are DSH176f (5'-CGTATGTGGCC TTCGGGC-3') and DSH1007r (5'-GCTTCCGTCTCC GGAAGCC-3'). Specificity of primer sequences was checked with the NCBI and RDP databases (<http://www.ncbi.nlm.nih.gov> and <http://rdp.cme.msu.edu/>) and resulted in at least one mismatch to 16S rRNA gene sequences of non-target organisms. Probe DSH176-HRP (5'-GCCCCAAGGCCACATACG-3') was used for FISH, in combination with helper probes DSH176-H1 (5'-GTATTACTCCAGTTTCC CA-3') and DSH176-H2 (5'-ATCCTTTGGCGATAAA TCTTTC-3').

### Sample preparation for CARD-FISH

In total, 5 ml of the *Prorocentrum* culture were fixed with 5 ml paraformaldehyde (4% w/v) for 1 h and filtered onto a 0.2- $\mu$ m filter (Nucleopore Track Etch PC MB 25 mm, Whatman Nr. 1110656, Whatman plc, Maidstone, Kent, UK) or a 10- $\mu$ m filter (Polycarbonate Membrane Filters, TCTP04700, Millipore Corporation, Billerica, MA, USA). Cells were washed eight times with distilled water. Reference strains *D. shibae* DFL12<sup>T</sup> (0 mismatches to probe DSH176) and *Sulfitobacter guttiformis* DSM11458<sup>T</sup> (two mismatches to probe DSH176) were filtered onto 0.2- $\mu$ m polycarbonate filters and fixed with paraformaldehyde (2% w/v) for 1 h, rinsed with 1 ml 1  $\times$  phosphate buffer solution and subsequently with 1 ml distilled water. Filters were air dried and stored at  $-20^{\circ}\text{C}$  until further processing.

### CARD-FISH

CARD-FISH (Catalyzed reporter deposition FISH) was performed according to Sekar *et al.* (2003). Hybridization conditions were as follows: 2 h of hybridization at 35  $^{\circ}\text{C}$  (60% formamide), 30 min washing at 37  $^{\circ}\text{C}$  (14 mM NaCl) and 30 min amplification at 37  $^{\circ}\text{C}$ . Per 400  $\mu$ l hybridization buffer, 15  $\mu$ l of the HRP (horseradish peroxidase) probe DSH176-HRP and 15  $\mu$ l of each unlabeled helper oligonucleotide (DSH176-H1 and DSH176-H2) were used (probe solutions 50 ng  $\mu$ l<sup>-1</sup>): Tyramine-HCl was labeled with fluorescein-5-isothiocyanate as described by Pernthaler *et al.* (2002). To avoid unspecific accumulation of dye in the cells, the last washing step in phosphate buffer solution (1  $\times$ ) amended with TritonX-100 (0.05%) was extended to 30 min. Counterstaining was performed with Vectashield-mounting medium with 4',6-Diamidino-2-phenylindole (1.5  $\mu$ g ml<sup>-1</sup>; Vector Laboratories, Peterborough, England).

### Comparative genomics

The Integrated Microbial Genomes Expert Review System (img/er <http://imgweb.jgi-psf.org>) (Markowitz *et al.*, 2008) was used for comparative analysis on the basis of the nine presently available completely sequenced genomes of the *Rhodobacteriales*: *Jannaschia* sp. CCS1 (Moran *et al.*, 2007), *Roseobacter denitrificans* (Swingley *et al.*, 2007), *Silicibacter pomeroyi* (Moran *et al.*, 2004), *Silicibacter* sp. TM1040 (Moran *et al.*, 2007), three strains of *Rh. sphaeroides* (Choudhary *et al.*, 2007) as well as *Paracoccus denitrificans* (JGI Institute, unpublished).

### CRISPR

Homologs of *cas* genes were detected using tools of the BLAST package (Altschul *et al.*, 1990). CRISPR arrays were detected by the CRISPR recognition tool (Bland *et al.*, 2007). Comparative analysis of

repeat sequences was done both manually and by CRISPRdb (Grissa *et al.*, 2007). The origin of spacer sequences was searched for in the CAMERA databases (Seshadri *et al.*, 2007) (<http://camera.calit2.net>).

Regarding the methods for dinoflagellate cultivation and sampling, isolation of genomic DNA, PCR, cloning and sequencing, determination of vitamin B<sub>12</sub> and coculture experiments, see Supplementary material 10.


## Results and discussion

### Ecological niche and phylogeny


*D. shibae* DFL12<sup>T</sup> was isolated from *P. lima* (Biebl *et al.*, 2005). This phototrophic dinoflagellate occurs in sand and sediment and to some extent also in the water column, but reaches its highest densities in biofilms on marine macroalgae, macrophytes, corals, mussels and oysters (Levasseur *et al.*, 2003; Vershinin *et al.*, 2005; Okolodkov *et al.*, 2007; Parsons and Preskitt, 2007).

Several closely related strains were isolated from the dinoflagellate *Alexandrium ostenfeldii* (Allgaier *et al.*, 2003). Using PCR with *D. shibae*-specific primers for the 16S rRNA gene and sequencing of the obtained bands for confirmation, *D. shibae* was also detected in cultures of *Protoceratium reticulatum*, a dinoflagellate isolated from the North Sea. All three host species are cosmopolitan, toxic algae causing diarrhetic shellfish poisoning (Aasen *et al.*, 2005). Finally, *D. shibae* was isolated from *Isochrysis galbana*, a member of the class *Haptophyceae* (Genbank accession EF512132). This nontoxic alga is cultivated as food for bivalves in aquaculture. Thus, *D. shibae* is not restricted to toxic dinoflagellates, but is associated with a variety of cosmopolitan marine microalgae. Figure 1 shows the phylogenetic position of all currently known *D. shibae* strains within the *Roseobacter* clade and indicates from which algae they were isolated. The similarity of these sequences to *D. shibae* DFL12<sup>T</sup> is between 97.2% and 99.8%, indicating that they probably belong to the same species, but additional data are needed (Rossello-Mora and Amann, 2001; Stackebrandt *et al.*, 2002; Gevers *et al.*, 2005).

A physical association between host algae and bacteria is suggested by the fact that *D. shibae* DFL12<sup>T</sup> was isolated from single washed cells of the dinoflagellate (Biebl *et al.*, 2005). Moreover, it can be seen attached to *P. lima* in coculture (Figure 2). In this study, dinoflagellates were separated from suspended bacteria using 10- $\mu$ m polycarbonate filters, washed eight times and stained using CARD-FISH with probes specific for *D. shibae* DFL12<sup>T</sup>. Many bacteria can be seen adhering to the surface of the dinoflagellate, and some are freely suspended in the medium. It remains to be seen if a genetic program coordinates the switch between planktonic growth and growth as an epibiont in *D. shibae*.



**Figure 1** Phylogenetic position of *Dinoroseobacter shibae* DFL12<sup>T</sup> and closely related strains and phylotypes within the *Rhodobacteraceae* (*Alphaproteobacteria*) based on 16S rRNA gene comparisons. All names written in boldface indicate organisms whose complete genome sequences have already been determined. The phylogenetic tree was generated using the maximum-likelihood method. Selected members of the *Gammaproteobacteria* were used as an outgroup (not shown). The eukaryotic host algae from which the strains were isolated are indicated next to the dotted brackets. *Pr.*, *Protoceeratium* (Dinophyceae); *I.*, *Isochrysis* (Haptophyceae); *P.*, *Prorocentrum* (Dinophyceae); *A.*, *Alexandrium* (Dinophyceae). Bar = 0.1 substitutions per nucleotide position.




**Figure 2** Specific attachment of *D. shibae* DFL12<sup>T</sup> to its dinoflagellate host *Prorocentrum lima*. (a) Differential interference contrast (DIC) image of dinoflagellates from a non-axenic culture to which cells of strain DFL12<sup>T</sup> had been added 2 months before; (b) Catalyzed reporter deposition fluorescent *in situ* hybridization (CARD-FISH) with *Dinoroseobacter*-specific 16S rRNA probes showing most cells of *D. shibae* DFL12<sup>T</sup> (green dots) closely attached to the dinoflagellate. Combined images generated by means of PICOLAY (<http://www.picolay.de>) from a series of six pictures taken at sequential focus levels. Bar = 20  $\mu$ m. The central round structure in the cell center seen in (a) is the pyrenoid, a specialized area of the plastid that contains high levels of Rubisco. It is typically surrounded by a starch sheath that can also be seen. The crystal-like ultrastructure of the pyrenoid (Kowalik, 1969) may be the reason for its orange appearance in the left cell in (b).

#### General characteristics of the genome

The *D. shibae* type strain DFL12<sup>T</sup> (=DSM 16493<sup>T</sup>=NCIMB 14021<sup>T</sup>) has a genome size of 4417 kbp, similar to the other fully sequenced *Roseobacter* genomes (Supplementary material S1). The genome contains 4198 protein-encoding genes.

About 28% of the derived gene products have no predicted function. The finished genome sequence indicates a circular conformation of the chromosome and the five plasmids (Figure 3). Previously the presence of plasmids with linear conformation was hypothesized for two *D. shibae* strains and



**Figure 3** Circular representation of the chromosome and plasmids of *D. shibae* DFL12<sup>T</sup>. The ring of the chromosome depicts (starting with the outer ring) (1) the scale in basepairs; (2) all protein-encoding genes, clockwise-transcribed genes in blue and counterclockwise-transcribed genes in yellow; (3) alien genes, that is, genes that are likely acquired by horizontal gene transfer; (4) genes related to DNA flexibility: transposases (blue), integrases (red) and recombinases (green); (5) rRNA and tRNA genes; (6) the GC content. The rings of the plasmids depict (1) the scale in base pairs; (2) all protein-encoding genes; (3) genes related to DNA flexibility; (4) rRNA and tRNA genes and (5) the GC content. Alien genes are not shown in the plasmids, because these contigs are too small for their calculation.

several related species (Pradella *et al.*, 2004). We investigated the plasmid structure in *D. shibae* DFL12 by a restriction assay combined with length separation through pulsed-field gel electrophoresis and could confirm the circular conformation (Supplementary material S2). This is in agreement with the absence of genes typically associated with linear

plasmids, for example, the telomere resolvase (Meinhardt *et al.*, 1997; Kobryn and Chaconas, 2001).

#### Comparative genomics

*D. shibae* DFL12<sup>T</sup> harbors all biogeochemically important traits reported previously from fully

sequenced *Roseobacter* isolates, namely anaerobic anoxygenic photosynthesis (*pufLM* genes), carbon monoxide oxidation (*cox* genes), aromatic compound degradation (*pcaGH* and *boxC* genes), sulfur oxidation (*soxB*), denitrification (*nirS/K*), nitrate assimilation (*nasA*), phosphonate use (*phn*), type IV secretion (*vir* genes) and dimethylsulfoniopropionate degradation (*dmdA*) (Supplementary material S3). Several families of transposases/integrases (61 and 43 copies, respectively) and site-specific recombinases/resolvases (11 copies) belong not only to the largest gene families, but also they are more frequent than in other members of the *Roseobacter* clade, indicating a large potential for DNA exchange in *D. shibae* DFL12<sup>T</sup>. Proteins involved in information processing usually represent the role categories showing the least variation between closely related organisms. Their comparative analysis is described in Supplementary material S1.

## The central energy metabolism and its regulation

### Central carbon metabolism

The central carbon metabolism is reconstructed in Supplementary material S4. Glucose breakdown is potentially possible through glycolysis (Embden-Meyerhoff-Parnas pathway) and the Entner-Doudoroff pathway, the latter being closely connected to the oxidative pentose phosphate pathway. Interestingly, a fluxom analysis using <sup>13</sup>C labeling techniques showed that glucose breakdown is carried out exclusively using the Entner-Doudoroff pathway; thus, the alternative routes are used only for anabolic purposes (Fürch *et al.*, 2009). The Calvin cycle is lacking, indicated by the absence of the key enzyme Rubisco, as in all *Roseobacter* strains (Swingley *et al.*, 2007). Carbon dioxide might be fixed by anaerobic enzymes either through (1) a two-step reaction involving pyruvate-orthophosphate dikinase and phosphoenolpyruvate carboxylase or through (2) pyruvate carboxylase. Genes for both carboxylases are present in all fully sequenced *Roseobacter* strains. Pyruvate carboxylase requires biotin as a cofactor, and hence the need to supply *D. shibae* DFL12<sup>T</sup> with this vitamin when it is grown on minimal medium (Biebl *et al.*, 2005). A fluxom analysis revealed that, indeed, this enzyme displays the major CO<sub>2</sub>-assimilating route in *D. shibae* (Fürch *et al.*, 2009). *D. shibae* DFL12<sup>T</sup> is the only fully sequenced *Roseobacter* strain harboring the arginine deiminase pathway, which allows one to carry out a survival fermentation process known from the *Bacillus* and *Pseudomonas* species (Eschbach *et al.*, 2004; Williams *et al.*, 2007).

### Electron transport chains

Complex electron transport systems were found (Supplementary material S5), including seven electron-donating primary dehydrogenases, such as for

L- and D-lactate, glycerol-3-phosphate, NADH and succinate, and respiratory dehydrogenases for glucose and gluconate. The presence of the aerobically performed anoxygenic photosynthesis further complicates the electron transport machinery. Just as in the mitochondria, which are derived from *Alpha-proteobacteria*, a cytochrome *bc*<sub>1</sub> complex is available. Electron flux can occur from ubiquinol, the reduced form of ubiquinone, through the *bc*<sub>1</sub> complex and different cytochrome *c* molecules to two types of cytochrome *c* oxidases to reduce molecular oxygen to water. Consequently, a close coordination between light-dependent cyclic electron transport (photosynthesis) and respiration is required, the mechanism of which remains to be elucidated.

A complete denitrification pathway and a dimethylsulfoxide (DMSO) reductase are available for the reduction of the alternative electron acceptors DMSO and nitrate. Interestingly, the organism uses the Nap (periplasmic) type nitrate reductase, which is oriented toward the periplasmic space and is active both under aerobic and anaerobic conditions, instead of the Nar (respiratory) type facing the cytoplasm, and is induced only under anaerobic/microaerophilic conditions. Anaerobic growth using nitrate as an electron acceptor was confirmed experimentally in a defined mineral medium (data not shown).

### Oxygen-dependent regulation of the central energy metabolism

We deduced the regulon for the global oxygen-dependent transcriptional regulator for fumarate and nitrate reduction, Fnr (Supplementary material S3). Six Fnr/Crp (cAMP receptor protein) type regulators are encoded by the genome of *D. shibae* DFL12<sup>T</sup>. However, only one of the deduced proteins carries the cysteine residues required for iron sulfur cluster formation. Two others resemble Fnr-type regulators such as Dnr and DnrD (transcription factor for denitrification gene expression). Interestingly, Dnr is located directly upstream of the *nos* operon encoding a nitrous oxide reductase, the enzyme catalyzing the transformation of N<sub>2</sub>O to N<sub>2</sub>, the last step of denitrification. Such a gene arrangement has not been found before in denitrifying bacteria, for example, in *R. denitrificans*.

Using a position weight matrix approach (Munch *et al.*, 2005), the genome was searched for potential Fnr-binding sites. In agreement with findings for other Gram-negative bacteria, the promoter region of genes for high-affinity oxygen-dependent cytochrome *c* oxidases (*cbb*<sub>3</sub>-type), alternative anaerobic systems including the NADH dehydrogenase, the quinone oxidoreductase and the corresponding cofactor biosynthesis genes (*hemN1*, *hemN2*, *hemA* and *moaC*) are all carrying Fnr-binding sites. No obvious Fnr boxes were found upstream of the various operons involved in denitrification and DMSO reduction. Interestingly, three universal

stress protein-like genes and eight potential regulator genes are also carrying Fnr boxes. Two of these regulators are suppressor proteins of the heat shock chaperone DnaK (DksA), and the function of the other six regulators is unknown. The Fnr box is also present in promoters of bacteriochlorophyll biosynthesis genes, but it is not possible to predict whether Fnr induces or represses the expression of the corresponding genes.

The Fnr regulator is not a unique feature of *D. shibae* DFL12<sup>T</sup>. Searching the complete genome sequences of the *Roseobacter* clade with the *D. shibae* Fnr by reciprocal best hit BLAST showed that one copy is present in all of them. Thus, the ability to fine-tune their metabolism to anoxic conditions is a general trait of the *Roseobacter* clade.

#### *Nutrients provided by the algal host and their uptake*

The pathways for the known growth substrates of *D. shibae* DFL12<sup>T</sup> (Biebl *et al.*, 2005) could be reconstructed from the genome sequence. Interestingly, most of them are Krebs cycle intermediates (fumarate, succinate, pyruvate and citrate). They may be released from the algal host after cell death or during photosynthesis. Photosynthate release from phytoplankton can be up to 40% of primary production (Wang and Douglas, 1997) and forms the basis of the microbial loop (Azam *et al.*, 1983). *D. shibae* DFL12<sup>T</sup> also uses acetate and glucose, important components of the marine dissolved organic carbon (Aluwihare *et al.*, 2002; Biersmith and Benner, 1998). Growth also occurs on fructose and glyceraldehyde-3-phosphate, products of the photosynthetic Calvin–Benson cycle. The dimethylsulfoniopropionate breakdown product DMSO can be used as an alternative electron acceptor by *D. shibae* DFL12<sup>T</sup>.

For the uptake of nutrients from the environment, tripartite ATP-independent periplasmic transporters (TRAP) seem to be preferentially used by *D. shibae* DFL12<sup>T</sup> rather than ABC-type transporters, which are only about half as frequent as in other members of the *Roseobacter* clade. TRAP transporters are composed of a periplasmic solute receptor (DctP or TAXI) and a secondary transporter (DctM(Q)) (Forward *et al.*, 1997). Transport is driven by an electrochemical ion gradient rather than by ATP hydrolysis, and this may be the reason why TRAP systems have their highest prevalence among marine bacteria (Rabus *et al.*, 1999; Mulligan *et al.*, 2007). Their design allows the coupling of high-affinity solute reception (formerly attributed only to primary transporters) with energy-efficient ion gradient-driven permeases. The substrate range of TRAP systems also comprises, besides C4-dicarboxylates, pyruvate and two other oxoacids, glutamate, sialic acid, ectoine and 2,3-diketogulonate. With 27 complete TRAP systems encoded in its genome, *D. shibae* DFL12<sup>T</sup> is second only to *S. pomeroyi* (28 complete systems). Efficient uptake of organic

nutrients available only in the sub-micromolar concentration range could provide members of the *Roseobacter* clade with a competitive advantage.


## Plasmid biology

### *Characteristic features of the plasmids*

The most striking observation regarding the extra-chromosomal elements is the long-range synteny between the 191 and the 126 kb plasmid (Figure 4). About 80% of the sequence of the smaller replicon, including a gene cluster of 47 kb, a *vir* operon and a short inverted region (shown in blue), are highly conserved. This pattern provides strong evidence for a common origin, thus justifying their designation as sister plasmids. The differences among the non-conserved regions result from genomic rearrangements likely assisted by transposition events, an explanation that is supported by the frequent occurrence of transposases and integrases (Supplementary material S6A). Moreover, comparative sequence analyses indicate that a common ancestor of the two sister plasmids may have been recruited through conjugational gene transfer. The *D. shibae* DFL12<sup>T</sup> chromosome has a GC content of 66%, which is in the upper range of *Roseobacters*, and the respective values for the plasmids vary between 60% and 69% (Supplementary material S6A). The two sister plasmids show the lowest GC content and the analysis with the Artemis software revealed a poor so-called ‘GC Frame Plot’ (Supplementary material 6B–D). This finding probably reflects their relatively recent recruitment in an evolutionary time scale and an early stage of sequence adaptation within the new host bacterium. This scenario is supported by the observation that in *D. shibae*, DFL12<sup>T</sup> *vir* operons are exclusively present on the sister plasmids (Figure 4). The *vir* gene cluster encoding the type IV secretion system is the structural prerequisite for the formation of sex pili required for conjugation. It is structurally highly conserved among *Roseobacter* strains and also present in *Silicibacter* sp. TM1040 (Moran *et al.*, 2007).

### *Plasmid replication and maintenance*

We identified the core functions for plasmid replication initiation, partitioning and stability representing the ‘functional heart’ of a plasmid (Supplementary material 7A–C). They provide the structural basis for compatibility versus incompatibility (Petersen *et al.*, 2009). We found three different types for replication initiation including the *repABC* module that is characteristic for the sister plasmids (Figure 4). Our phylogenetic analyses document that their crucial replicases (*repC*) are only distantly related, indicating that at least one of them has been exchanged by a gene transfer event. The respective *repABC* modules hence belong to



**Figure 4** Synteny plot of the 191 and 126 kb plasmids of *D. shibae* DFL12<sup>T</sup>. Long-range homologies are shown with red bars and an inverted region is printed in blue. The left and right borders of the plasmids are completely conserved. Homologous *vir* clusters and the distantly related *repABC*-type replicons as well as the adjacent toxin/antidote addition systems for plasmid stability are shown in detail. Adjacent genes that are not involved in plasmid core functions are indicated with dotted lines. The analysis was calculated with the Artemis Comparison Tool (WebACT, <http://www.webact.org/WebACT/generate>).

different compatibility groups (Petersen *et al.*, 2009), the prerequisite for a stable coexistence of the syntenous plasmids. Finally, at least four of the five replicons contain putative addiction systems for plasmid maintenance (Zielenkiewicz and Ceglowski, 2001). They are represented by operons encoding a stable toxin as well as a less stable antidote (Figure 4). These selfish units indirectly ensure the maintenance of the low copy replicons in the offspring, as plasmid loss would be lethal for the host cell.

#### Metabolic capabilities and localization of genes

The extrachromosomal replicons of *D. shibae* DFL12<sup>T</sup> can roughly be classified according to their genetic composition (Supplementary material S7D). The two sister plasmids are characterized by the *vir* gene cluster (see above), whereas the other plasmids harbor gene clusters for the degradation of aromatic compounds (153 kb), *cox* operons that presumably contribute to energy production (72 kb) and a wealth

of genes for sugar metabolism (86 kb). The latter replicon represents an aged and possibly dying plasmid owing to its conspicuous amount of pseudogenes and transposases. Transposition may dominate intracellular genetic rearrangements as documented for *thiC*, an essential gene for thiamine (vitamin B<sub>1</sub>) biosynthesis (Vander Horn *et al.*, 1993). In contrast to all other genes of this pathway that are localized on the *D. shibae* DFL12<sup>T</sup> chromosome, altogether three copies of *thiC* are located on different plasmids (191 kb, 153 kb and 86 kb). The genomic neighborhood of an integrase, resolvase and transposase strongly supports an underlying dispersal mechanism based on DNA transposition.

## Interaction with other organisms

### Quorum sensing

*D. shibae* DFL12<sup>T</sup> produces three different quorum-sensing signals from the group of AHLs, namely a



C<sub>8</sub>-homoserine lactone (C8-HSL), and two AHLs with a side chain of 18 carbon atoms, C<sub>18</sub>-en-HSL and C<sub>18</sub>-dien-HSL (Wagner-Döbler et al., 2005). This is the maximum side chain length for AHLs and it was, for the first time, detected in *Sinorhizobium meliloti*, a root nodule-forming symbiont (Marketon et al., 2002), where this signal downregulates motility at high cell density (Hoang et al., 2008). The unsaturations found in the C<sub>18</sub> side chain of DFL12<sup>T</sup> are novel and may be critical for signal specificity. The genome analysis predicted three autoinducer synthases (*luxI* type) and five *luxR* type genes for AHL-controlled transcriptional regulators (Supplementary material S3). *luxI*<sub>1</sub> and *luxI*<sub>2</sub> of *D. shibae* DFL12<sup>T</sup> are located on the chromosome adjacent to a *luxR* gene with an interspacer region of appr. 100 kb, whereas *luxI*<sub>3</sub> is on the 86-kb plasmid without an adjacent *luxR* gene. *Roseobacter* strains usually have one or two autoinducer synthases, whereas up to four have been found in root nodule-forming *Rhizobia* (Case et al., 2008). The orphan *luxR* genes may allow eavesdropping on competing bacterial species in the environment (Case et al., 2008).

#### Production of secondary metabolites

Polyketide synthases (PKSs) catalyze the synthesis of polyketides, a large class of secondary metabolites, from acetyl-CoA precursors. *D. shibae* DFL12<sup>T</sup> harbors a 12-gene cluster containing two adjacent modular type I PKS genes among hypothetical and putative proteins (Supplementary material S3). The first PKS gene is composed of only one module containing the domains essential for chain elongation ( $\beta$ -ketoacyl synthase domain, acyltransferase domain, and two ACP (acyl carrier protein) domains), a 2-nitropropane dioxygenase domain and a  $\beta$ -ketoacyl-ACP reductase domain. No reductive domain required for the modification of the keto-group could be identified. The second PKS gene contains two modules. In the N-terminal module, a  $\beta$ -ketoacyl synthase domain and an acyltransferase domain were identified; the C-terminal module shows similarity to a phosphopantetheinyltransferase domain (PKS loading enzyme). The other genes in this cluster, such as the two putative short-chain dehydrogenases or the two putative NAD-dependent epimerases/dehydratases, may be involved in chain modification reactions to give the final active product.

Close homologs of these genes are not found in the *Roseobacter* clade, but a similar gene cluster is harbored by *Solibacter usitatus* (*Acidobacteria*) and *Gloeobacter violaceus* (*Cyanobacteria*), suggesting that these genes may have been acquired by horizontal gene transfer. Thus, *D. shibae* DFL12<sup>T</sup> encodes a multiprotein complex that may catalyze the synthesis of a novel polyketide that could have a role in interactions with competing microbes. The dinoflagellate surface is colonized by a range of other bacteria, including *Vibrios* and *Flavobacteria*.


Tropodithietic acid, an antibiotic produced by at least two *Roseobacter* isolates, is especially active against *Vibrios* (Brinkhoff et al., 2004; Bruhn et al., 2005) and is encoded by a different pathway (Geng et al., 2008).

#### Resistance against viral attack

Viruses are the most abundant biological entity on the planet (Williamson et al., 2008). By lysing a significant fraction of bacterial communities, bacteriophages influence the cycling of organic matter in the sea (Fuhrman, 1999; Weinbauer, 2004; Brussaard et al., 2008; Danovaro et al., 2008) and the genetic diversity of their host populations (Muhling et al., 2005). Continuous viral attack accompanies red tides (Tomaru and Nagasaki, 2004) and contributes to their sudden collapse (Nagasaki et al., 2005; Rhodes et al., 2008). Thus, resistance against phage attack must be highly adaptive in the habitat of *D. shibae* DFL12<sup>T</sup>, which has evolved the most complex viral defense system of all *Roseobacter* strains sequenced to date.

CRISPRs (clustered regularly interspaced small palindromic repeats) are small, transcribed DNA spacers separated by short palindromic repeats that are located next to a cluster of *cas* (CRISPR associated) genes. Many spacer sequences have been shown to originate from bacteriophages, leading to the current hypothesis that CRISPRs are involved in an RNAi-like mechanism resulting in antiviral resistance (Haft et al., 2005; Makarova et al., 2006; Barrangou et al., 2007). Two CRISPRs were detected in the genome of *D. shibae* DFL12<sup>T</sup>, whereas no other completed genome from the *Rhodobacterales* contains more than one. Five organisms of this group revealed a CRISPR (Supplementary material S8). Remarkably, four of them lack the characteristic CRISPR marker gene *cas1*, which is reportedly present in all CRISPR-containing organisms, with *Pyrococcus abyssi*, as formerly, a single exception. Therefore, *cas1* is apparently no reliable marker for CRISPRs in the *Rhodobacterales*.

Figure 5 shows that the two CRISPR/*cas* gene clusters of *D. shibae* DFL12<sup>T</sup> have totally different structures. Comparative sequence analyses indicated that they are complete and independent units derived from different microbial lineages. It is most likely that their acquisition occurred through horizontal gene transfer, as suggested earlier (Godde and Bickerton, 2006). To discover the origin of the CRISPR spacer sequences in *D. shibae* DFL12<sup>T</sup>, these sequences were compared with the marine metagenomics database, CAMERA (Seshadri et al., 2007), but no homologs were found, most likely because the habitats of *D. shibae* DFL12<sup>T</sup> have not yet been searched for phages. Diversification of the CRISPR region is inferred to be a population-level response to the rapidly changing selective pressure of phage predation (Tyson and Banfield, 2008), and hence phage predation must represent a significant selective force in the habitat of *D. shibae* DFL12<sup>T</sup>.



**Figure 5** Genomic context of the two CRISPRs of *D. shibae* DFL12<sup>T</sup>. The numbers indicate the base position in the genome. Barcode-like series of vertical black bars symbolize CRISPR arrays. The colored gene arrows show the orientation, length and assignment to functional gene groups: Green—*cas* core genes; 1 = *cas1*; 2 = *cas2*; 5 = *cas5*. Purple—*csn1* gene, specific for subtype *Neisseria meningitidis*. Yellow—*cas* genes specific for subtype *Escherichia coli*; 2 = *cse2*; 3 = *cse3*; 4 = *cse4*. Orange—genes for polynucleotide processing; i = integrase; r = recombinase; e = excisionase.

## Synthesis and utilization of vitamin B<sub>12</sub>

### Vitamin B<sub>12</sub>-dependent enzymatic reactions


Most Bacteria and Archaea including *D. shibae* DFL12<sup>T</sup> have enzymes that require B<sub>12</sub> as a cofactor. The genome revealed a gene encoding a B<sub>12</sub>-dependent methionine synthase, catalyzing the final step in methionine biosynthesis. A gene encoding an enzyme that catalyzes the same reaction in a B<sub>12</sub>-independent manner was also found. Genes encoding a B<sub>12</sub>-dependent ribonucleotide reductase and a methylmalonyl-CoA mutase were also found, but genes for other B<sub>12</sub>-dependent enzymes (glutamate mutase, ethanolamine ammonia lyase, diol dehydratase and glycerol dehydratase) were not present.

### Cobalamin biosynthesis

Vitamin B<sub>12</sub>, which is only synthesized *de novo* by some Bacteria and Archaea, is derived from uroporphyrinogen III, a precursor in the synthesis of heme, siroheme, cobamides, chlorophylls and the methanogenic F430 (Scott and Roessner, 2002; Warren et al., 2002; Rodionov et al., 2003). At least 25 enzymes are uniquely involved (Roth et al., 1996). The genome of *D. shibae* DFL12<sup>T</sup> harbors all the genes required for *de novo* B<sub>12</sub> synthesis (Supplementary material S9), similar to other fully sequenced members of the *Roseobacter* clade, for example, *R. denitrificans* and *S. pomeroyi*. However, it has an unusual route for vitamin B<sub>12</sub> synthesis. Two different biosynthetic routes for cobalamin are known in Bacteria: (i) an oxygen-dependent (aerobic) pathway and (ii) an oxygen-independent (anaerobic) pathway (Martens et al., 2002; Scott and Roessner, 2002; Warren et al., 2002). They differ in

the first part, the insertion of cobalt into the corrin ring, whereas the second part is common for both the routes. In the aerobic route, cobalt insertion is performed by an ATP-dependent cobalt chelatase, which is encoded in *P. denitrificans* by *cobN*, *cobS* and *cobT*. In the anaerobic pathway, single subunit ATP-independent cobalt chelatases such as CbiX of *B. megaterium* are performing the reaction. *D. shibae* DFL12<sup>T</sup> contains putative genes for both types of chelatases and in this respect it is unique among the fully sequenced *Rhodobacterales*. Other pathway-specific genes were also present, but *cobG*, regarded as a signature gene for the aerobic route (Scott and Roessner, 2002), was absent. The gene for an alternative enzyme found in *Rhodobacter capsulatus*, *CobZ*, was also not present (McGoldrick et al., 2005). Thus, *D. shibae* DFL12<sup>T</sup> might have the property to synthesize B<sub>12</sub> by either the aerobic or the anaerobic pathway depending on the availability of oxygen. Alternatively, it might synthesize B<sub>12</sub> by an unusual combination of both pathways.

We experimentally confirmed that the *de novo* pathway for vitamin B<sub>12</sub> synthesis is functional in *D. shibae* DFL12<sup>T</sup>. The bacteria can grow in a defined mineral medium, but to exclude that traces of vitamin B<sub>12</sub> might be present even in such media, we compared the B<sub>12</sub> content of cells grown in the presence and absence of methionine. *D. shibae* DFL12<sup>T</sup> has two methionine synthases, one requiring B<sub>12</sub> and another that is B<sub>12</sub> independent. In the absence of methionine, the content of B<sub>12</sub> in the cells should increase, as the B<sub>12</sub>-dependent methionine synthase requires B<sub>12</sub> as a cofactor. The vitamin B<sub>12</sub> content of the cells was 340 ± 36 ng mg<sup>-1</sup> protein in the presence of methionine and increased to 450 ± 26 ng mg<sup>-1</sup> protein in the absence of methio-



**Figure 6** The bacterium *D. shibae* DFL12<sup>T</sup> provides thiamine (B<sub>1</sub>) and cobalamin (B<sub>12</sub>) to the dinoflagellate *Prorocentrum minimum* in coculture. Growth of the algae is shown as the amount of chlorophyll *a* (absorbance at 664 nm). Symbols show cultures on L1 mineral salts medium  $\Delta$  lacking thiamine;  $\square$  lacking cobalamin and  $\bullet$  control (complete L1 medium). On day 22 (dotted line), 1% of the cultures lacking thiamine or cobalamin were transferred to fresh medium of the same composition. One half of the culture was provided with the lacking vitamin (open symbols), and the other half was inoculated with *D. shibae* DFL12<sup>T</sup> (filled symbols). The inset shows log cell densities of *D. shibae* DFL12<sup>T</sup> at the time of inoculation and at the end of the experiment (cfu and direct microscopic counts). A full colour version of this figure is available at *The ISME Journal* online.

nine, showing that a complete *de novo* pathway for vitamin B<sub>12</sub> synthesis is functional.

#### Symbiosis with dinoflagellates


Besides supplying its own B<sub>12</sub>-dependent enzymes with the cofactor, the versatility in B<sub>12</sub> production might also contribute to a successful symbiosis with the dinoflagellate. Axenic (bacteria-free) cultures of dinoflagellates can be obtained on defined mineral media, but supplementation with three vitamins is necessary, namely thiamine (B<sub>1</sub>), cobalamin (B<sub>12</sub>) and biotin (B<sub>7</sub> or vitamin H) (Guillard and Hargraves, 1993). *D. shibae* DFL12<sup>T</sup> is able to synthesize two of these, B<sub>12</sub> and B<sub>1</sub>. It is auxotrophic for biotin, similar to many *Roseobacter* species (Shiba, 1991), and thus this vitamin must be provided by other sources in nature to both the dinoflagellate and the *Roseobacter* species.

To experimentally determine whether *D. shibae* DFL12<sup>T</sup> can provide B<sub>1</sub> and B<sub>12</sub> to its dinoflagellate host, we performed coculture experiments. Axenic cultures for *P. lima* are not available in culture collections, but as the symbiosis of *D. shibae* with algae is not strictly host specific, we reasoned that proof of principle might be obtained using the closely related dinoflagellate *P. minimum* CCMP1329.

First, auxotrophy of *P. minimum* for B<sub>1</sub> and B<sub>12</sub> was investigated, as it had not been tested earlier (Croft et al., 2005). Supplementary material S10 shows that *P. minimum* was not able to grow on a synthetic mineral medium from which either B<sub>1</sub> or B<sub>12</sub> had been omitted. Next, either B<sub>1</sub> or B<sub>12</sub> were replaced by  $1.1 \times 10^6$  cells per ml of *D. shibae*

DFL12<sup>T</sup>. In the experiment shown in Figure 6, the dinoflagellate was first cultivated for 22 days without bacteria; there was no growth without thiamine, and cobalamin became growth limiting after about 17 days. On day 22, the cultures were divided. One half was provided with the lacking vitamin, and the other half was inoculated with *D. shibae* DFL12<sup>T</sup>. Figure 6 shows that *D. shibae* DFL12<sup>T</sup> stimulated the growth of the alga in the absence of B<sub>1</sub> or B<sub>12</sub>, and that the alga allowed the bacterium to grow too (inlay in Figure 6). From an initial density of  $1.1 \times 10^6$  cfu ml<sup>-1</sup>, they increased to  $5.1 \times 10^6$  cfu ml<sup>-1</sup>. Bacterial densities determined by direct microscopic counts were even higher by more than one order of magnitude. The density of the dinoflagellate at the end of the experiment was appr.  $2.5 \times 10^6$  cells ml<sup>-1</sup>, as determined by microscopic counts. Supplementation with pure vitamins resulted in faster recovery of the dinoflagellate culture in the case of B<sub>12</sub>, and in much stronger recovery in the case of B<sub>1</sub> compared with inoculation with *D. shibae* DFL12<sup>T</sup>. The bacteria were entirely dependent on carbon sources excreted by the algae, and thus may have been initially starved in the non-growing algal culture. The concentration of thiamine present in the cultivation medium is 200-fold higher than that of B<sub>12</sub>, and therefore the amount synthesized by the relatively low density of bacteria might not have been sufficient.

Experiments with the haptophyte *I. galbana* CCMP1323, from which *D. shibae* DFL12<sup>T</sup> has also been isolated and which is known to be auxotrophic for B<sub>12</sub> (Croft et al., 2005), showed similar results for cobalamin, but this alga did not require thiamine for



***Prorocentrum minimum***  
**CCMP1329.**  
Media are (from left to right) L1 medium, L1 medium lacking cobalamin, and L1 medium lacking cobalamin inoculated with *D. shibae* DFL12<sup>T</sup>.

***Prorocentrum minimum***  
**CCMP1329.**  
Media are (from left to right) L1 medium, L1 medium lacking thiamine, and L1 medium lacking thiamine inoculated with *D. shibae* DFL12<sup>T</sup>.

***Isochrysis galbana***  
**CCMP1323.**  
Media are (from left to right) L1 medium, L1 medium lacking cobalamin, L1 medium lacking cobalamin inoculated with 10 µl *D. shibae* DFL12<sup>T</sup> and L1 medium lacking cobalamin inoculated with 100 µl *D. shibae* DFL12<sup>T</sup>.

**Figure 7** Cultures of the axenic dinoflagellate *Prorocentrum minimum* CCMP1329 (a and b) and the axenic haptophyte *Isochrysis galbana* CCMP1323 (c) inoculated with *D. shibae* DFL12<sup>T</sup> in the absence of vitamin B<sub>12</sub> and (a and c) or vitamin B<sub>1</sub> (b).

growth. Figure 7 shows the effects of cocultivation with *D. shibae* DFL12<sup>T</sup> on cultures of *P. minimum* and *I. galbana* in media lacking cobalamin or thiamine.

## Discussion

### *Adaptation to periodic anoxia*

*D. shibae* DFL12<sup>T</sup> is clearly optimized for respiratory modes of energy conservation. Its electron transport systems are fueled both by organic carbon com-

pounds and light, with ubiquinone as a central intermediate pool. However, the arginine deiminase survival fermentation pathway and the ability to use alternative electron acceptors (nitrate, DSMO) allow *D. shibae* DFL12<sup>T</sup> to sustain an active energy metabolism in the complete absence of oxygen. Several of the key genes involved are also found in other *Roseobacter* species. Therefore, the current paradigm that the *Roseobacter* clade comprises only obligate aerobes needs to be revised. Periodic anoxia is regularly encountered in photosynthetic biofilms at night as a result of intense respiration, as shown,

for example, for microbial mats (Steunou *et al.*, 2008). Moreover, steep gradients of oxygen and nitrate are present in marine sediments (Lorenzen *et al.*, 1998), and as microorganisms are easily transported by bioturbation to deeper layers (Pischedda *et al.*, 2008), the ability of *D. shibae* DFL12<sup>T</sup> to switch between alternative electron acceptors is crucial.

#### *D. shibae* DFL12<sup>T</sup> is able to provide the dinoflagellate host with vitamins

The genome data suggest that *D. shibae* DFL12<sup>T</sup> is able to synthesize two nutrients that are essential and potentially growth limiting for their hosts, vitamin B<sub>12</sub> (cobalamin) and vitamin B<sub>1</sub> (thiamine). The pathways for both of them have unique features: Altogether, three copies of the thiamine synthetase, *thiC*, are present on different plasmids (191 kb, 153 kb and 86 kb), whereas all the other thiamine synthesis genes are located on the chromosome. Synthesis of vitamin B<sub>12</sub> should be possible by an oxygen-dependent as well as an oxygen-independent pathway in *D. shibae* DFL12<sup>T</sup>, whereas organisms studied to date use only one of these pathways, pointing to both the importance of anaerobic conditions in the life of *D. shibae* DFL12<sup>T</sup> and to the importance of B<sub>12</sub>.

We experimentally confirmed that a pathway for *de novo* synthesis of B<sub>12</sub> is functional in *D. shibae* DFL12<sup>T</sup> and that the bacteria can provide both B<sub>1</sub> and B<sub>12</sub> to the dinoflagellate *P. minimum* in a defined coculture. Thus, a symbiotic relationship between *D. shibae* DFL12<sup>T</sup> and its dinoflagellate host may exist, which is based on an exchange of micronutrients synthesized at extremely low concentrations but high metabolic costs by the bacteria against photosynthate leaking from the algae during photosynthesis. As *D. shibae* is associated with several phylogenetically diverse species of algae, this symbiosis is not host specific. The dinoflagellate may replace *D. shibae* with other bacteria, provided they carry out the same essential functions, for example, synthesize B<sub>12</sub>. Conversely, *D. shibae* may live with another algae if it excretes photosynthate, dimethylsulfoniopropionate and so on. A variety of symbioses ranging from loose mutualistic interactions to obligate endosymbiotic relationships are increasingly discovered on the basis of genomic data and novel experimental methods (Moran, 2006). However, *D. shibae* DFL12<sup>T</sup> could also be a scavenger of dead algae during the collapse of algal blooms. The metabolic flexibility as shown in the genome suggests that it might even switch from a symbiotic to a parasitic mode of life.

#### Growth limitation of phytoplankton by vitamins in the sea

Although 50% of all dinoflagellate species require B<sub>12</sub>, they may not be dependent on a symbiosis with

bacteria to obtain it (Croft *et al.*, 2005), as the requirements are so low that ambient concentrations in seawater may suffice (Droop, 2007). New methods have been used to determine the pmolar concentrations of B<sub>1</sub> and B<sub>12</sub> *in situ* (Okbami and Sanudo-Wilhelmy, 2005). B<sub>12</sub> concentrations correlate with bacterial densities at certain times and in some locations and decreased 90% during dinoflagellate blooms, relative to pre-bloom values (Gobler *et al.*, 2007). B<sub>12</sub> colimitation with iron was, for the first time, shown in coastal waters of the Antarctic peninsula (Panzeca *et al.*, 2006) and later on in the Ross Sea, one of the most productive areas in the Southern Ocean (Bertrand *et al.*, 2007). In this study, the bacterial densities are low; cyanobacteria, which are able to synthesize B<sub>12</sub>, are absent, and UV irradiance is high, all accounting for low ambient levels of B<sub>12</sub>. These phytoplankton communities were iron limited, but adding B<sub>12</sub> together with iron resulted in an additional increase in biomass. One of the dominant dinoflagellates present at the time was *P. minimum*, which was shown here to be auxotrophic for B<sub>12</sub>, supporting the results of Bertrand *et al.* Thus, B<sub>12</sub> can clearly be growth limiting in the sea under certain conditions. The composition of the bacterial flora may therefore have consequences for the succession of phytoplankton species (Panzeca *et al.*, 2008). Field investigations of bacterial communities associated with algae at such a level of detail, deciphering not only phylogenetic composition but also metabolic function, have not yet been carried out, but the breathtaking progress of sequencing whole microbial communities (Nealson and Venter, 2007) and, in the future, even single cells will eventually change this.

#### Horizontal and vertical gene transfer and the adaptive *Roseobacter* gene pool

The members of the *Roseobacter* clade are characterized by a significant fraction of extrachromosomal elements in their genomes (Pradella *et al.*, 2004), with *D. shibae* DFL12<sup>T</sup> currently housing the largest number of plasmids among the fully sequenced strains. Conjugation obviously has a crucial role in the horizontal spread of these genetic elements, even if the experimental proof is still lacking. A conspicuous example is the presence of a 47-kb gene cluster on the two sister plasmids of *D. shibae* DFL12<sup>T</sup> and on several other extrachromosomal elements of the *Roseobacter* clade, which also harbor the *vir* gene cluster required for conjugation. Genes that have been found on *Roseobacter* plasmids include not only typical plasmid-encoded traits such as degradation of aromatic compounds, but also key genes for biogeochemical cycles, for example, carbon monoxide oxidation in *D. shibae* DFL12<sup>T</sup> and even the complete photosynthesis gene cluster in *R. litoralis* (Pradella *et al.*, 2004) and integrated microbial genomes). The horizontal transfer of these traits within the *Roseobacter* clade is

consistent with their patchwork distribution in the phylogenetic tree (Buchan *et al.*, 2005). Our data show that the boundary between the chromosome and the plasmid is dynamic, as outsourcing of chromosomal genes to plasmids can be observed and increases the shared adaptive gene pool in the lineage.

Besides plasmids, phages also may mediate horizontal gene transfer. In total, 5% of the genome of *S. TM1040* consists of prophages (Chen *et al.*, 2006; Moran *et al.*, 2007). A phage-like structure, the so-called GTA (gene transfer agent), was discovered many years ago in *Rh. capsulatus*. GTA particles contain only a few kilobases of DNA, are not inducible by mitomycin and do not form plaques. They contain small random pieces of host DNA, rather than a phage genome, suggesting that their main activity is the lateral transfer of host DNA. A search for the 15 GTA genes from *Rh. capsulatus* in complete and draft bacterial genomes showed that they are exclusively present in *Alphaproteobacteria*, and that all but one of the genome sequences in the *Roseobacter* clade have a GTA-like gene cluster, including *D. shibae* DFL12<sup>T</sup> (Biers *et al.*, 2008). Thus, the genetic adaptability of the *Roseobacter* gene pool appears to be because of the richness of extrachromosomal elements, some of which are conjugative, the mobilization of genes from the chromosome to the plasmid, and a phage-like gene transfer mechanism unique in *Alphaproteobacteria*.

### Outlook

*Roseobacter* bacteria are often associated with marine algae and reach high abundances during algal blooms. Our findings shed new light on this association, suggesting that it might be a symbiotic relationship in which a bacterium is required by the alga for the supply of essential nutrients, which can be growth limiting under certain conditions. Conversely, the bacteria may scavenge on dead algae and possibly switch between both types of interactions.

Studying the specificity and molecular mechanisms of this relationship in detail, both in the laboratory and in the sea, remains a fascinating field for future analyses.

### Acknowledgements

Jonathan Eisen is most gratefully acknowledged for helpful comments on an earlier version of this paper. Many thanks to the computer department at HZI for providing the facilities for the genome jamboree, and to all employees at JGI for fast and efficient feedback and support at all times. Special thanks to Tracey Riggins at the Provasoli-Guillard Center for Culture of Marine Phytoplankton (CCMP), Maine, USA, and Maike Lorenz at the Experimental Phycology and Culture Collection of Algae (SAG), Göttingen, Germany, for sending the algal cultures so

promptly! Finally, the skillful technical assistance of Bettina Elxnat is very much appreciated.

We thank the Ministry of Science and Culture (MWK), Lower Saxony, Germany and the Lower Saxonian Branch of the Volkswagen Foundation for financial support within the priority program on Marine Biotechnology (ZN1225) and within the program 'Comparative Genome Analysis of Representative Members of the Roseobacter Clade' (ZN2235). The help of the Moore Foundation, USA, is gratefully acknowledged. Sequencing and finishing of *D. shibae* DFL12<sup>T</sup> were performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program and by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48, Lawrence Berkeley National Laboratory under contract No. DE-AC03-76SF00098 and Los Alamos National Laboratory under contract No. W-7405-ENG-36.

### References

- Aasen J, Samdal IA, Miles CO, Dahl E, Briggs LR, Aune T. (2005). Yessotoxins in Norwegian blue mussels (*Mytilus edulis*): uptake from *Protoceratium reticulatum*, metabolism and depuration. *Toxicon* **45**: 265–272.
- Alavi M, Miller T, Erlandson K, Schneider R, Belas R. (2001). Bacterial community associated with *Pfiesteria*-like dinoflagellate cultures. *Environ Microbiol* **3**: 380–396.
- Allgaier M, Uphoff H, Felske A, Wagner-Döbler I. (2003). Aerobic anoxygenic photosynthesis in *Roseobacter* clade bacteria from diverse marine habitats. *Appl Environ Microbiol* **69**: 5051–5059.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403–410.
- Aluwihare LI, Repeta DJ, Chen RF. (2002). Chemical composition and cycling of dissolved organic matter in the Mid-Atlantic bight. *Deep Sea Res Part II Top Stud Oceanogr* **49**: 4421–4437.
- Azam F, Fenchel T, Gray JS, Meyer-Reill LA, Thingstad F. (1983). The ecological role of water-column microbes in the sea. *Mar Ecol Prog Ser* **10**: 257–263.
- Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S *et al.* (2007). CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **315**: 1709–1712.
- Bertrand EM, Saito MA, Rose JM, Riesselman CR, Lohan MC, Noble AE *et al.* (2007). Vitamin B-12 and iron colimitation of phytoplankton growth in the Ross sea. *Limnol Oceanogr* **52**: 1079–1093.
- Biebl H, Allgaier M, Tindall BJ, Koblizek M, Luensdorf H, Pukall R *et al.* (2005). *Dinoroseobacter shibae* gen. nov., sp. nov., a new aerobic phototrophic bacterium isolated from dinoflagellates. *Int J Syst Evol Microbiol* **55**: 1089–1096.
- Biebl H, Wagner-Döbler I. (2006). Growth and bacteriochlorophyll *a* formation in taxonomically diverse aerobic anoxygenic phototrophic bacteria in chemostat culture: influence of light regimen and starvation. *Process Biochem* **41**: 2153–2159.
- Biers EJ, Wang K, Pennington C, Belas R, Chen F, Moran MA. (2008). Occurrence and expression of

- gene transfer agent genes in marine bacterioplankton. *Appl Environ Microbiol* **74**: 2933–2939.
- Biersmith A, Benner R. (1998). Carbohydrates in phytoplankton and freshly produced dissolved organic matter. *Mar Chem* **63**: 131–144.
- Bland C, Ramsey TL, Sabree F, Lowe M, Brown K, Kyrpides NC et al. (2007). CRISPR recognition tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. *BMC Bioinformatics* **8**: 209.
- Brinkhoff T, Bach G, Heidorn T, Liang L, Schlingloff A, Simon M. (2004). Antibiotic production by a *Roseobacter* clade-affiliated species from the German Wadden Sea and its antagonistic effects on indigenous isolates. *Appl Environ Microbiol* **70**: 2560–2565.
- Brinkhoff T, Giebel HA, Simon M. (2008). Diversity, ecology, and genomics of the *Roseobacter* clade: a short overview. *Arch Microbiol* **189**: 531–539.
- Bruhn JB, Nielsen KF, Hjelm M, Hansen M, Bresciani J, Schulz S et al. (2005). Ecology, inhibitory activity, and morphogenesis of a marine antagonistic bacterium belonging to the *Roseobacter* clade. *Appl Environ Microbiol* **71**: 7263–7270.
- Brussaard CP, Wilhelm SW, Thingstad F, Weinbauer MG, Bratbak G, Heldal M et al. (2008). Global-scale processes with a nanoscale drive: the role of marine viruses. *ISME J* **2**: 575–578.
- Buchan A, Gonzalez JM, Moran MA. (2005). Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* **71**: 5665–5677.
- Case RJ, Labbate M, Kjelleberg S. (2008). AHL-driven quorum-sensing circuits: their frequency and function among the Proteobacteria. *ISME J* **2**: 345–349.
- Chen F, Wang K, Stewart J, Belas R. (2006). Induction of multiple prophages from a marine bacterium: a genomic approach. *Appl Environ Microbiol* **72**: 4995–5001.
- Choudhary M, Zanhua X, Fu YX, Kaplan S. (2007). Genome analyses of three strains of *Rhodobacter sphaeroides*: evidence of rapid evolution of chromosome II. *J Bacteriol* **189**: 1914–1921.
- Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. (2005). Algae acquire vitamin B12 through a symbiotic relationship with bacteria. *Nature* **438**: 90–93.
- Danovaro R, Dell'Anno A, Corinaldesi C, Magagnini M, Noble R, Tamburini C et al. (2008). Major viral impact on the functioning of benthic deep-sea ecosystems. *Nature* **454**: 1084–1087.
- Droop MR. (2007). Vitamins, phytoplankton and bacteria: symbiosis or scavenging? *J Plankton Res* **29**: 107–113.
- Eschbach M, Schreiber K, Trunk K, Buer J, Jahn D, Schobert M. (2004). Long-term anaerobic survival of the opportunistic pathogen *Pseudomonas aeruginosa* via pyruvate fermentation. *J Bacteriol* **186**: 4596–4604.
- Ewing B, Green P. (1998). Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res* **8**: 186–194.
- Ewing B, Hillier L, Wendl MC, Green P. (1998). Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res* **8**: 175–185.
- Forward JA, Behrendt MC, Wyborn NR, Cross R, Kelly DJ. (1997). TRAP transporters: a new family of periplasmic solute transport systems encoded by the *dctPQM* genes of *Rhodobacter capsulatus* and by homologs in diverse gram-negative bacteria. *J Bacteriol* **179**: 5482–5493.
- Fuhrman JA. (1999). Marine viruses and their biogeochemical and ecological effects. *Nature* **399**: 541–548.
- Fürch T, Preusse M, Tomasch J, Zech H, Wagner-Döbler I, Rabus R et al. (2009). Metabolic fluxes in the central carbon metabolism of *Dinoroseobacter shibae* and *Phaeobacter gallaeciensis*, to members of the marine *Roseobacter* clade. submitted to BMC Microbiology.
- Geng H, Bruhn JB, Nielsen KF, Gram L, Belas R. (2008). Genetic dissection of tropodithietic acid biosynthesis by marine roseobacters. *Appl Environ Microbiol* **74**: 1535–1545.
- Gevers D, Cohan FM, Lawrence JG, Spratt BG, Coenye T, Feil EJ et al. (2005). Opinion: re-evaluating prokaryotic species. *Nat Rev Microbiol* **3**: 733–739.
- Gobler CJ, Norman C, Panzeca C, Taylor GT, Sanudo-Wilhelmy SA. (2007). Effect of B-vitamins (B1, B12) and inorganic nutrients on algal bloom dynamics in a coastal ecosystem. *Aquat Microb Ecol* **49**: 181–194.
- Godde JS, Bickerton A. (2006). The repetitive DNA elements called CRISPRs and their associated genes: evidence of horizontal transfer among prokaryotes. *J Mol Evol* **62**: 718–729.
- Gordon D, Abajian C, Green P. (1998). Consed: a graphical tool for sequence finishing. *Genome Res* **8**: 195–202.
- Grissa I, Vergnaud G, Pourcel C. (2007). The CRISPRdb database and tools to display CRISPRs and to generate dictionaries of spacers and repeats. *BMC Bioinformatics* **8**: 172.
- Guillard RRL, Hargraves PE. (1993). *Stichochrysis immobilis* is a diatom, not a chrysophyte. *Phycologia* **32**: 234–236.
- Haft DH, Selengut J, Mongodin EF, Nelson KE. (2005). A guild of 45 CRISPR-associated (Cas) protein families and multiple CRISPR/Cas subtypes exist in prokaryotic genomes. *PLoS Comput Biol* **1**: e60.
- Han CS, Chain P. (2006). Finishing repeat regions automatically with Dupfinisher Arabnia, H. R. and Valafar H. 141–146. CSREA Press. Proceedings of the 2006 International Conference on Bioinformatics & Computational Biology. 26-6-2006.
- Hasegawa Y, Martin JL, Giewat MW, Rooney-Varga JN. (2007). Microbial community diversity in the phycosphere of natural populations of the toxic alga, *Alexandrium fundyense*. *Environ Microbiol* **9**: 3108–3121.
- Hoang HH, Gurich N, lez JE. (2008). Regulation of motility by the ExpR/Sin quorum-sensing system in *Sinorhizobium meliloti*. *J Bacteriol* **190**: 861–871.
- Howard EC, Henriksen JR, Buchan A, Reisch CR, Burgmann H, Welsh R et al. (2006). Bacterial taxa that limit sulfur flux from the ocean. *Science* **314**: 649–652.
- Kobryn K, Chaconas G. (2001). The circle is broken: telomere resolution in linear replicons. *Curr Opin Microbiol* **4**: 558–564.
- Kowalik K. (1969). The crystal lattice of the pyrenoid matrix of *Prorocentrum micans*. *J Cell Sci* **5**: 251–269.
- Lafay B, Ruimy R, de Traubenberg CR, Breittmayer V, Gauthier MJ, Christen R. (1995). *Roseobacter algicola* sp. nov., a new marine bacterium isolated from the phycosphere of the toxin-producing dinoflagellate *Prorocentrum lima*. *Int J Syst Bacteriol* **45**: 290–296.
- Levasseur M, Couture JY, Weise AM, Michaud S, Elbraechter M, Sauve G et al. (2003). Pelagic and epiphytic summer distributions of *Prorocentrum lima* and *P. mexicanum* at two mussel farms in the Gulf of St. Lawrence, Canada. *Aquat Microb Ecol* **30**: 283–293.
- Lorenzen J, Larsen LH, Kjaer T, Revsbech NP. (1998). Biosensor determination of the microscale distribution

- of nitrate, nitrate assimilation, nitrification, and denitrification in a diatom-inhabited freshwater sediment. *Appl Environ Microbiol* **64**: 3264–3269.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar *et al.* (2004). ARB: a software environment for sequence data. *Nucleic Acids Res* **32**: 1363–1371.
- Makarova KS, Grishin NV, Shabalina SA, Wolf YI, Koonin EV. (2006). A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. *Biol Direct* **1**: 7.
- Marketon MM, Gronquist MR, Eberhard A, Iez JE. (2002). Characterization of the *Sinorhizobium meliloti* sinR/sinI locus and the production of novel N-acyl homoserine lactones. *J Bacteriol* **184**: 5686–5695.
- Markowitz VM, Szeto E, Palaniappan K, Grechkin Y, Chu K, Chen IM *et al.* (2008). The integrated microbial genomes (IMG) system in 2007: data content and analysis tool extensions. *Nucleic Acids Res* **36**: D528–D533.
- Martens JH, Barg H, Warren MJ, Jahn D. (2002). Microbial production of vitamin B12. *Appl Microbiol Biotechnol* **58**: 275–285.
- Mayali X, Franks PJ, Azam F. (2008). Cultivation and ecosystem role of a marine *Roseobacter* clade-affiliated cluster bacterium. *Appl Environ Microbiol* **74**: 2595–2603.
- McGoldrick HM, Roessner CA, Raux E, Lawrence AD, McLean KJ, Munro AW *et al.* (2005). Identification and characterization of a novel vitamin B12 (cobalamin) biosynthetic enzyme (CobZ) from *Rhodobacter capsulatus*, containing flavin, heme, and Fe-S cofactors. *J Biol Chem* **280**: 1086–1094.
- Meinhardt F, Schaffrath R, Larsen M. (1997). Microbial linear plasmids. *Appl Microbiol Biotechnol* **47**: 329–336.
- Miller TR, Belas R. (2004). Dimethylsulfoniopropionate metabolism by *Pfiesteria*-associated *Roseobacter* spp. *Appl Environ Microbiol* **70**: 3383–3391.
- Miller TR, Belas R. (2006). Motility is involved in *Silicibacter* sp. TM 1040 interaction with dinoflagellates. *Environ Microbiol* **8**: 1648–1659.
- Miller TR, Hnilicka K, Dziedzic A, Desplats P, Belas R. (2004). Chemotaxis of *Silicibacter* sp. strain TM1040 toward dinoflagellate products. *Appl Environ Microbiol* **70**: 4692–4701.
- Moran MA, Belas R, Schell MA, Gonzalez JM, Sun F, Sun S *et al.* (2007). Ecological genomics of marine *Roseobacters*. *Appl Environ Microbiol* **73**: 4559–4569.
- Moran MA, Buchan A, Gonzalez JM, Heidelberg JF, Whitman WB, Kiene RP *et al.* (2004). Genome sequence of *Silicibacter pomeroyi* reveals adaptations to the marine environment. *Nature* **432**: 910–913.
- Moran MA, Miller WL. (2007). Resourceful heterotrophs make the most of light in the coastal ocean. *Nat Rev Microbiol* **5**: 792–800.
- Moran NA. (2006). Symbiosis. *Curr Biol* **16**: R866–R871.
- Muhling M, Fuller NJ, Millard A, Somerfield PJ, Marie D, Wilson WH *et al.* (2005). Genetic diversity of marine *Synechococcus* and co-occurring cyanophage communities: evidence for viral control of phytoplankton. *Environ Microbiol* **7**: 499–508.
- Mulligan C, Kelly DJ, Thomas GH. (2007). Tripartite ATP-independent periplasmic transporters: application of a relational database for genome-wide analysis of transporter gene frequency and organization. *J Mol Microbiol Biotechnol* **12**: 218–226.
- Munch R, Hiller K, Grote A, Scheer M, Klein J, Schobert M *et al.* (2005). Virtual Footprint and PRODORIC: an integrative framework for regulon prediction in prokaryotes. *Bioinformatics* **21**: 4187–4189.
- Nagasaki K, Takao Y, Shirai Y, Mizumoto H, Tomaru Y. (2005). Molecular ecology of microalgal viruses. *Virus* **55**: 127–132.
- Nealson KH, Venter JC. (2007). Metagenomics and the global ocean survey: what's in it for us, and why should we care? *ISME J* **1**: 185–187.
- Okbami Michael M, Sanudo-Wilhelmy SA. (2005). Direct determination of vitamin B1 in seawater by solid-phase extraction and high-performance liquid chromatography quantification. *Limnol Oceanogr Methods* **3**: 241–246.
- Okolodkov YB, Campos-Bautista G-L, Gonzalez-Gonzalez JAG, Hoppenrath M, Arenas V. (2007). Seasonal changes of benthic and epiphytic dinoflagellates in the Veracruz reef zone, Gulf of Mexico. *Aquat Microb Ecol* **47**: 223–237.
- Pan Y, Cembella AD, Quilliam MA. (1999). Cell cycle and toxin production in the benthic dinoflagellate *Prorocentrum lima*. *Mar Biol* **134**: 541–549.
- Panzeca C, Beck AJ, Leblanc K, Taylor GT, Hutchins DA, Sanudo-Wilhelmy SA. (2008). Potential cobalt limitation of vitamin B12 synthesis in the North Atlantic Ocean. *Global Biogeochem Cycles* **22**: GB2029, doi:10.29/2007GB003124.
- Panzeca C, Tovar-Sanchez A, Agusti S, Reche I, Duarte CM, Taylor GT *et al.* (2006). B vitamins as regulators of phytoplankton dynamics. *Eos* **87**: 593, 596.
- Parsons ML, Preskitt LB. (2007). A survey of epiphytic dinoflagellates from the coastal waters of the island of Hawai'i. *Harmful Algae* **6**: 658–669.
- Pernthaler A, Pernthaler J, Amann R. (2002). Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. *Appl Environ Microbiol* **68**: 3094–3101.
- Petersen J, Brinkmann H, Pradella S. (2009). Diversity and evolution of *repABC* type plasmids in *Rhodobacteriales*. *Environ Microbiol* **1462**–2920.
- Pischedda L, Poggiale JC, Cuny P, Gilbert F. (2008). Imaging oxygen distribution in marine sediments. The importance of bioturbation and sediment heterogeneity. *Acta Biotheor* **56**: 123–135.
- Pradella S, Allgaier M, Hoch C, Paeuker O, Stackebrandt E, Wagner-Döbler I. (2004). Genome organization and localization of the *pufLM* genes of the photosynthesis reaction center in phylogenetically diverse marine *Alphaproteobacteria*. *Appl Environ Microbiol* **70**: 3360–3369.
- Prokic I, Bruemmer F, Brigge T, Goertz H, Gerdt G, Schuett C *et al.* (1998). Bacteria of the Genus *Roseobacter* Associated with the Toxic Dinoflagellate *Prorocentrum lima*. *Protist* **149**: 347–357.
- Rabus R, Jack DL, Kelly DJ, Saier Jr MH. (1999). TRAP transporters: an ancient family of extracytoplasmic solute-receptor-dependent secondary active transporters. *Microbiology* **145**: 3431–3445.
- Rhodes CJ, Truscott JE, Martin AP. (2008). Viral infection as a regulator of oceanic phytoplankton populations. *J Mar Syst* **74**: 216–226.
- Rodionov DA, Vitreschak AG, Mironov AA, Gelfand MS. (2003). Comparative genomics of the vitamin B12 metabolism and regulation in prokaryotes. *J Biol Chem* **278**: 41148–41159.



- Rossello-Mora R, Amann R. (2001). The species concept for prokaryotes. *FEMS Microbiol Rev* **25**: 39–67.
- Roth JR, Lawrence JG, Bobik TA. (1996). Cobalamin (coenzyme B12): synthesis and biological significance. *Annu Rev Microbiol* **50**: 137–181.
- Scott AI, Roessner CA. (2002). Biosynthesis of cobalamin (vitamin B12). *Biochem Soc Trans* **30**: 613–620.
- Sekar R, Pernthaler A, Pernthaler J, Warnecke F, Posch T, Amann R. (2003). An improved protocol for quantification of freshwater *Actinobacteria* by fluorescence in situ hybridization. *Appl Environ Microbiol* **69**: 2928–2935.
- Selje N, Simon M, Brinkhoff T. (2004). A newly discovered *Roseobacter* cluster in temperate and polar oceans. *Nature* **427**: 445–448.
- Seshadri R, Kravitz SA, Smarr L, Gilna P, Frazier M. (2007). CAMERA: a community resource for metagenomics. *PLoS Biol* **5**: e75.
- Shiba T. (1991). *Roseobacter litoralis* gen. nov., sp. nov., and *Roseobacter denitrificans* sp. nov., aerobic pink-pigmented bacteria which contain bacteriochlorophyll a. *Syst Appl Microbiol* **14**: 140–145.
- Stackebrandt E, Frederiksen W, Garrity GM, Grimont PA, Kampf P, Maiden MC *et al*. (2002). Report of the *ad hoc* committee for the re-evaluation of the species definition in bacteriology. *Int. J Syst Evol Microbiol* **52**: 1043–1047.
- Steunou AS, Jensen SI, Brecht E, Becraft ED, Bateson MM, Kilian O *et al*. (2008). Regulation of *nif* gene expression and the energetics of N<sub>2</sub> fixation over the diel cycle in a hot spring microbial mat. *ISME J* **2**: 364–378.
- Swingley WD, Sadekar S, Mastrian SD, Matthies HJ, Hao J, Ramos H *et al*. (2007). The complete genome sequence of *Roseobacter denitrificans* reveals a mixotrophic rather than photosynthetic metabolism. *J Bacteriol* **189**: 683–690.
- Tomaru Y, Nagasaki K. (2004). Widespread occurrence of viruses lytic to the bivalve-killing dinoflagellate *Heterocapsa circularisquama* along the western coast of Japan. *Plankton Biol Ecol* **51**: 1–6.
- Tyson GW, Banfield JF. (2008). Rapidly evolving CRISPRs implicated in acquired resistance of microorganisms to viruses. *Environ Microbiol* **10**: 200–207.
- Vander Horn PB, Backstrom AD, Stewart V, Begley TP. (1993). Structural genes for thiamine biosynthetic enzymes (thiCEFGH) in *Escherichia coli* K-12. *J Bacteriol* **175**: 982–992.
- Vershinin AO, Moruchkov AA, Leighfield T, Sukhanova IN, Pan'kov SL, Morton SL *et al*. (2005). Potentially toxic algae in the coastal phytoplankton of the north-east Black Sea in 2001–2002. *Oceanology* **45**: 224–232.
- Wagner-Döbler I, Biebl H. (2006). Environmental biology of the marine *Roseobacter* lineage. *Annu Rev Microbiol* **60**: 255–280.
- Wagner-Döbler I, Thiel V, Eberl L, Allgaier M, Bodor A, Meyer S *et al*. (2005). Discovery of complex mixtures of novel long-chain quorum sensing signals in free-living and host-associated marine *Alphaproteobacteria*. *Chembiochem* **6**: 2195–2206.
- Wang JT, Douglas AE. (1997). Nutrients, signals, and photosynthate release by symbiotic algae: the impact of taurine on the dinoflagellate alga *Symbiodinium* from the sea anemone *Aiptasia pulchella*. *Plant Physiol* **114**: 631–636.
- Warren MJ, Raux E, Schubert HL, Escalante-Semerena JC. (2002). The biosynthesis of adenosylcobalamin (vitamin B12). *Nat Prod Rep* **19**: 390–412.
- Weinbauer MG. (2004). Ecology of prokaryotic viruses. *FEMS Microbiol Rev* **28**: 127–181.
- Williams HD, Zlosnik JE, Ryall B. (2007). Oxygen, cyanide and energy generation in the cystic fibrosis pathogen *Pseudomonas aeruginosa*. *Adv Microb Physiol* **52**: 1–71–1–71.
- Williamson SJ, Rusch DB, Yooseph S, Halpern AL, Heidelberg KB, Glass JI *et al*. (2008). The Sorcerer II Global Ocean Sampling Expedition: metagenomic characterization of viruses within aquatic microbial samples. *PLoS ONE* **3**: e1456.
- Yoch DC. (2002). Dimethylsulfoniopropionate: its sources, role in the marine food web, and biological degradation to dimethylsulfide. *Appl Environ Microbiol* **68**: 5804–5815.
- Zielenkiewicz U, Ceglowski P. (2001). Mechanisms of plasmid stable maintenance with special focus on plasmid addiction systems. *Acta Biochim Pol* **48**: 1003–1023.

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)