Tissue-specific evaluation of suitable reference genes for RT-qPCR in the pond snail, *Lymnaea stagnalis*

Alexander P. Young¹, Carmen F. Landry¹, Daniel J. Jackson² and Russell C. Wyeth¹

¹ Department of Biology, St. Francis Xavier University, Antigonish, NS, Canada

² Department of Geobiology, Georg-August Universität Göttingen, Göttingen, Germany

ABSTRACT

Reverse transcription quantitative PCR (RT-qPCR) is a robust technique for the quantification and comparison of gene expression. To obtain reliable results with this method, one or more reference genes must be employed to normalize expression measurements among treatments or tissue samples. Candidate reference genes must be validated to ensure that they are stable prior to use in qPCR experiments. The pond snail (*Lymnaea stagnalis*) is a common research organism, particularly in the areas of learning and memory, and is an emerging model for the study of biological asymmetry, biomineralization, and evolution and development. However, no systematic assessment of qPCR reference genes has been performed in this animal. Therefore, the aim of our research was to identify stable reference genes to normalize gene expression data from several commonly studied tissues in *L. stagnalis* as well as across the entire body. We evaluated a panel of seven reference genes across six different tissues in *L. stagnalis* with RT-qPCR. The genes included: *elongation factor 1-alpha*, *glyceraldehyde-3-phosphate dehydrogenase*, *beta-actin*, *beta-tubulin*, *ubiquitin*, *prenylated rab acceptor protein 1*, and a voltage gated potassium channel. These genes exhibited a wide range of expression levels among tissues. The tissue-specific stability of each of the genes was consistent when measured by the standard stability assessment algorithms: geNorm, NormFinder, BestKeeper, and RefFinder. Our data indicate that the most stable reference genes vary among the tissues that we examined (central nervous system, tentacles, lips, penis, foot, mantle). Our results were generally congruent with those obtained from similar studies in other molluscs. Given that a minimum of two reference genes are recommended for data normalization, we provide suggestions for strong pairs of reference genes for single- and multi-tissue analyses of RT-qPCR data in *L. stagnalis*.

Submitted 29 April 2019

Accepted 13 September 2019

Published 15 October 2019

Corresponding author

Alexander P. Young, ayoung@stfx.ca

Academic editor

Björn Brembs

Additional Information and
Declarations can be found on
page 12

DOI [10.7717/peerj.7888](https://doi.org/10.7717/peerj.7888)

© Copyright
2019 Young et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Aquaculture, Fisheries and Fish Science, Genetics, Molecular Biology, Neuroscience

Keywords Gene expression, Normalization, Mollusc, Gastropod neurobiology

BACKGROUND

Lymnaea stagnalis is a freshwater snail with an extensive history of proximate mechanism research. Previous studies have primarily focused on its neurobiology, as well as its endocrinology, immunology, and shell morphology (Chase, 2002; Benjamin, 2008). The central nervous system (CNS) is relatively simple, with approximately 20,000 neurons,

many of which are large and easily identifiable, making them amenable for studies of learning, memory, motor pattern generation, neuronal regeneration, and synapse formation (Lukowiak, 2016; Elliott & Susswein, 2002; Chase, 2002; Lukowiak et al., 2003; Getz et al., 2018). Parallel investigations into the neuroendocrine (Koene, 2010; Pirger et al., 2010) and immune (Van Der Knaap, Adema & Sminia, 1993; Gust et al., 2013a) systems of *L. stagnalis* and other gastropods have improved our understanding of reproductive behavior, immunity and toxicology of *L. stagnalis*. Additionally, *L. stagnalis* has been used as a research organism to study the molecular mechanisms that guide shell formation (Boer & Witteveen, 1980; Ebanks, O'Donnell & Grosell, 2010; Hohagen & Jackson, 2013; Jackson, Herlitze & Hohagen, 2016; Herlitze et al., 2018). More recently, *L. stagnalis* has also been used to study the molecular basis of body asymmetry (Davison et al., 2016; Abe & Kuroda, 2019).

A broad range of traditional techniques have been used in past studies, but modern molecular genetics have yet to be thoroughly incorporated into the study of *L. stagnalis*. Methods to study the genetics of *L. stagnalis* such as reverse transcription quantitative PCR (RT-qPCR), in situ hybridization, and transcriptomics (among others) have been established but have not yet been used extensively (Feng et al., 2009; Herlitze et al., 2018). The few studies using RT-qPCR have spanned several topics including chirality, learning, and memory (Sadamoto et al., 2010; Foster, Lukowiak & Henry, 2015; Davison et al., 2016; Korneev et al., 2018; Dong et al., 2018). Additionally, a small number of studies have used in situ hybridization in this species, primarily in sections and whole mounts of the CNS, as well as larvae (Dirks et al., 1989; Boer et al., 1992; Croll & Van Minnen, 1992; Sadamoto et al., 2004). Our goal here is to continue to build the foundation for further molecular genetics studies in *L. stagnalis* via validation of candidate reference genes for RT-qPCR.

RT-qPCR is a robust technique for the quantification of the mRNA for a specific transcript. It can be used alongside in situ hybridization or other visualization techniques to establish patterns of gene expression in animals. For such experiments, relative quantification against one or more reference genes is the most common choice to compare gene expression across samples (Schmittgen & Livak, 2008; Ruijter et al., 2009). Historically, highly expressed cellular maintenance genes have been haphazardly selected as reference genes for qPCR experiments in many species because expression of such genes was thought to be inherently stable (Kozera & Rapacz, 2013). However, many of these genes have been shown to have unstable expression in several species of plants and animals, highlighting the importance of species-specific validation prior to use (Dheda et al., 2004; Barber et al., 2005; De Jonge et al., 2007; Tong et al., 2009; Eissa et al., 2016). Therefore, validation that candidate reference genes have stable expression is critical for the acquisition of accurate qPCR data and the experimental approaches that incorporate this technique.

To our knowledge, the only previous example of reference gene validation in *L. stagnalis* is for *EF1α* in the CNS (Foster, Lukowiak & Henry, 2015). Other experiments have used reference genes without validation, including *elongation factor 1-alpha* (*EF1α*) (Gust et al., 2013b; Shimizu et al., 2013), *glyceraldehyde-3-phosphate dehydrogenase*

(GAPDH) (Aleksic & Feng, 2012), beta-actin (ACTB) (Senatore & Spafford, 2012; Hatakeyama et al., 2013; Carter et al., 2015) and beta-tubulin (TUBB) (Bavan et al., 2012; Korneev et al., 2013; Gust et al., 2013b; Flynn et al., 2014; Carter et al., 2015; Benatti et al., 2017). To establish a more rigorous foundation for future research using RT-qPCR, we present an analysis of seven candidate reference genes across six tissues of interest (tentacles, lips, foot, penis, mantle, and CNS) in *L. stagnalis*. We also provide a combined analysis with data from all tissues to demonstrate which genes are likely to be most stably expressed in whole-body preparations of *L. stagnalis*.

We investigated the stability of *Lst-EF1α*, *Lst-GAPDH*, *Lst-ACTB*, *Lst-TUBB*, and *Lst-ubiquitin* (*UBI*) as they are commonly employed reference genes. Messenger RNA transcripts encoding the prenylated rab acceptor protein 1 (*Rapac1*) and a voltage gated potassium channel (*VGKC*) were also assessed as analogs of these genes have recently been found to be stable in the terrestrial gastropod *Cepaea nemoralis* (Affenzeller, Cerveau & Jackson, 2018).

METHODS

Care of snails

Animal use was consistent with the Canadian Council for Animal Care guidelines. A colony of *L. stagnalis* was bred and maintained in the animal care facility at StFX. The animals were exposed to a photoperiod matched to natural daylight patterns in Antigonish, Nova Scotia. The water in the animals' tanks was changed three times per week. The animals were fed fish food and romaine lettuce ad libitum plus sinking protein pellets once per week.

Total RNA extraction

Total RNA was collected from each *L. stagnalis* tissue of interest. To prepare for RNA isolation, animals were anesthetized in 0.125% 1-phenoxy-2-propanol dissolved in *Lymnaea* saline for up to 30 min (Wyeth et al., 2009). The animals were dissected to isolate the six tissues of interest: CNS, tentacles, lips, penis, foot, and mantle. The tissues of each type from two animals were combined for each replicate to ensure adequate yields of RNA from each tissue, three replicates were produced with this method. Tissues were flash frozen in liquid nitrogen, shredded with razor blades and added to 500 µL of TRIzol reagent (Thermo Fisher, Waltham, MA, USA). Tissue solutions were thoroughly homogenized with a PowerGen 125 tissue homogenizer (Thermo Fisher, Waltham, MA, USA). Total RNA was extracted from the homogenized tissue via addition of 0.5 volumes of a 24:1 solution of chloroform and isoamyl alcohol. The aqueous layer was separated, and RNA was precipitated with isopropyl alcohol. RNA precipitate was transferred to the spin column of the E.Z.N.A.[®] Mollusc RNA Kit (Omega Bio-Tek, Norcross, GA, USA) where the RNA was washed and subjected to the on-column DNase I treatment as per the manufacturer's instructions. To confirm the effectiveness of the DNase I treatment, PCR was performed with primer sets for *Lst-tyrosine hydroxylase* (forward: 5'-CCCACG TGTATCGTCATCTTA-3'; reverse: 5'-ATCTTCTCCTCGCAAACC-3') and *Lst-EF1α* with 20 ng of total RNA as a template as this reaction would not work in the absence

of genomic DNA (gDNA) contamination. *Lst-tyrosine hydroxylase* was chosen as one of the gDNA contamination controls as it is frequently amplified by us in all of the tissues of interest as a part of a larger research project on neural development. No amplification was observed for either gene in any of the RNA samples after 40 cycles, thus the RNA samples were judged to be free of gDNA contamination.

Total RNA was quantified in a QuBit 3.0 (Thermo Fisher, Waltham, MA, USA). Yields of 50–200 ng/μL dissolved in 30 μL of molecular grade water were common for roughly 40 mg of tissue mass. A sample of total RNA was measured with a spectrophotometer to confirm a 260:280 ratio of 1.8–2.0. Lastly, five μL of each RNA sample was denatured in five μL of 2× RNA Loading Dye (New England Biolabs, Ipswich, MA, USA) at 75 °C for 10 min and then immediately placed on ice. The RNA samples were loaded on a 1% agarose non-denaturing gel and run alongside a single stranded RNA ladder (New England Biolabs, Ipswich, MA, USA) to verify that the 28S and 18S rRNA bands were intact and the RNA samples were not degraded.

Reverse transcription PCR

Reverse transcription-PCR was performed with iScript Reverse Transcription Supermix for RT-qPCR (Bio-Rad Laboratories, Hercules, CA, USA) which contains a mixture of oligo(dT) primers and random hexamers. For RT-PCR reactions, 500 ng of RNA was added to four μL of iScript Supermix and topped up to 20 μL with molecular grade water as recommended by the manufacturer. RT-PCR took place in a Bio-Rad CFX Connect thermocycler (Bio-Rad Laboratories, Hercules, CA, USA). The RT-PCR program ran at 25 °C for 5 min, 46 °C for 20 min and 95 °C for 1 min.

Design, validation, and optimization of primers for quantitative PCR

Candidate primers intended for qPCR were designed with CLC Main Workbench software (Qiagen, Hilden, Germany). Primers were designed to have a length of 19–23 nucleotides, a melting temperature between 55 and 62 °C, a GC content between 40% and 60% and produce a product no more than 200 bp long. Additionally, the GC content of each primer was kept within 10% of its counterpart, and the melting temperature of each was kept within 3 °C of its counterpart.

Two to six sets of primers were designed for each gene so that optimal primers could be selected for the qPCR reactions. The primers were tested on combined cDNA samples to minimize tissue-specific bias. Melt curves were performed to verify that one product was amplified. If the primers produced a single product, then five μL of the PCR reactions were run on a 2% (w/v) agarose gel at 60 V for roughly 30 min alongside a 50 bp DNA Ladder (New England BioLabs, Ipswich, MA, USA). The agarose gels were analyzed with a Bio-Rad ChemiDoc (Bio-Rad Laboratories, Hercules, CA, USA) to visualize the size of the product. Ultimately, one set of primers was selected for each candidate reference gene based on the quality of the PCR product. The primers for each candidate reference gene are listed in [Table 1](#).

After the optimal primer sets were selected based on reaction specificity, primer efficiencies were calculated for each primer set. Primer efficiency curves were generated

Table 1 Description of all primers used to amplify candidate reference genes in *Lymnaea stagnalis*. Primers labeled “For” are forward primers and primers labeled “Rev” are reverse primers, all sequences are written in the 5’ to 3’ direction. The amplification efficiency was determined from each reference gene primer set following RT-qPCR with five 1:5 serial dilutions of total RNA from 100 ng.

Gene	Primer Sequence (5’ to 3’)	Function	Product Size (bp)	Tm (°C)	Efficiency	r ²	Accession
<i>Lst-ACTB</i>	For [AGGCCAACAGAGAAAAGA] Rev [AGATGCGTACAGAGAGAG]	Cell structure and motility	97	56	2.12	0.999	KX387883
<i>Lst EF1α</i>	For [ACCACAACCTGGCCACTTGATC] Rev [CCATCTCTTGGGCCTCTTTCT]	Delivery of tRNA for protein synthesis	85	59	2.00	0.998	MH687364
<i>Lst-GAPDH</i>	For [CAACAACCGACAAAGCAA] Rev [CATAACAAACATAGGGGCA]	Carbohydrate metabolism	93	55	1.82	0.988	MH687363
<i>Lst-Rapac1</i>	For [GGCTCTTTCTTTCCCTTTGT] Rev [TTCCTGCTCTTCTTGCGT]	Cellular trafficking	124	58	1.82	0.989	MH687365
<i>Lst-TUBB</i>	For [GGCTAGGGGATGAAGATGA] Rev [AGGATGAGGGTGAATTTGA]	Microtubule element—cell structure	130	56	1.80	0.993	KX387887
<i>Lst-UBI</i>	For [GTATTGTGGTGCTGGTGTGTTT] Rev [GCTTCCTCCTCTGGTTTGT]	Regulate protein function	105	59	1.94	0.993	MH687367
<i>Lst-VGKC</i>	For [TGGCTTCCTGCTTCTCTGT] Rev [GCTTCTGTCGTTGTTTTGCT]	Maintenance of cell membrane potential	99	60	1.82	0.997	MH687366

from RT-qPCR reactions on serial dilutions of RNA. The parent reaction contained 100 ng total RNA and four 1:5 dilutions were performed to generate five Cq values.

Quantitative PCR

The minimum information for publication of quantitative real-time (MIQE) PCR experiments guidelines were followed throughout the collection of qPCR data ([Bustin et al., 2009](#)). Amplification of all genes was detected with SyBR Green dye which generates fluorescence based on the synthesis of double-stranded DNA. The reactions contained two µL of cDNA with 10 µL of Bio-Rad SsoAdvanced Universal SyBR Mix, 600 nM forward and reverse primer concentration, and topped to 20 µL with DEPC H₂O. Each replicate of *L. stagnalis* tissue was subjected to qPCR reactions in triplicate. The qPCR reactions took place in a Bio-Rad CFX Connect thermocycler running a custom program. The custom qPCR program consisted of 95 °C for 30 s; 40 cycles of 95 °C for 15 s, 55 °C for 30 s. The plate was read by the machine to measure fluorescence at the end of each cycle.

Data analysis

The expression stability of each gene was assessed with four computational algorithms: geNorm ([Vandesompele et al., 2002](#)), NormFinder ([Andersen, Jensen & Ørntoft, 2004](#)), BestKeeper ([Pfaffl et al., 2004](#)), and RefFinder ([Xie et al., 2012](#)). GeNorm ranks candidates by their expression stability (*M*) values that are assigned following pair-wise variation measurements among genes. Genes with *M* < 1.5 are considered to be stable. The NormFinder algorithm compares intra-group variation (i.e., mRNA levels of one gene within the tissue of interest) to inter-group variation (i.e., mRNA levels of other genes in the same tissues) and assigns genes a stability value based on variation among Cq values. BestKeeper builds a stability index based on repeated pair-wise correlation analyses

Figure 1 Representative image of PCR products for each reference gene. A five μ L sample of each PCR product was run on a 2% agarose gel that also contained a no primer control (NPC).

Full-size [DOI: 10.7717/peerj.7888/fig-1](https://doi.org/10.7717/peerj.7888/fig-1)

between every reference gene and judges reference gene stability based on the standard deviation (SD) from the index, genes that have a smaller SD will be ranked more highly by BestKeeper. Finally, RefFinder was used to combine the ranked results from each algorithm and assign each gene an overall rank.

Data analyses were performed to compare the stability of among candidate reference genes in each tissue of interest. Following the tissue-specific analyses, we also combined all data for analysis together to simulate the conditions of a whole-body analysis. This approach allowed for the identification of which genes are most stable in each tissue of interest and provided evidence for which genes would be the strongest reference genes in an RT-qPCR experiment that used RNA from a whole *L. stagnalis* body. Given that a minimum of two reference genes are typically recommended for proper RT-qPCR data normalization, the recommended reference genes are generally expressed in terms of pairs.

RESULTS

Primer specificity and efficiency

Primers for all candidate reference genes were evaluated to ensure that they could produce consistent results and not amplify off-target products or generate primer dimers. Following amplification, each primer pair produced amplicons that yielded single bands at the correct size after electrophoresis in 2% agarose gels (Fig. 1). Additionally, no amplification was observed in controls that lacked reverse transcriptase in the RT-PCR or lacked cDNA template or primers in qPCR. Thus, primer pairs specifically amplified a single cDNA target. Based on the standard curves, primer set efficiencies ranged from 90% (*GAPDH* and *Rapac1*) to 106% (*ACTB*) with correlation coefficients (R^2) of >0.980 (Table 1).

Tissue-specific expression profiles and stability of reference genes

The candidate reference genes showed a variety of expression levels across individual tissues (Fig. 2). *Lst-UBI* produced the smallest mean Cq values and displayed the smallest Cq ranges for most tissues (Table 2). Conversely, *Lst-VGKC* mRNA was the least abundant in every tissue and displayed large Cq ranges.

The algorithms used to measure the stability of each candidate reference gene were highly congruent in their tissue-specific rankings. Therefore, only the results from the RefFinder analysis (which combines the rankings from the other three algorithms) are presented in Fig. 3 (results for other algorithms reported in Figs. S1–S3). *Lst-GAPDH* and

Figure 2 Expression levels for each gene in each tissue based on Cqs. Data was collected from three replicates of *L. stagnalis* tissue and each reaction was performed in triplicate. The tissues examined were: (A) CNS, (B) tentacles, (C) lips, (D) penis, (E) foot, and (F) mantle. Box plot: upper and lower box limits indicate 25th and 75th percentiles, line of division in box indicates the median, and whiskers indicate the minimum/maximum values.

[Full-size !\[\]\(666e09182d4cd268646ea700ea60dcdf_img.jpg\) DOI: 10.7717/peerj.7888/fig-2](https://doi.org/10.7717/peerj.7888/fig-2)

Table 2 Mean Cq values and standard errors of the mean (SEM) of reference genes for tissues in *Lymnaea stagnalis*. Means were calculated from three replicates of *L. stagnalis* tissue, each reaction was performed in triplicate.

Tissue	<i>Lst</i> -ACTB		<i>Lst</i> -EF1α		<i>Lst</i> -GAPDH		<i>Lst</i> -Rapac1		<i>Lst</i> -TUBB		<i>Lst</i> -UBI		<i>Lst</i> -VGKC	
	Cq mean	SEM	Cq mean	SEM	Cq mean	SEM	Cq mean	SEM	Cq mean	SEM	Cq mean	SEM	Cq mean	SEM
CNS	19.27	0.67	19.49	0.53	21.10	0.56	26.81	0.81	18.30	0.76	16.72	0.20	29.27	0.82
Tentacles	17.42	0.20	18.50	0.22	20.14	0.21	27.36	0.43	17.99	0.36	16.41	0.19	28.86	0.23
Lips	18.19	0.27	19.14	0.23	21.01	0.21	26.97	0.19	18.39	0.29	17.11	0.27	29.26	0.28
Penis	18.30	0.38	17.87	0.25	19.71	0.25	26.44	0.35	22.05	1.98	16.55	0.16	30.60	0.23
Foot	18.08	0.41	18.91	0.48	20.22	0.46	25.72	0.46	18.53	0.25	16.69	0.24	30.48	0.17
Mantle	17.67	0.26	18.91	0.24	20.41	0.34	27.57	0.39	22.16	0.42	16.24	0.16	30.78	0.31
Overall	18.15	0.17	18.80	0.15	20.43	0.16	26.81	0.20	19.57	0.43	16.62	0.09	29.87	0.19

Figure 3 RefFinder rankings for all candidate reference genes by tissue. RefFinder rankings for the (A) CNS, (B) tentacles, (C) lips, (D) penis, (E) foot, and (F) mantle. RefFinder calculates rankings as the geometric mean of the rankings assigned by geNorm, NormFinder, and BestKeeper. Genes are ranked in order from the least stable to the most stable in each panel. [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.7888/fig-3](https://doi.org/10.7717/peerj.7888/fig-3)

Lst-EF1α were the most stable candidate reference genes in the CNS and penis. *Lst-UBI* and *Lst-GAPDH* were the most stable in the tentacles, *Lst-GAPDH* and *Lst-Rapac1* were the most stable in the lips, *Lst-UBI* and *Lst-ACTB* were the most stable in the foot, and *Lst-UBI* and *Lst-VGKC* were the most stable in the mantle. Overall, identification of the most reliable candidate reference gene was tissue dependent. However, according to geNorm, all genes except for *Lst-ACTB* and *Lst-TUBB* were acceptable as a reference genes for all tissues as they met the minimum stability threshold of $M < 1.50$ for heterogeneous tissue samples.

Expression levels and stability of reference genes in a combined tissue analysis

To assess which candidate reference genes could be the most stable in whole-body preparations of *L. stagnalis*, the data from all tissues was combined and analyzed together with geNorm, NormFinder, BestKeeper, and RefFinder. The candidate reference genes spanned a wide range of expression levels when data from all tissue types was combined. Based on the Cq values, *Lst-UBI* was the most highly expressed gene (mean Cq 16.62 ± 0.09) and had the smallest overall Cq range whereas *Lst-VGKC* was the least expressed gene and had the highest biological variability as measured by the standard error of the mean (mean Cq 29.87 ± 0.19 ; Table 2). All four algorithms used to assess expression stability were highly congruent in their ranking of candidate reference genes in the combined analysis (Fig. 4). GeNorm, NormFinder, and RefFinder rated *Lst-GAPDH* and *Lst-EF1α* as the top two most stable reference genes. BestKeeper ranked *Lst-UBI* as the most stable with *Lst-EF1α* and *Lst-GAPDH* as the second and third most stable, respectively.

DISCUSSION

We employed several established algorithms to analyze mRNA abundance data and identify stably expressed genes that are suitable as reference genes for RT-qPCR in *L. stagnalis*. We assessed seven candidate reference genes (*Lst-ACTB*, *Lst-EF1α*, *Lst-GAPDH*, *Lst-Rapac1*, *Lst-TUBB*, *Lst-UBI*, and *Lst-VGKC*) in six tissues of interest (CNS, tentacles, lips, penis, foot, and mantle) and provided a separate analysis of all tissues combined. There were variable patterns of expression stability among the genes in the different tissues, but the tissue-specific rankings produced by the different algorithms were highly congruent. *Lst-GAPDH* and *Lst-EF1α* were the most suitable pair of reference genes in the CNS and penis. However, *Lst-UBI* also was one of the two most stable genes the tentacles, foot, and mantle when paired with *Lst-GAPDH*, *Lst-ACTB*, and *Lst-VGKC*, respectively. Finally, *Lst-GAPDH* and *Lst-Rapac1* were the most stable in the lips. In the combined analysis, *Lst-GAPDH*, *Lst-EF1α*, and *Lst-UBI* were the most stable genes according to all algorithms. Thus, for future experiments using either whole animals or tissues or tissue combinations not included in our analysis, we recommend these three genes as the first candidates for validation as reference genes.

Previous measurements of RT-qPCR reference gene stability has revealed some consistent patterns across the molluscs. *EF1α* appears to be the most effective reference

Figure 4 Combined analysis rankings assigned to each candidate reference gene. Data from all tissues was combined and assigned ranks by (A) geNorm, (B) NormFinder, (C) BestKeeper, and (D) RefFinder. *Indicates that both genes were equally recommended. [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.7888/fig-4](https://doi.org/10.7717/peerj.7888/fig-4)

gene in several mollusc species in addition to the CNS and penis of *L. stagnalis* (Table 3). In one study of *L. stagnalis*, *EF1α* was identified to remain stable in the CNS under heat stress (Foster, Lukowiak & Henry, 2015). Given that *EF1α* has also been identified to be highly stable in several other molluscs, it appears to be a strong candidate reference gene across the phylum (Morga et al., 2010; Wan et al., 2011; Cubero-Leon et al., 2012; Mauriz et al., 2012; Moreira et al., 2014; García-Fernández et al., 2016; Huan, Wang & Liu, 2016). An analysis of reference genes in the freshwater snail *Bellamya aeruginosa* showed

Table 3 Summary of the stability rankings of reference genes from studies conducted in molluscs.

Organism	Most stable gene(s)	Least stable gene(s)	Other genes tested	Reference
Gastropods				
<i>Haliotis discus</i>	<i>EF1α/RPL5</i>	<i>18S rRNA</i>	<i>ACTB, BGLU, CY, GAPDH, H2A, HPRT, SDHA, TUBB, UBC, CYP4</i>	Wan et al. (2011)
<i>Bellamyia aeruginosa</i>	<i>RPL7</i>	<i>ACTB</i>	<i>18S rRNA, EF1α, GAPDH, TUBB, H2A, DRP2</i>	Liu et al. (2016)
<i>Cepaea nemoralis</i>	<i>EF1α/ACTB</i>	<i>GAPDH</i>	<i>DNARP, FIB3, GTP8, Rapac1, RNAP, TUBA, UBI, VGKC</i>	Affenzeller, Cerveau & Jackson (2018)
<i>Lymnaea stagnalis</i>	<i>EF1α/GAPDH</i>	<i>ACTB/TUBB</i>	<i>VGKC, UBI, Rapac1</i>	Present study
Cephalopods				
<i>Octopus vulgaris</i>	<i>TUBA/UBI</i>	<i>18S rRNA</i>	<i>16S rRNA, ACTB, EF1α, TUBA</i>	Sirakov et al. (2009)
<i>Octopus vulgaris</i>	<i>UBI</i>	<i>ACTB</i>	<i>18S rRNA, EF1α, GAPDH, TUBA</i>	García-Fernández et al. (2016)
Bivalves				
<i>Ostrea edulis</i>	<i>EF1α/GAPDH</i>	<i>ACTB</i>	<i>UBI, RPL5</i>	Morga et al. (2010)
<i>Mytilus edulis</i>	<i>EF1α/18S rRNA</i>	<i>ACTB</i>	<i>28S rRNA, TUBA, HEL</i>	Cubero-Leon et al. (2012)
<i>Mytilus galloprovincialis</i>	<i>EF1α</i>	<i>ACTB</i>	<i>18S rRNA, TUBA</i>	Moreira et al. (2014)
<i>Ruditapes philippinarum</i>	<i>TUBA</i>	<i>ACTB</i>	<i>18S rRNA, EF1α</i>	Moreira et al. (2014)
<i>Crassostrea gigas</i>	<i>EF1α</i>	<i>RPS18</i>	<i>ACTB, ARF1, GAPDH, HNRPQ, UBC</i>	Huan, Wang & Liu (2016)
<i>Mytilus galloprovincialis</i>	<i>GAPDH/RPS4</i>	<i>NAD4/18S rRNA</i>	<i>ACTB, COX1, GAPDH, RPS27, TIF5A</i>	Martínez-Escauriaza et al. (2018)

Note:

ARF1, adp-ribosylation factor 1; BGLU, beta-glucuronidase; COX1, cytochrome c oxidase subunit 1; CY, cyclophilin; CYP4, cytochrome p450 family 4; DNARP, DNA repair protein; DRP2, DNA-directed RNA polymerase II; FIB3, fibronectin type III domain containing protein; GTP8, GTP-binding protein; H2A, histone H2A; HEL, RNA helicase; HNRPQ, heterogeneous nuclear ribonucleoprotein q; HPRT, hypoxanthine phosphoribosyltransferase 1; NAD4, NADH dehydrogenase subunit 4; RNAP, RNA-directed DNA polymerase; RPL5, ribosomal protein L5; RPL7, ribosomal protein L7; RPS4, 40S ribosomal protein S4; RPS27, 40S ribosomal protein S27; SDHA, succinate dehydrogenase; UBC, ubiquitin-conjugating enzyme.

that *EF1α* and *GAPDH* were stable in the tentacles and penis but more variable in the foot ([Liu et al., 2015](#)), similar to our findings (albeit with some discrepancies of rankings in the tentacles and foot). *GAPDH* has also been shown to be a stable reference gene in bivalves ([Morga et al., 2010](#); [Martínez-Escauriaza et al., 2018](#)) but was reported as unsuitable in abalone and octopus ([Wan et al., 2011](#); [García-Fernández et al., 2016](#)). Thus, *GAPDH* is not consistent across the molluscs. Additionally, *GAPDH* is highly stable in the terrestrial gastropod *C. nemoralis* within a given season, but expression levels in this species are subject to great variation between seasons ([Affenzeller, Cerveau & Jackson, 2018](#)). In our data, *Lst-UBI* had a very consistent expression profile between tissues compared to other reference genes and was found to be highly stable in the foot, mantle, and tentacles as well as in the combined analysis. *UBI* has also shown promise in other molluscs ([Sirakov et al., 2009](#); [García-Fernández et al., 2016](#); [Affenzeller, Cerveau & Jackson, 2018](#)). *Alpha-tubulin* (*TUBA*) has generally produced positive results as a reference gene in several molluscan species ([Sirakov et al., 2009](#); [Cubero-Leon et al., 2012](#); [Moreira et al., 2014](#)). However, *TUBA* was unstable in *C. nemoralis* ([Affenzeller, Cerveau & Jackson, 2018](#)) and we also found *Lst-TUBB* to be highly unstable. *ACTB* has generally produced negative results as a candidate reference gene in molluscs ([Cubero-Leon et al., 2012](#);

Moreira et al., 2014; Liu et al., 2015; García-Fernández et al., 2016; Huan, Wang & Liu, 2016) and the results here from *L. stagnalis* are largely congruent with these previous findings, although *Lst-ACTB* was stable specifically in the foot. Finally, *Rapac1* and *VGKC* were identified as suitable reference genes in *C. nemoralis* (Affenzeller, Cerveau & Jackson, 2018). These genes did show relative stability in our analyses of the mantle and lips, but were relatively unstable in the other tissues examined.

Despite the heterogeneity in top-ranked reference genes among the tissues, there was some consistency in expression stability across all tissues. Though *Lst-EF1 α* , *Lst-GAPDH*, and *Lst-UBI* were not the top ranked in every tissue as judged by RefFinder, they were relatively stable in all tissues tested which is reflected in the stability values provided by geNorm. Thus, we recommend these genes for use in multi-tissue comparisons in *L. stagnalis*. In particular, *Lst-EF1 α* and *Lst-GAPDH* are a favorable pair as they have the most similar Cq ranges. Importantly, despite the demonstrated stability of these genes under baseline conditions reported here, it would be crucial to directly verify their stability under any experimental conditions (Kozera & Rapacz, 2013).

CONCLUSION

The validation of stable reference genes is necessary for the acquisition of reliable gene expression data. Therefore, it is important to perform a species-specific verification of reference gene stability before undertaking RT-qPCR experiments. The variable expression of the seven genes among the six tissue types we investigated in *L. stagnalis* demonstrates that it is critical to select reference genes based on the tissues of interest. The results presented here should guide the selection of reference genes for tissue-specific RT-qPCR and thereby assist with future studies of gene expression in the snails.

ACKNOWLEDGEMENTS

We thank Dr. Scott Cummins for advice, the animal care staff at St. Francis Xavier University for maintaining the *Lymnaea stagnalis* colony, and both Hisayo Sadamoto and anonymous reviewer for constructive comments.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Support was received from the Canadian Foundation for Innovation (Grant 25196 to Dr. Cory Bishop, Grant to 17075 Dr. Angela Beye, and Grant 19286 to Dr. Russell Wyeth), the Natural Sciences and Engineering Research Council of Canada (Discovery grant RGPIN-2015-04957 to Dr. Russell Wyeth, CGS-M and a Michael Smith Foreign Study Supplement to Alexander Young), the Deutsche Forschungsgemeinschaft (Grant JA 2108/6-1 to Dr. Daniel Jackson), and St. Francis Xavier University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Canadian Foundation for Innovation: 25196, 17075, 19286.

Natural Sciences and Engineering Research Council of Canada: RGPIN-2015-04957.

Deutsche Forschungsgemeinschaft: JA 2108/6-1.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Alexander P. Young conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Carmen F. Landry performed the experiments, analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Daniel J. Jackson contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft, supplied transcriptomes used to identify *L. stagnalis* gene sequences.
- Russell C. Wyeth conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The novel reference gene sequences (GAPDH, EF1a, Rapac1, UBI, VGKC) are available at GenBank: [MH687363–MH687367](#).

Data Availability

The following information was supplied regarding data availability:

The raw data set is available at Figshare: Young, Alex (2019): Primary dataset—RefGene ALL Raw Cqs (500 ng).xlsx. figshare. Dataset. [DOI 10.6084/m9.figshare.8329937.v1](https://doi.org/10.6084/m9.figshare.8329937.v1)

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.7888#supplemental-information>.

REFERENCES

- Abe M, Kuroda R. 2019.** The development of CRISPR for a mollusc establishes the formin *Lsdia1* as the long-sought gene for snail dextral/sinistral coiling. *Development* **146**(9):dev175976 [DOI 10.1242/dev.175976](https://doi.org/10.1242/dev.175976).
- Affenzeller S, Cerveau N, Jackson DJ. 2018.** Identification and validation of reference genes for qPCR in the terrestrial gastropod *Cepaea nemoralis*. *PLOS ONE* **13**(8):e0201396 [DOI 10.1371/journal.pone.0201396](https://doi.org/10.1371/journal.pone.0201396).
- Aleksic M, Feng Z-P. 2012.** Identification of the role of C/EBP in neurite regeneration following microarray analysis of a *L. stagnalis* CNS injury model. *BMC Neuroscience* **13**(1):2 [DOI 10.1186/1471-2202-13-2](https://doi.org/10.1186/1471-2202-13-2).

- Andersen CL, Jensen JL, Ørntoft TF. 2004. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Research* 64(15):5245–5250 DOI 10.1158/0008-5472.CAN-04-0496.
- Barber RD, Harmer DW, Coleman RA, Clark BJ. 2005. GAPDH as a housekeeping gene: analysis of GAPDH mRNA expression in a panel of 72 human tissues. *Physiological Genomics* 21(3):389–395 DOI 10.1152/physiolgenomics.00025.2005.
- Bavan S, Straub VA, Webb TE, Ennion SJ. 2012. Cloning and characterization of a P2X receptor expressed in the central nervous system of *Lymnaea stagnalis*. *PLOS ONE* 7(11):e50487 DOI 10.1371/journal.pone.0050487.
- Benatti C, Colliva C, Blom JY, Ottaviani E, Tascadda F. 2017. Transcriptional effect of serotonin in the ganglia of *Lymnaea stagnalis*. *Invertebrate Survival Journal* 14:251–258.
- Benjamin P. 2008. *Lymnaea*. *Scholarpedia* 3(1):4124 DOI 10.4249/scholarpedia.4124.
- Boer HH, Montagne-Wajer C, Van Minnen J, Ramkema M, De Boer P. 1992. Functional morphology of the neuroendocrine sodium influx-stimulating peptide system of the pond snail, *Lymnaea stagnalis*, studied by in situ hybridization and immunocytochemistry. *Cell and Tissue Research* 268(3):559–566 DOI 10.1007/BF00319163.
- Boer HH, Witteveen J. 1980. Ultrastructural localization of carbonic anhydrase in tissues involved in shell formation and ionic regulation in the pond snail *Lymnaea stagnalis*. *Cell and Tissue Research* 209(3):383–390 DOI 10.1007/BF00234753.
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clinical Chemistry* 55(4):611–622 DOI 10.1373/clinchem.2008.112797.
- Carter CJ, Rand C, Mohammad I, Lepp A, Vesprini N, Wiebe O, Carlone R, Spencer GE. 2015. Expression of a retinoic acid receptor (RAR)-like protein in the embryonic and adult nervous system of a protostome species. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 324(1):51–67 DOI 10.1002/jez.b.22604.
- Chase R. 2002. *Behavior and its neural control in gastropod molluscs*. Oxford: Oxford University Press.
- Croll RP, Van Minnen J. 1992. Distribution of the peptide Ala-Pro-Gly-Trp-NH₂ (APGWamide) in the nervous system and periphery of the snail *Lymnaea stagnalis* as revealed by immunocytochemistry and in situ hybridization. *Journal of Comparative Neurology* 324(4):567–574 DOI 10.1002/cne.903240409.
- Cubero-Leon E, Ciocan CM, Minier C, Rotchell JM. 2012. Reference gene selection for qPCR in mussel, *Mytilus edulis*, during gametogenesis and exogenous estrogen exposure. *Environmental Science and Pollution Research* 19(7):2728–2733 DOI 10.1007/s11356-012-0772-9.
- Davison A, McDowell GS, Holden JM, Johnson HF, Koutsovoulos GD, Liu MM, Hulpiau P, Van Roy F, Wade CM, Banerjee R, Yang F, Chiba S, Davey JW, Jackson DJ, Levin M, Blaxter ML. 2016. Formin is associated with left-right asymmetry in the pond snail and the frog. *Current Biology* 26(5):654–660 DOI 10.1016/j.cub.2015.12.071.
- De Jonge HJM, Fehrmann RSN, De Bont ESJM, Hofstra RMW, Gerbens F, Kamps WA, De Vries EGE, Van Der Zee AGJ, Te Meerman GJ, Ter Elst A. 2007. Evidence based selection of housekeeping genes. *PLOS ONE* 2(9):e898 DOI 10.1371/journal.pone.0000898.
- Dheda K, Huggett JF, Bustin SA, Johnson MA, Rook G, Zumla A. 2004. Validation of housekeeping genes for normalizing RNA expression in real-time PCR. *BioTechniques* 37(1):112–119 DOI 10.2144/04371RR03.

- Dirks RW, Raap AK, Van Minnen J, Vreugdenhil E, Smit AB, Van Der Ploeg M. 1989. Detection of mRNA molecules coding for neuropeptide hormones of the pond snail *Lymnaea stagnalis* by radioactive and non-radioactive in situ hybridization: a model study for mRNA detection. *Journal of Histochemistry & Cytochemistry* 37(1):7–14 DOI 10.1177/37.1.2642295.
- Dong N, Senzel A, Li K, Lu TZ, Guo C-H, Aleksic M, Feng Z-P. 2018. MEN1 tumor suppressor gene is required for long-term memory formation in an aversive operant conditioning model of *Lymnaea stagnalis*. *Neuroscience* 379:22–31 DOI 10.1016/j.neuroscience.2018.02.018.
- Ebanks SC, O'Donnell MJ, Grosell M. 2010. Characterization of mechanisms for Ca^{2+} and $\text{HCO}_3^-/\text{CO}_3^{2-}$ acquisition for shell formation in embryos of the freshwater common pond snail *Lymnaea stagnalis*. *Journal of Experimental Biology* 213(23):4092–4098 DOI 10.1242/jeb.045088.
- Eissa N, Hussein H, Wang H, Rabbi MF, Bernstein CN, Ghia J-E. 2016. Stability of reference genes for messenger RNA quantification by real-time PCR in mouse dextran sodium sulfate experimental colitis. *PLOS ONE* 11(5):e0156289 DOI 10.1371/journal.pone.0156289.
- Elliott CJH, Susswein AJ. 2002. Comparative neuroethology of feeding control in molluscs. *Journal of Experimental Biology* 205:877–896.
- Feng Z-P, Zhang Z, Van Kesteren RE, Straub VA, Van Nierop P, Jin K, Nejatbakhsh N, Goldberg JI, Spencer GE, Yeoman MS, Wildering W, Coorsen JR, Croll RP, Buck LT, Syed NI, Smit AB. 2009. Transcriptome analysis of the central nervous system of the mollusc *Lymnaea stagnalis*. *BMC Genomics* 10(1):451 DOI 10.1186/1471-2164-10-451.
- Flynn N, Getz A, Visser F, Janes TA, Syed NI. 2014. Menin: a tumor suppressor that mediates postsynaptic receptor expression and synaptogenesis between central neurons of *Lymnaea stagnalis*. *PLOS ONE* 9(10):e111103 DOI 10.1371/journal.pone.0111103.
- Foster NL, Lukowiak K, Henry TB. 2015. Time-related expression profiles for heat shock protein gene transcripts (HSP40, HSP70) in the central nervous system of *Lymnaea stagnalis* exposed to thermal stress. *Communicative & Integrative Biology* 8(3):e1040954 DOI 10.1080/19420889.2015.1040954.
- García-Fernández P, Castellanos-Martínez S, Iglesias J, Otero JJ, Gestal C. 2016. Selection of reliable reference genes for RT-qPCR studies in *Octopus vulgaris* paralarvae during development and immune-stimulation. *Journal of Invertebrate Pathology* 138:57–62 DOI 10.1016/j.jip.2016.06.003.
- Getz AM, Wijdenes P, Riaz S, Syed NI. 2018. Uncovering the cellular and molecular mechanisms of synapse formation and functional specificity using central neurons of *Lymnaea stagnalis*. *ACS Chemical Neuroscience* 9(8):1928–1938 DOI 10.1021/acschemneuro.7b00448.
- Gust M, Fortier M, Garric J, Fournier M, Gagné F. 2013a. Immunotoxicity of surface waters contaminated by municipal effluents to the snail *Lymnaea stagnalis*. *Aquatic Toxicology* 126:393–403 DOI 10.1016/j.aquatox.2012.09.001.
- Gust M, Fortier M, Garric J, Fournier M, Gagné F. 2013b. Effects of short-term exposure to environmentally relevant concentrations of different pharmaceutical mixtures on the immune response of the pond snail *Lymnaea stagnalis*. *Science of the Total Environment* 445–446:210–218 DOI 10.1016/j.scitotenv.2012.12.057.
- Hatakeyama D, Okuta A, Otsuka E, Lukowiak K, Ito E. 2013. Consolidation of long-term memory by insulin in *Lymnaea* is not brought about by changing the number of insulin receptors. *Communicative & Integrative Biology* 6(3):e23955 DOI 10.4161/cib.23955.
- Herlitze I, Marie B, Marin F, Jackson DJ. 2018. Molecular modularity and asymmetry of the molluscan mantle revealed by a gene expression atlas. *GigaScience* 7(6):giy056 DOI 10.1093/gigascience/giy056.

- Hohagen J, Jackson DJ. 2013. An ancient process in a modern mollusc: early development of the shell in *Lymnaea stagnalis*. *BMC Developmental Biology* 13(1):27 DOI 10.1186/1471-213X-13-27.
- Huan P, Wang H, Liu B. 2016. Assessment of housekeeping genes as internal references in quantitative expression analysis during early development of oyster. *Genes & Genetic Systems* 91(5):257–265 DOI 10.1266/ggs.16-00007.
- Jackson DJ, Herlitze I, Hohagen J. 2016. A whole mount in situ hybridization method for the gastropod mollusc *Lymnaea stagnalis*. *Journal of Visualized Experiments* 15:109 DOI 10.3791/53968.
- Koene JM. 2010. Neuro-endocrine control of reproduction in hermaphroditic freshwater snails: Mechanisms and evolution. *Frontiers in Behavioral Neuroscience* 4:167 DOI 10.3389/fnbeh.2010.00167.
- Korneev SA, Kemenes I, Bettini NL, Kemenes G, Staras K, Benjamin PR, O'Shea M. 2013. Axonal trafficking of an antisense RNA transcribed from a pseudogene is regulated by classical conditioning. *Scientific Reports* 3(1):1027 DOI 10.1038/srep01027.
- Korneev SA, Vavoulis DV, Naskar S, Dyakonova VE, Kemenes I, Kemenes G. 2018. A CREB2-targeting microRNA is required for long-term memory after single-trial learning. *Scientific Reports* 8(1):3950 DOI 10.1038/s41598-018-22278-w.
- Kozera B, Rapacz M. 2013. Reference genes in real-time PCR. *Journal of Applied Genetics* 54(4):391–406 DOI 10.1007/s13353-013-0173-x.
- Liu Q, Lei K, Ma Q, Qiao F, Li Z, An L. 2016. Ribosomal protein L7 as a suitable reference gene for quantifying gene expression in gastropod *Bellamya aeruginosa*. *Environmental Toxicology and Pharmacology* 43:120–127 DOI 10.1016/j.etap.2016.03.007.
- Liu L-L, Zhao H, Ma T-F, Ge F, Chen C-S, Zhang Y-P. 2015. Identification of valid reference genes for the normalization of RT-qPCR expression studies in human breast cancer cell lines treated with and without transient transfection. *PLOS ONE* 10(1):e0117058 DOI 10.1371/journal.pone.0117058.
- Lukowiak K. 2016. Stress, memory, forgetting and what, *Lymnaea* can tell us about a stressful world. In: Saleuddin S, Mukai S, eds. *Physiology of Molluscs Volume 2: A Collection of Selected Reviews*. Palm Bay: Apple Academic Press and CRC Press, 68–97.
- Lukowiak K, Haque Z, Spencer G, Varshay N, Sangha S, Syed N. 2003. Long-term memory survives nerve injury and the subsequent regeneration process. *Learning & Memory* 10(1):44–54 DOI 10.1101/lm.48703.
- Martínez-Escauriaza R, Lozano V, Pérez-Parallé ML, Pazos AJ, Sánchez JL. 2018. Validation of reference genes in mussel *Mytilus galloprovincialis* tissues under the presence of okadaic acid. *Journal of Shellfish Research* 37(1):93–102 DOI 10.2983/035.037.0108.
- Mauriz O, Maneiro V, Pérez-Parallé ML, Sánchez JL, Pazos AJ. 2012. Selection of reference genes for quantitative RT-PCR studies on the gonad of the bivalve mollusc *Pecten maximus* L. *Aquaculture* 370-371:158–165 DOI 10.1016/j.aquaculture.2012.10.020.
- Moreira R, Pereiro P, Costa MM, Figueras A, Novoa B. 2014. Evaluation of reference genes of *Mytilus galloprovincialis* and *Ruditapes philippinarum* infected with three bacteria strains for gene expression analysis. *Aquatic Living Resources* 27(3–4):147–152 DOI 10.1051/alr/2014015.
- Morga B, Arzul I, Faury N, Renault T. 2010. Identification of genes from flat oyster *Ostrea edulis* as suitable housekeeping genes for quantitative real time PCR. *Fish & Shellfish Immunology* 29(6):937–945 DOI 10.1016/j.fsi.2010.07.028.
- Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper – Excel-based

- tool using pair-wise correlations. *Biotechnology Letters* **26**(6):509–515
DOI 10.1023/B:BILE.0000019559.84305.47.
- Pirger Z, Laszlo Z, Hiripi L, Hernadi L, Toth G, Lubics A, Reglodi D, Kemenes G, Mark L. 2010. Pituitary adenylate cyclase activating polypeptide (PACAP) and its receptors are present and biochemically active in the central nervous system of the pond snail *Lymnaea stagnalis*. *Journal of Molecular Neuroscience* **42**(3):464–471 DOI 10.1007/s12031-010-9361-x.
- Ruijter JM, Ramakers C, Hoogaars WMH, Karlen Y, Bakker O, Van Den Hoff MJB, Moorman AFM. 2009. Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Research* **37**(6):e45 DOI 10.1093/nar/gkp045.
- Sadamoto H, Kitahashi T, Fujito Y, Ito E. 2010. Learning-dependent gene expression of CREB1 isoforms in the molluscan brain. *Frontiers in Behavioral Neuroscience* **4**:25 DOI 10.3389/fnbeh.2010.00025.
- Sadamoto H, Sato H, Kobayashi S, Murakami J, Aonuma H, Ando H, Fujito Y, Hamano K, Awaji M, Lukowiak K, Urano A, Ito E. 2004. CREB in the pond snail *Lymnaea stagnalis*: cloning, gene expression, and function in identifiable neurons of the central nervous system. *Journal of Neurobiology* **58**(4):455–466 DOI 10.1002/neu.10296.
- Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative C_T method. *Nature Protocols* **3**(6):1101–1108 DOI 10.1038/nprot.2008.73.
- Senatore A, Spafford JD. 2012. Gene transcription and splicing of T-Type channels are evolutionarily-conserved strategies for regulating channel expression and gating. *PLOS ONE* **7**(6):e37409 DOI 10.1371/journal.pone.0037409.
- Shimizu K, Iijima M, Setiamarga DHE, Sarashina I, Kudoh T, Asami T, Gittenberger E, Endo K. 2013. Left-right asymmetric expression of dpp in the mantle of gastropods correlates with asymmetric shell coiling. *EvoDevo* **4**(1):15 DOI 10.1186/2041-9139-4-15.
- Sirakov M, Zarrella I, Borra M, Rizzo F, Biffali E, Arnone MI, Fiorito G. 2009. Selection and validation of a set of reliable reference genes for quantitative RT-PCR studies in the brain of the cephalopod mollusc *Octopus vulgaris*. *BMC Molecular Biology* **10**(1):70 DOI 10.1186/1471-2199-10-70.
- Tong Z, Gao Z, Wang F, Zhou J, Zhang Z. 2009. Selection of reliable reference genes for gene expression studies in peach using real-time PCR. *BMC Molecular Biology* **10**(1):71 DOI 10.1186/1471-2199-10-71.
- Van Der Knaap WPW, Adema CM, Sminia T. 1993. Invertebrate blood cells: morphological and functional aspects of the haemocytes in the pond snail *Lymnaea stagnalis*. *Comparative Haematology International* **3**(1):20–26 DOI 10.1007/BF00394923.
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* **3**(7):research0034.1 DOI 10.1186/gb-2002-3-7-research0034.
- Wan Q, Whang I, Choi CY, Lee J-S, Lee J. 2011. Validation of housekeeping genes as internal controls for studying biomarkers of endocrine-disrupting chemicals in disk abalone by real-time PCR. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **153**(3):259–268 DOI 10.1016/j.cbpc.2010.11.009.
- Wyeth RC, Croll RP, Willows AOD, Spencer AN. 2009. 1-Phenoxy-2-propanol is a useful anaesthetic for gastropods used in neurophysiology. *Journal of Neuroscience Methods* **176**(2):121–128 DOI 10.1016/j.jneumeth.2008.08.028.
- Xie F, Xiao P, Chen D, Xu L, Zhang B. 2012. miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Molecular Biology* **80**(1):75–84 DOI 10.1007/s11103-012-9885-2.