

REVIEW

Dysfunction of RAB39B-Mediated Vesicular Trafficking in Lewy Body Diseases

David J. Koss, PhD,^{1*}  Susanna Campesan, PhD,² Flaviano Giorgini, PhD,² and Tiago F. Outeiro, PhD^{1,3,4,5*}

¹Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

²Department of Genetics and Genome Biology, University of Leicester, University Road, Leicester, UK

³Department of Experimental Neurodegeneration, Center for Biostructural Imaging of Neurodegeneration, University Medical Center Goettingen, Goettingen, Germany

⁴Max Planck Institute for Experimental Medicine, Goettingen, Germany

⁵Scientific employee with a honorary contract at Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), Göttingen, Germany

ABSTRACT: Intracellular vesicular trafficking is essential for neuronal development, function, and homeostasis and serves to process, direct, and sort proteins, lipids, and other cargo throughout the cell. This intricate system of membrane trafficking between different compartments is tightly orchestrated by Ras analog in brain (RAB) GTPases and their effectors. Of the 66 members of the RAB family in humans, many have been implicated in neurodegenerative diseases and impairment of their functions contributes to cellular stress, protein aggregation, and death. Critically, RAB39B loss-of-function mutations are known to be associated with X-linked intellectual disability and with rare early-onset Parkinson's disease. Moreover, recent studies have highlighted altered RAB39B expression in idiopathic cases of

several Lewy body diseases (LBDs). This review contextualizes the role of RAB proteins in LBDs and highlights the consequences of RAB39B impairment in terms of endosomal trafficking, neurite outgrowth, synaptic maturation, autophagy, as well as alpha-synuclein homeostasis. Additionally, the potential for therapeutic intervention is examined via a discussion of the recent progress towards the development of specific RAB modulators. © 2021 The Authors. *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society

Key Words: RAB39B; Lewy body diseases; alpha-synuclein; endocytosis; neurodegeneration

RAB Proteins and Their Regulation

The intracellular system of membrane trafficking is an essential aspect of cell physiology, being at the core of mechanisms for exocytosis, endocytosis, movement, and degradation of cargo within the cell. Ras analog in brain (RAB) GTPases, the largest branch of the Ras-like small GTPase superfamily, are the master regulators of cellular vesicle traffic, with 66 members having been

described in humans, 31 in *Drosophila*, and 11 in yeast.^{1,2} RAB proteins spatio-temporally regulate membrane docking, tethering, and movements along the cytoskeleton during the various steps of trafficking processes.³ These processes are driven by their ability to act as molecular switches oscillating from cytosolic inactive-GDP-bound, to membrane-associated active-GTP-bound states.⁴ Crucial to the interaction of RAB proteins with membrane vesicles are several key

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

***Correspondence to:** Dr. D.J. Koss and Dr. T. Outeiro, Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK, NE4 5PL; E-mail: david.koss@newcastle.ac.uk and E-mail: tiago.outeiro@newcastle.ac.uk

Relevant conflicts of interest/financial disclosures: The authors declare they have no conflict of interests financially or otherwise.

Funding agencies: T.F.O. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy (EXC 2067/1- 390729940) and by SFB1286 (Project B8). S.C. and F.G. are supported by funding from Parkinson's UK (G-1802). D.K., T.F.O., and F.G. are supported by Alzheimer's Research UK Newcastle Network Centre. T.F.O. and D.J.K. are supported by the Lewy Body Society (LBS-0007).

Received: 1 December 2020; **Revised:** 9 March 2021; **Accepted:** 12 March 2021

Published online 3 May 2021 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28605




FIG. 1. Ras analog in brain (RAB) regulation. **(A)** RAB protein activation cycle. The newly synthesized RAB, in the GDP-bound inactive form, is recognized by RAB escort protein (REP) (1). REP presents the RAB to a geranylgeranyl transferase (GGT), which geranylgeranylates the RAB on one or two carboxy-terminal Cys residues (prenylation) (2). Prenylation allows the RAB to associate with membranes (3). A guanine nucleotide exchange factor (GEF) catalyzes the exchange of GDP for GTP which activates the RAB. The GTP-bound active RAB associates with multiple effectors (4) and is then converted back to the GDP-bound inactive form by hydrolysis of GTP, which is catalyzed by a GTPase activating protein (GAP) (5). The RAB GDP dissociation inhibitor (GDI) recruits and maintains the GDP-bound RAB in the cytosol (6) until it is removed by a GDI displacement factor (GDF) which allows the RAB to associate with a membrane, restarting the cycle (3). **(B)** RAB protein cascade. During the maturing of a membrane a RAB cascade is achieved with the effectors of each RAB being a GEF for the next RAB and a GAP for the previous RAB. [Color figure can be viewed at wileyonlinelibrary.com]

regulators such as RAB prenylation escort protein (REP), which promotes geranylgeranyltransferase (GGTase)-mediated C-terminus prenylation of newly synthesized RABs (a prerequisite for their association with membranes),⁵ and specific membrane-associated guanine nucleotide exchange factors (GEFs), which recruit and activate RABs by promoting the exchange

of GDP to GTP. Activated RABs interact with a wide range of effector proteins whose functions include cargo sorting, vesicle formation, movement, tethering, and fusion.^{4,6} GTPase activating proteins (GAPs) accelerate the hydrolysis of GTP into GDP and inactivate RABs, which are then extracted from the membrane and chaperoned to the cytosol by a GDP dissociation

inhibitor (GDI), providing a pool of inactive RABs ready to be reutilized.⁷ GDI displacement factor (GDF) can subsequently promote GDI release and the RAB activation cycle can recommence (Fig. 1A).

As vesicles mature from one membrane compartment to the next, they associate with different RABs, and the specificity of each RAB can be orchestrated by cascades of RABs, GEFs, and GAPs (Fig. 1B). These RAB cycles are intimately connected with SNARE cycles, thereby regulating the fusion of vesicles with the target organelles/compartments.

Neuronal RABs

Given the highly specialized, dynamic, and polarized nature of neurons it is unsurprising that the maintenance and function of these long-lived cell types critically depend on vesicle transport, exocytosis, and endocytosis.⁸⁻¹⁰ Accordingly, several RABs, such as RAB3A and RAB6B, are specifically expressed in the brain, and many other RABs are enriched in the brain.¹¹ Together these neuron-specific RABs complement non-cell-type-specific RABs in order to orchestrate a variety of critical functions in neuronal homeostasis, such as neurite outgrowth and axon or dendrite formation,¹²⁻¹⁴ neurotransmitter release,¹⁵⁻¹⁸ the recycling or degradation of synaptic or endosomal vesicles,¹⁹⁻²⁹ and synaptic plasticity.³⁰⁻³⁵

Thus, due to the near ubiquitous involvement of RABs in neuronal homeostasis, it is unsurprising that their dysregulation as part of neurodegenerative processes has been widely reported.⁹ Nevertheless, the direct association of RAB gene mutations with neuropathy are rare, with the critical exceptions of RAB7A gene mutations associated with Charcot-Marie-Tooth disease type 2B (CMT2B),³⁶ RAB18 gene mutations with Warburg Micro syndrome,³⁷ and RAB39B with Parkinson's disease (PD).

Lewy Body Diseases

PD and the related dementia with Lewy bodies (DLB) are pathologically defined by the presence of α -synuclein (aSyn)-rich intraneuronal Lewy bodies (LBs) and are collectively referred to as Lewy body diseases (LBDs). PD is the most common movement disorder and is characterized by resting tremor, bradykinesia, rigidity, and postural instability.³⁸ The motor features are caused by the progressive loss of dopaminergic neurons in the substantia nigra pars compacta, concomitantly with the presentation nigral and brainstem LBs.³⁹ In contrast, DLB cases exhibit limbic/neocortical predominate LBs and present with visual hallucinations, cognitive fluctuations, rapid eye movement (REM) sleep behavior disorder, and one or more features of PD.⁴⁰ Despite the

identification of aSyn as the major component of LB pathology, the precise role of these protein inclusions in neurodegeneration remains unclear.

To date, numerous cellular stressors and impairments have associated with neurodegeneration, including oxidative stress, endoplasmic reticulum (ER) stress, DNA damage, mitochondrial dysfunction, and vesicular-mediated protein and lipid trafficking and degradation.^{41,42} Strikingly, although most LBD cases are sporadic, around 20 genes are associated with genetically inherited forms of PD including *SNCA* (aSyn), *LRRK2* (leucine-rich repeat kinase2), *VPS35* (vacuolar sortin protein 35), *PINK1* (PTEN-induced putative kinase1), *PARK2/Parkin*, and the subject of this review, *RAB39B*. All proteins encoded by these genes have been implicated in membrane trafficking and/or RAB function⁴³⁻⁴⁵ (see Table 1). Over the last decade, the identification of *RAB39B* mutations as causative in the occurrence of rare early-onset forms of PD has served to further highlight dysfunctional vesicular trafficking as a potential pathogenic source of disease.

RAB39B Mutations and Dysfunction

Originally identified as a mutation locus for X-linked intellectual disability (XLID), the initial study of XLID families reported a likely benign silent *RAB39B* mutation (c.543A > G p.T181T) within the cohort.⁹⁵ However, several *RAB39B* mutations were later identified as causative for XLID.⁹⁶ Affected families presenting with mild mental impairment and macrocephaly and individual cases demonstrating additional symptoms including autism spectrum disorder and/or seizure occurrence.⁹⁶ An association with early-onset PD in addition to XLID was reported in a follow-up study of Australian kindred and a genetically distinct Wisconsin family.⁶⁷ In addition to the symptomatic PD presentation, postmortem neuropathological examination of an individual Australian kindred confirmed the presence of cortical and subcortical LBs, neurofibrillary tangles (NFTs), and subcortical atrophy.⁶⁷ Similar observations were reported within a member of the Wisconsin family, in which LBs and NFTs were also apparent alongside subcortical atrophy and iron deposition.⁹⁷

To date, a number of *RAB39B* mutations have now been associated with XLID and early-onset PD, the majority of these mutations results in a total loss of *RAB39B* expression, although examples of reduced protein stability (C.503 > A p.T168K)⁶⁷ and altered function (c.574G > A p.G192R)⁹⁸ have also been reported (see Table 2 for full details). Interestingly, *RAB39B* duplication is also linked to XLID, suggesting that tight regulation of *RAB39B* activity is essential for physiological development. However, there are no reports of PD-like symptomology and neuropathological examination is currently unavailable.¹⁰⁴ Regardless,

TABLE 1 Familial Parkinson's disease genes and Ras analog in brain (RAB) protein associations

| Genes | Interacting RAB | Functional outcome |
|--------|--|--|
| aSyn | RAB1A | Overexpression rescues aSyn-induced ER-Golgi traffic defects ^{46,47} |
| | RAB3A | Stabilizes aSyn on synaptic membranes; overexpression rescues aSyn toxicity in animal models ^{46,48-52} |
| | RAB5A/B | Modulates aSyn clearance and spreading; interacts with mutant aSyn disrupting endocytosis ^{48,53-58} |
| | RAB7 | Modify aSyn clearance and spreading; overexpression rescues aSyn ^{48,49,55,57,59-66} |
| | RAB8 | |
| | RAB11A | |
| | RAB13 | |
| | RAB27A | Modifies aSyn aggregation, clearance and toxicity ⁵⁹ |
| RAB27A | Modulates secretion of aSyn ⁶² | |
| RAB39B | Modulates steady-state levels of aSyn, oligomerization and toxicity ^{59,67} | |
| LRRK2 | RAB3A/B/C/D | LRRK2 kinase substrates ^{68,69} |
| | RAB5 | Implicated with LRRK2 and Rab11 in <i>Drosophila</i> synaptic vesicle recycling ^{70,71} |
| | RAB7L1 | Phosphorylated by LRRK2. LRRK2 and Rab7L1 interact in the endolysosomal system ^{70,72-79} |
| | RAB8A | LRRK2 kinase substrate, interacts with LRRK and RAB7L1 in endosomal homeostasis. LRRK2-mediated phosphorylation of RAB8A leads to centrosomal alterations ^{68,78,80-82} |
| | RAB10 | LRRK2 kinase substrate, involved with LRRK and RAB7L1 in endosomal homeostasis ^{68,78,80-82} |
| | RAB12 | LRRK2 kinase substrate ⁶⁸ |
| | RAB32 | Directly interacts with LRRK and is linked to SNX6/retromer trafficking at the Golgi ^{83,84} |
| | RAB35 | LRRK2 kinase substrate ⁶⁸ |
| | RAB43 | LRRK2 kinase substrate ⁶⁸ |
| VPS35 | RAB5 | Implicated with VPS35, LRRK, and Rab11 in <i>Drosophila</i> synaptic vesicle endocytosis ⁷¹ |
| | RAB7 | Recruits retromer on endosomes via interactions with the Vps sub-complex ⁸⁵⁻⁸⁷ |
| | RAB7L1 | Indication of functional relationship between LRRK, RAB7L1, and VPS35 ⁷² |
| PINK1 | RAB8A/B | PINK1-induced phosphorylation alters the ability of RAB8A to interact with its GEF Rabin8 ⁸⁸⁻⁹⁰ |
| | RAB13 | Phosphorylated after PINK1 activation ⁸⁸ |
| PARKIN | RAB5 | Recruited to damaged mitochondria after Parkin-mediated ubiquitination of RABGEF1 ⁹¹ |
| | RAB7A | Recruited to damaged mitochondria after Parkin-mediated ubiquitination of RABGEF1. Parkin also regulates the activity of Rab7 in the endo-lysosomal pathway ⁹¹⁻⁹⁴ |

Common familial Parkinson disease (PD) genes, alpha-synuclein (aSyn), leucine-rich repeat kinase 2 (LRRK2), vacuolar protein sorting-associated protein 35 (VSP35), PTEN-induced kinase (PINK1) and PARKIN are listed alongside their associated RAB proteins and functional outcome. Abbreviations: RAB, Ras analog in brain; ER, endoplasmic reticulum; GEF, guanine nucleotide exchange factor.

despite the identification of a number of XLID- and LBD-affected families which carry *RAB39B* mutations, large-scale studies of LBD cohorts have failed to find significant prevalence among Caucasian or Asian populations.¹⁰⁵⁻¹⁰⁸ Thus, *RAB39B* mutations are likely rare and do not seem to contribute to the occurrence of LBD outwith XLID cases. However, the symptomatic and pathological recapitulation of LBDs following the loss of RAB39B clearly highlights the protein's functions as critical for neuronal viability and for aSyn homeostasis. Consequently, alterations in RAB39B may still contribute to the pathogenesis of LBDs in the wider population. Indeed, ourselves and others have recently

reported on the loss and/or redistribution of RAB39B in idiopathic LBD variants.^{97,109}

RAB39B and Endosomal Trafficking

The human *RAB39B* gene, identified in 2002, encodes for a protein with 74.2% homology with RAB39A.¹¹⁰ Phylogenetic analysis and subfamily segregation suggests that in addition to RAB39A, RAB39B is also closely related to RAB 2, 4, 11A/B, and 25, each of which are involved in the trafficking of endosomes.^{111,112}

TABLE 2 *RAB39B* gene mutations, symptomology, and neuropathology

| Mutations | Molecular consequence | Symptomology | Pathology |
|---------------------------|---|---|--|
| c.21C > A p.Y7X | Nonsense mutation/loss of expression | Mental impairment/autism/seizures | Macrocephaly ⁹⁶ |
| c.215 + 1G > A | Intronic mutation/loss of expression | Mental impairment/autism/seizures | Macrocephaly ⁹⁶ |
| 45 kb deletion | Gene deletion/loss of expression | Mental impairment (non-progressive)PD symptomology/onset ~45 years of age | Macrocephaly Crt and SCrt LBs/NFTs Iron deposition ⁶⁷ |
| C.503C > A p.T168K | Missense mutation Reduce stability/loss of expression | Mental impairment (non-progressive) /seizuresPD symptomology /onset ~20 years of age | Macrocephaly Crt and SCrt LBs/NFTs Iron deposition SCrt atrophy ⁶⁷ |
| c.557G > A p.T186X | Nonsense mutation /loss of expression | Mild mental impairment (non-progressive)PD symptomology/onset~39 years oldExecutive function deficits and mood disorder | NR ⁹⁹ |
| c.574G > A p.G192R | Missense mutation Impaired membrane association and function | Mild mental impairment (non-progressive) PD symptomology/onset ~50 years of age | NR ⁹⁸ |
| c.428C > G p.A143G | Missense mutation | PD symptomology/onset 47 years of age | NR ⁹⁸ |
| c.624_626delGAG p.R209del | Deletion mutation | PD symptomology/onset 67 years of age | NR ⁹⁸ |
| c.432delA p.T145Tfs*3 | Deletion mutation/loss of expression | Mental impairmentPD symptomology/onset 29 years of age | Abnormal DAT and SPECT signals Iron deposition ¹⁰⁰ |
| c.123G > T p.V41V | Silent mutation In silico cryptic splice site determined | Normal intellectual capacityPD symptomology/onset 45 years of age | NR ¹⁰⁰ |
| c.536dupA p.I180Afs*48 | Duplication mutation/loss of expression | Mental impairmentPD symptomology/onset ~12 years of age | GP atrophyBG calcification ¹⁰¹ |
| c.137dupT p.S47L.fs*44 | Duplication mutation/loss of expression | Mental impairment PD symptomology/onset ~60 years of ageExecutive function deficits and mood disorders | Abnormal SPECT signalSN and GP atrophyBG calcification ¹⁰² |
| c.371delA p.K124S.fs*10 | Deletion mutation/loss of expression | Mental impairmentPD symptomology/onset ~44 years of ageExecutive function deficits and mood disorders | Abnormal SPECT signalSN and GP atrophy ¹⁰² |
| c.559G > T p.E187X | Missense mutation | Mental impairment/autism Motor impairment/tremor | NR ¹⁰³ |
| 0.5 Mb dul Xq28 | Duplication of RAB39B and 7 other genes | Mental impairment | NR ¹⁰⁴ |

Mutations are cited as per changes in codon (c.) and protein amino acid sequence, mutations resulting in frameshifts (fs) and position of induced stop codon (*) are also indicated. Molecular consequences, generalized symptomology, and onset age of Parkinson's disease symptoms are provided.

Abbreviations: Crt, cortical; sCrt, subcortical; PD, Parkinson's disease; LB, Lewy body; NFT, neurofibrillary tangle; NR, not reported; DAT, dopamine transporter; SPECT, single photon emission computed tomography; GP, globus pallidus; BG, basal ganglia; SN, substantia nigra.

Despite its assignment to a group of endosomal trafficking RABs, little is definitively understood about the exact function of RAB39B in this process. At a conceptual level the endosomal system regulates the fate of endocytosed cargo, which is initially sorted in early endosomes, from which the internalized proteins are trafficked for either degradation in late endosomes, ultimately entering lysosomal pathways, or are sorted for the return to the plasma membrane via exocytosis. The process of exocytosis can be further spilt into either a direct rapid route (endosomes to plasma membrane) or via slow endocytic recycling compartments at times also involving retrograde transport from the endosomes to the Golgi apparatus.¹¹³

In both mice and humans RAB39B is highly enriched in brain neurons and is developmentally upregulated after birth, with expression being highest within the hippocampus, neocortex, and substantia nigra.^{96,97} When expressed in various cell types including in primary hippocampal neurons, RAB39B colocalizes with VAMP4 and syntaxin 16, markers of retrograde Golgi trafficking, as well as with ER, Golgi, ER-Golgi trafficking markers, but also in early endosomes and within slow endosomal recycling and post-Golgi secretory pathways where it partially colocalizes with RAB11.^{67,96} The mediation of endocytotic recycling via RAB39B is further supported by its association with myosin Va, a post-Golgi actin-based motor protein,¹¹⁴ as well as its partial colocalization with trans-Golgi network (TGN) protein p230, known to influence transport from the TGN to the plasma membrane.⁶⁷ Recently, RAB39B has also been localized with ER and ER-Golgi trafficking markers.¹¹⁵ Collectively, these studies implicate RAB39B in a variety of trafficking events predominately associated with endocytotic retrograde and/or early-stage anterograde secretory transport (Fig. 2A). Therefore, alterations in RAB39B function likely have widespread consequences for cellular trafficking and without further investigation the predominant consequences for neuronal homeostasis as a result of disrupted vesicle trafficking are difficult to discern. Nevertheless, a number of key studies have begun to shed light on this.

RAB39B and Neurite Elongation and Synaptic Development

At a subcellular level, RAB39B is enriched in the growth cones (GCs) of developing neurites. Intriguingly, both knockdown and overexpression of RAB39B results in a reduction in GC number and in neurite length.^{96,104} This impaired neuritic outgrowth likely contributes to the reduced number of presynaptic terminals also observed following the modulation of RAB39B expression.^{96,104} Such deficits may relate to

the improper regulation of membrane remodeling at the GC leading edge, as efficient endosomal recycling is required for removal and insertion of guidance and adhesion-based receptors and lipids.^{116,117} Although the exact mechanism for this disruption of neuronal maturation has not been established, such impaired neurite outgrowth and pathfinding likely contribute to the developmental abnormalities seen in those carrying loss of function mutations within the *RAB39B* gene.

RAB39B and Glutamatergic Receptor Maturation/Modulation

RAB39B also affects the maturation of AMPA receptors (AMPA) subunits, a process that involves a change from the predominantly Ca²⁺-permeable GluA1 AMPAR subunits to include Ca²⁺-impermeable GluA2-3 subunits,¹¹⁸ thereby promoting the adoption of the classical Na⁺-based electrochemical signalling of AMPARs. Knockdown of RAB39B in neurons results in the accumulation of GluA2 and GluA3 subunits within the cell body, which fail to traffic into the dendrites, ultimately reducing their surface expression and altering AMPAR-mediated postsynaptic currents.¹¹⁹ In this context, RAB39B appears to interact with protein interacting with C kinase (PICK1), which itself associates with GluA2 within the ER and mediates the trafficking of GluA2 subunits through the Golgi and into an early secretory pathway. In contrast, studies focused on a known GEF for RAB39B, C9orf72, have found that following C9orf72 knockout, a loss of postsynaptic RAB39B is observed alongside a corresponding increase in GluA1 postsynaptic localization without change in GluA2 levels.¹²⁰

Independent of the finer details of which subunits are altered, the loss of AMPAR regulation likely has widespread functional consequences. Such changes may be of particular relevance for neurodegeneration as both outcomes favor an increase in Ca²⁺ permeability and thus may increase vulnerability to excitotoxic events.¹²¹ The altered complement of AMPAR subunit expression is consistent with the histopathological study of the cortical expression of these receptors in LBD.¹²² Therefore loss of RAB39B function, either as a consequence of gene mutation or pathological disruption (as discussed below), may contribute to changes in AMPA receptor subunit composition and to a progressive increase in neuronal vulnerability towards Ca²⁺-mediated degenerative insults, thought to participate in the cell death associated with LBDs.¹²³ Despite the findings of these in vitro-based investigations, a recent study of *RAB39B* gene knockout mice has observed a reduction of postsynaptic NMDA receptor subunits as oppose to AMPA receptors, suggesting that in vivo deficits in synaptic function may differ from those established within in vitro models.¹²⁴




FIG. 2. RAB39B-mediated vesicular transport and site of interaction with alpha-synuclein (aSyn) homeostasis. **(A)** Vesicular localization of RAB39B, based on colocalization studies, is indicated alongside key proteins impacted by RAB39B impairments. RAB39B mediates trafficking from the endoplasmic reticulum (ER) to Golgi, influences GluA2 surface expression (i). Localization of RAB39B within the early secretory vesicle, where it may colocalize with membrane-bound aSyn (ii). RAB39B-mediated regulation of endosomal trafficking, localized to early endosomes (EE) and recycling endosomes (RE) alongside internalized GluA1 subunits and aSyn (iii). The association of RAB39B within late endosomes (LE) which feed into lysosomes, generated from RAB39B-positive phagophores is shown alongside its potential to influence aSyn degradation (iv). **(B)** Sites of interaction between a loss of RAB39B function and aSyn accumulation and aggregation. Exocytotic release pathway via vesicular-bound aSyn and endocytotic trafficking of aggregate-prone aSyn via EE and ER towards re-release and via LE to the lysosomal degradation is shown. A loss of RAB39B at each point in the processing pathway (indicated via transparent ball) may lead to increased aSyn retention, accumulation, and aggregation. The disruption of this pathway independent from a loss of function mutation within the RAB39B gene may trigger the deposition of aSyn, in turn trapping key trafficking proteins inclusive of RAB39B, further impeding the homeostatic clearance of aSyn.

RAB39B and Autophagy

RAB39B may also participate in the formation of autophagosomes from the ER membrane. Defects in autophagy due to loss of C9orf72 can be rescued by the expression of constitutively active RAB39B, but not other RABs.¹²⁵ Consistently, endogenous RAB39B colocalizes with the lysosomal marker LAMP1,⁶⁷ and

when overexpressed RAB39B associates with a member of the phosphatidylinositol 3-kinases (PI3Ks) complex initiator of autophagosome formation, Beclin 1.¹²⁶ However, when investigated at endogenous levels the association of RAB39B with Beclin 1¹²⁷ and the localization of RAB39B with LAMP1¹¹⁵ has not been replicated. Nevertheless, in RAB39B gene knockout mice and in responses to the downregulation of RAB39B

gene expression in mouse N2A cells a reduction in autophagolysosome formation has been observed, suggesting decreased autophagic flux.¹²⁴ When autophagy was induced with rapamycin, this impairment was eliminated, indicating that a loss of RAB39B expression impairs basal autophagy, but not autophagy induction. Notably, rapamycin treatment improved defects in synaptic plasticity and memory observed in RAB39B gene knockout mice, suggesting that autophagy plays a central role in phenotypes observed with RAB39B deficiency.¹²⁴

Regulation of aSyn Homeostasis Via RAB39B

Of clear relevance to the association of RAB39B with PD is the potential of the protein to modulate intracellular levels of aSyn. Initial studies suggested that the experimental knockdown of RAB39B resulted in an overall reduction in steady-state levels of aSyn in primary neuronal preparations.⁶⁷ Yet, in contrast, in neuroglioma cells, a reduction of RAB39B expression facilitated aSyn oligomerization and aggregation and was associated with increased cellular toxicity.⁵⁹ As the former finding is contrary to the accumulation of aSyn observed in human postmortem studies, it may be that this contradiction is a consequence of interactions between the role of RAB39B in synaptic development and that of a direct interaction with aSyn homeostasis. Indeed, during development an absence of RAB39B may perturb the development and maturation of synapses, resulting in a reduction of several synaptic proteins required for normal function, including aSyn, whilst beyond the developmental period, the same absence of RAB39B may promote the cellular retention of aSyn and its aggregation.

Although the mechanisms by which a loss of RAB39B leads to the dyshomeostasis of aSyn is unknown, the disruption of autophagic clearance has recently been proposed as central to this, in line with the impairments seen following the expression of mutations within the RAB39B GEF C9orf72¹²⁸ and would appear to be supported by the autophagy deficits in RAB39B knockout mice.¹²⁴ Whilst such a proposition is interesting it remains unclear why such generalized failure of lysosomal clearance would preferentially lead to the accumulation and aggregation of aSyn over other aggregate prone proteins, and indeed several studies have failed to find an impact upon lysosomal degradation following the loss of RAB39B.^{126,127} Equally, it must also be considered that Lewy pathology is not commonly reported in cases of amyotrophic lateral sclerosis and frontotemporal disease associated with the loss of function hexanucleotide repeat expansion of the

c9orf72 gene¹²⁹ and thus such a close relationship between the mode of cellular dysfunction induced by a loss of RAB39B activity and the loss of c9orf72 activity would seem unlikely. Nevertheless, rare cases of PD linked to c9orf72 mutation have been reported,¹³⁰ perhaps indicative of a partial overlap in defective pathways, with innate or environmental factors modifying the cellular outcome to one more closely aligned with those mediated by the loss of RAB39B.

Here, we would propose that the function of RAB39B relates not only to autophagy but also to endosomal trafficking particularly within retrograde trafficking vesicles, early and late endosomes endosomes,⁶⁷ and recycling endosomes where it colocalizes with RAB11.⁹⁶ Thus, akin to other PD-associated mutations within endosomal trafficking regulators such as the retrograde trafficking protein VPS35,^{131,132} the early endosomal-associated DNAJ13¹³³ and the RAB-regulating LRRK2,⁶⁸ the loss of RAB39B may perturb essential endosomal trafficking of aSyn which in addition to compromised autophagic clearance leads to its aggregation.

Such a model would suggest that the compromised endosomal pathways are likely to impact upon several different pools of aSyn including the significant portion of the aSyn which is processed for extracellular release either through exosomal and/or exocytotic pathways.¹³⁴ In neurons, vesicles containing intraluminal aSyn are released via an atypical ER-Golgi-independent exocytosis pathway, in a process which is intimately linked with lysosomal degradation rates.^{135,136} Rather critically, vesicular aSyn has a greater propensity for aggregation compared to aSyn within the cytoplasm,^{135,137} suggesting that the rapid processing of vesicular-bound aSyn is of high importance to minimize the potential for aSyn deposition. Although the exact regulation of aSyn exocytosis remains to be resolved, it is known that the process requires the activity of the RAB11a,^{135,137} which is both spatially⁹⁶ and functional^{111,112} related to RAB39B.

The endosomal transport of aSyn is not only relevant to the regulation of de novo synthesized aSyn but also to the extracellular pools of aSyn, particularly oligomeric and aggregates species, which enter cells via a variety of internalization processes.¹³⁴ Internalized aSyn oligomers and aggregates rapidly enter endosomal pathways colocalizing with early and late endosomal markers and are largely trafficked into the lysosomes for degradation.^{138,139} However, again, a significant proportion of the internalized aSyn can also be recycled back into the extracellular environment, utilizing a similar RAB11-dependent atypical ER-Golgi-independent pathway as employed for the release of intracellularly produced aSyn.⁶⁰

Thus, efficient endosomal trafficking of both endogenous and exogenous aSyn pools would appear as critical not only to the regulation of total intracellular aSyn

abundance but also in the clearance of aggregation-prone aSyn species. Moreover, these studies highlight aSyn endosomal trafficking as a major potential site of dysfunction, which may underlie the association of RAB39B mutations and aSyn aggregation.

Whilst the exact point of dysfunction within the endosomal pathway induced by an absence of RAB39 is unknown, be that in early, recycling, or retrograde endosomes or indeed in the regulation of autophagy, it is noteworthy that the axonal and synaptic expression of RAB39B^{96,120} is in line with reports of the initial sites of aSyn aggregate formation within these subcellular compartments preceding their trafficking and maturation into somatic LBs.¹⁴⁰⁻¹⁴³ Thus, in the absence of a key regulator of endosomal trafficking, such as RAB39B, the entrance of vesicle-bound aSyn into either degradation or exocytosis pathways may stall, with the prolonged retention of aSyn leading to aggregation and deposition. Likewise, in accordance with several studies reporting the disruption of endosomal trafficking following aSyn overexpression,^{46,144,145} should the abundance of vesicular-bound aSyn exceed the endosomal handling capacity of the cell, blockage and aggregation within the pathway may ensue. The impact of such initial deposition of aSyn with endosomal vesicles to cellular homeostasis would be two-fold, as in addition to the formation of toxic aSyn species, the formation of Lewy pathology around cargo-bearing vesicles may entrap a variety of essential lipids and proteins including RAB proteins themselves, further propagating intracellular aggregation (Fig. 2B). This secondary consequence of vesicular aggregation formation is consistent with our previous observation of sequestration of RAB11a and RAB13 in cellular models of aSyn deposition as well as the coaggregation of RAB39B in a subpopulation of LBs in idiopathic DLB cases,¹⁰⁹ which together implicate the dysfunction of key RAB proteins for both familial cases of LBD and idiopathic variants. Furthermore, should a loss of functional RAB39B be mediated by its inclusion within LBs, affected neurons maybe further impacted by downstream alterations of synaptic homeostasis as a consequence of altered glutamatergic signalling and autophagic deficits.

Pharmacological Targeting of RAB Proteins

Although it must be acknowledged that the direct targeting of RAB39B activity in those carrying XLID-associated loss of function mutations would be fruitless, there may be sufficient functional overlap in the system such that the targeting of closely related RAB proteins such as RAB11 may be a beneficial line of investigation. Furthermore, the above outlined potential of a wider relevancy for targeting RAB proteins, RAB39B, and

others in idiopathic cases has prompted a clear interest in identifying therapeutic strategies aimed at their modulation.

Conceptually, a number of interaction sites for targeting RAB proteins exist including RAB prenylation, either acting at the C-terminus of the RAB protein itself or at the GGTase enzyme to regulate membrane interactions, the modulation of RAB activation either at the point of GTP binding, or in the expression and colocalization of GEFs, GAPs, and GDIs or indeed via the expression and turnover of RAB proteins themselves.¹⁴⁶ Nevertheless, targeting RAB proteins is in general challenging due to the high sequence homology amongst GTPase families and the strong affinity of small GTPase for GTP (~pM), largely negating attempts for competitive nucleotide antagonism. To date, the majority of efforts have been focused on reducing the activation of RABs in line with their overactivation in cancer.¹⁴⁷ This work has led to a number of promising, albeit rather non-specific, compounds such as the broad-spectrum GTPases inhibitor CID1067700¹⁴⁸ and several GGTases inhibitors such as psoromic acid¹⁴⁹ and 3-(3-pyridyl)-2-hydroxy-2-phosphonopropanoic acid.¹⁵⁰ Despite the potential for improved specificity to be gained from targeting GEF, GAPs, and GDIs, relatively little success has been achieved in this respect; however, their continued characterization may yet offer the best opportunities for the development of small molecules targeting RAB activity.^{4,151} Such approaches have proven fruitful at least within the related Ras GTPase families, with Ras GEF inhibitors NSC-658497 against SOS1¹⁵² and NPPD against TRIO-GEF1D,¹⁵³ as well as an inhibitor of the Rho GAP, male germ cell Rac GAP known as MINC1¹⁵⁴ having been established. In this respect the identification of a number of RAB39B GEFs such as C9orf72¹⁵⁵ and DENN domain (DENND) proteins DENND5A/B¹⁵⁶ and GAPs such as TBC1D18/RABGAP1L and RUTBC3^{157,158} may prove advantageous in the search for pharmacological targets.

In parallel, progress in identifying allosteric sites and modulators of RAB proteins is also being made.¹⁵⁹ High-throughput screens have identified promising RAB activators, these derivatives of salicylic, indole, and nicotinic acid stabilize the GTP-bound structure of RAB2 and RAB7, independent of associated GEF and GAP, yet lack robust specificity also activating Ras and other Ras-related GTPases.¹⁶⁰ Nevertheless, this recent progress rebukes the former “undruggable” status of GTPases and the emerging compounds may serve as the basis for future improved drug development. Despite much work still being required to elucidate any potential mechanisms of direct pharmacological interventions, several drugs have been identified to modulate downstream pathways affected by RAB gene mutations.¹⁴⁶ For example, the treatment of neuroblastoma

cells expressing CMT2B mutant RAB7 genes with valproic acid can overcome deficient neurite outgrowth via readdressing disruptions to C-Jun N-terminal kinase pathways.¹⁶¹ Furthermore, the targeting of cholesterol to the plasma membrane can overcome the RAB11-mediated deficit in cholesterol esterification within Niemann–Pick type C1 fibroblasts.¹⁶² Thus, the targeting of RABs directly or correcting impacted RAB functions may prove beneficial in the further treatment of neurodegenerative conditions.

In this regard, it is of relevance that in striatal neurons, the orphan G-protein-coupled receptor 52 (GRP52) was found to enhance HTT toxicity via the activation of a RAB39B GEF, acting in opposition to the Rabgap11 GAP which is epistatically expressed in relation to GPR52.¹⁶³ The identification of a regulatory receptor capable of mediating changes in the activity of RAB39B clearly holds potential for the development of relevant agonists or antagonists in order to readdress alterations in RAB39B activity. Nevertheless, given the potential for enhanced toxicity of some neurodegenerative proteins, careful consideration following extensive investigation will have to be conducted when seeking to modulate levels/activity of this protein within the aging brain.

Outlook

Clearly much remains to be determined about the intricacies of intracellular trafficking routes and how neurons utilize such pathways for the regulation and clearance of aSyn. Whilst genetic associations place RAB39B at the centre of aSyn dysfunction and thus LB deposition, the specific point of interaction between the two is not currently determined. Future research focused on the uptake, transport, and release of vesicular-bound aSyn following the manipulation of RAB39B activity is now required to further delineate the consequences of aSyn retention and aggregation. Moreover, studies investigating the potential for compensation and recovery of this system via the augmented activation of related RAB proteins may serve to validate various therapeutic strategies.

Similarly, should RAB39B prove a viable future target, establishing the route of clearance and thus the fate of RAB39B trafficked aSyn should be considered of high importance and will likely further inform potential therapeutic approaches. For example, whilst current studies focused on the health of individual cells would appear to support a protective role for RAB39B-mediated trafficking in the clearance of intracellular aSyn, caution must be exercised, as if RAB39B functions serve to facilitate the extracellular release of aSyn as opposed to its ultimate degradation, its further activation may augment prion-like spread of pathology and thus be detrimental in the context of the whole

brain as opposed to that of a single cell. Despite the low frequency of RAB39B mutations within the human population and the confounding XLID presentation of carriers, the recent observations made by ourselves and others of a disruption of RAB39B subcellular distribution and also of its sequestration in LBs in idiopathic cases of LBD,^{97,109} further strengthens the need to clarify the relationship between RAB39B and aSyn and indeed its role in the formation of concomitant disease-modifying pathologies.

In the absence of mutations within its gene and/or the genes of its effectors, it is unknown how RAB39B may contribute to the development of age-related neurodegeneration onset. It is plausible that declining RAB39B levels or activation in line with age may lead to intercellular trafficking becoming vulnerable to disruption. This aging-related decline in RAB39B may in turn sensitize neurons towards previously subthreshold stressors, inclusive of familial and risk gene mutations which summate to impact upon vesicular trafficking and consequently facilitate the accumulation and aggregation of aSyn. Critically, whilst indeed some key endocytosis proteins are increased in their expression with age,^{164–166} a detailed profile of how the expression of each RAB protein and its modulators (GEFs/GAPs, etc.) alters over the course of human aging is as yet undefined. Thus, the extent to which such an age-induced vulnerability contributes to a tipping point of aggregation and cellular dysfunction is uncertain. Nevertheless continued research into the functional roles of RAB39B in protein and lipid trafficking will serve to clarify its significance in cellular development, homeostasis, and in the pathology of LBDs and other neurodegenerative diseases. ■

Acknowledgments: D.K. and T.F.O. are supported by the Lewy Body Society (LBS-0007). T.F.O. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC 2067/1- 390729940, and SFB1286 (Project B8). S.C. and F.G. are supported by funding from Parkinson's UK (G-1802). Initial work into RAB39B was supported by an award from the Alzheimer's Research UK Newcastle Network Centre to D.K., T.F.O., and F.G.

References

- Li G, Marlin MC. Rab family of GTPases. *Methods Mol Biol* 2015;1298:1–15. https://doi.org/10.1007/978-1-4939-2569-8_1
- Zhang J, Schulze KL, Hiesinger PR, et al. Thirty-one flavors of drosophila Rab proteins. *Genetics* 2007;176:1307–1322. <https://doi.org/10.1534/genetics.106.066761>
- Zerial M, McBride H. Rab proteins as membrane organizers. *Nat Rev Mol Cell Biol* 2001;2:107–117. <https://doi.org/10.1038/35052055>
- Müller MP, Goody RS. Molecular control of Rab activity by GEFs, GAPs and GDI. *Small GTPases* 2018;9:5–21. <https://doi.org/10.1080/21541248.2016.1276999>
- Leung KF, Baron R, Seabra MC. Thematic review series: lipid post-translational modifications. Geranylgeranylation of Rab GTPases.

- J Lipid Res 2006;47:467–475. <https://doi.org/10.1194/jlr.R500017-JLR200>
6. Hutagalung AH, Novick PJ. Role of Rab GTPases in membrane traffic and cell physiology. *Physiol Rev* 2011;91:119–149. <https://doi.org/10.1152/physrev.00059.2009>
 7. Barr F, Lambright DG. Rab GEFs and GAPs. *Curr Opin Cell Biol* 2010;22:461–470. <https://doi.org/10.1016/j.ccb.2010.04.007>
 8. Wang D, Chan C-C, Cherry S, Hiesinger PR. Membrane trafficking in neuronal maintenance and degeneration. *Cell Mol Life Sci* 2013;70:2919–2934. <https://doi.org/10.1007/s00018-012-1201-4>
 9. Kiral FR, Kohrs FE, Jin EJ, Hiesinger PR. Rab GTPases and membrane trafficking in neurodegeneration. *Curr Biol* 2018;28:R471–r486. <https://doi.org/10.1016/j.cub.2018.02.010>
 10. Villarroel-Campos D, Gastaldi L, Conde C, Caceres A, Gonzalez-Billault C. Rab-mediated trafficking role in neurite formation. *J Neurochem* 2014;129:240–248. <https://doi.org/10.1111/jnc.12676>
 11. D'Adamo P, Masetti M, Bianchi V, et al. RAB GTPases and RAB-interacting proteins and their role in the control of cognitive functions. *Neurosci Biobehav Rev* 2014;46(Pt 2):302–314. <https://doi.org/10.1016/j.neubiorev.2013.12.009>
 12. Di Giovanni S, Knights CD, Rao M, et al. The tumor suppressor protein p53 is required for neurite outgrowth and axon regeneration. *EMBO J* 2006;25:4084–4096. <https://doi.org/10.1038/sj.emboj.7601292>
 13. Shikanai M, Yuzaki M, Kawachi T. Rab family small GTPases-mediated regulation of intracellular logistics in neural development. *Histol Histopathol* 2018;33:765–771. <https://doi.org/10.14670/hh-11-956>
 14. Veleri S, Punnakkal P, Dunbar GL, Maiti P. Molecular insights into the roles of Rab proteins in intracellular dynamics and neurodegenerative diseases. *Neuromolecular Med* 2018;20:18–36. <https://doi.org/10.1007/s12017-018-8479-9>
 15. Pavlos NJ, Grønberg M, Riedel D, et al. Quantitative analysis of synaptic vesicle Rabs uncovers distinct yet overlapping roles for Rab3a and Rab27b in Ca²⁺-triggered exocytosis. *J Neurosci* 2010;30:13441–13453. <https://doi.org/10.1523/jneurosci.0907-10.2010>
 16. Graf ER, Daniels RW, Burgess RW, Schwarz TL, DiAntonio A. Rab3 dynamically controls protein composition at active zones. *Neuron* 2009;64:663–677. <https://doi.org/10.1016/j.neuron.2009.11.002>
 17. Yu E, Kanno E, Choi S, et al. Role of Rab27 in synaptic transmission at the squid giant synapse. *Proc Natl Acad Sci* 2008;105:16003–16008. <https://doi.org/10.1073/pnas.0804825105>
 18. Star EN, Newton AJ, Murthy VN. Real-time imaging of Rab3a and Rab5a reveals differential roles in presynaptic function. *J Physiol* 2005;569:103–117. <https://doi.org/10.1113/jphysiol.2005.092528>
 19. Semerdjieva S, Shortt B, Maxwell E, et al. Coordinated regulation of AP2 uncoating from clathrin-coated vesicles by rab5 and hRME-6. *J Cell Biol* 2008;183:499–511. <https://doi.org/10.1083/jcb.200806016>
 20. McLauchlan H, Newell J, Morrice N, Osborne A, West M, Smythe E. A novel role for Rab5-GDI in ligand sequestration into clathrin-coated pits. *Curr Biol* 1998;8:34–45. [https://doi.org/10.1016/S0960-9822\(98\)70018-1](https://doi.org/10.1016/S0960-9822(98)70018-1)
 21. Wucherpfennig T, Wilsch-Bräuninger M, González-Gaitán M. Role of drosophila Rab5 during endosomal trafficking at the synapse and evoked neurotransmitter release. *J Cell Biol* 2003;161:609–624. <https://doi.org/10.1083/jcb.200211087>
 22. Shimizu H, Kawamura S, Ozaki K. An essential role of Rab5 in uniformity of synaptic vesicle size. *J Cell Sci* 2003;116:3583–3590. <https://doi.org/10.1242/jcs.00676>
 23. Wit H d, Lichtenstein Y, Kelly RB, Geuze HJ, Klumperman J, van der Sluijs P. Rab4 regulates formation of synaptic-like microvesicles from early endosomes in PC12 cells. *Mol Biol Cell* 2001;12:3703–3715. <https://doi.org/10.1091/mbc.12.11.3703>
 24. Esteves da Silva M, Adrian M, Schätzle P, et al. Positioning of AMPA receptor-containing endosomes regulates synapse architecture. *Cell Rep* 2015;13:933–943. <https://doi.org/10.1016/j.celrep.2015.09.062>
 25. Uytterhoeven V, Kuenen S, Kasproicz J, Miskiewicz K, Verstreken P. Loss of Skywalker reveals synaptic endosomes as sorting stations for synaptic vesicle proteins. *Cell* 2011;145:117–132. <https://doi.org/10.1016/j.cell.2011.02.039>
 26. Sheehan P, Zhu M, Beskow A, Vollmer C, Waites CL. Activity-dependent degradation of synaptic vesicle proteins requires Rab35 and the ESCRT pathway. *J Neurosci* 2016;36:8668–8686. <https://doi.org/10.1523/jneurosci.0725-16.2016>
 27. Guerra F, Bucci C. Multiple roles of the small GTPase Rab7. *Cell* 2016;5:34. <https://doi.org/10.3390/cells5030034>
 28. Binotti B, Jahn R, Chua JJ. Functions of Rab proteins at presynaptic sites. *Cells* 2016;5. <https://doi.org/10.3390/cells5010007>
 29. Binotti B, Pavlos NJ, Riedel D, et al. The GTPase Rab26 links synaptic vesicles to the autophagy pathway. *eLife* 2015;4:e05597. <https://doi.org/10.7554/eLife.05597>
 30. Brown TC, Tran IC, Backos DS, Esteban JA. NMDA receptor-dependent activation of the small GTPase Rab5 drives the removal of synaptic AMPA receptors during hippocampal LTD. *Neuron* 2005;45:81–94. <https://doi.org/10.1016/j.neuron.2004.12.023>
 31. Park M, Penick EC, Edwards JG, Kauer JA, Ehlers MD. Recycling endosomes supply AMPA receptors for LTP. *Science* 2004;305:1972–1975. <https://doi.org/10.1126/science.1102026>
 32. Brown TC, Correia SS, Petrok CN, Esteban JA. Functional compartmentalization of endosomal trafficking for the synaptic delivery of AMPA receptors during long-term potentiation. *J Neurosci* 2007;27:13311–13315. <https://doi.org/10.1523/jneurosci.4258-07.2007>
 33. Gerges NZ, Backos DS, Esteban JA. Local control of AMPA receptor trafficking at the postsynaptic terminal by a small GTPase of the Rab family. *J Biol Chem* 2004;279:43870–43878. <https://doi.org/10.1074/jbc.m404982200>
 34. Bacaj T, Ahmad M, Jurado S, Malenka RC, Südhof TC. Synaptic function of Rab11Fip5: selective requirement for hippocampal long-term depression. *J Neurosci* 2015;35:7460–7474. <https://doi.org/10.1523/jneurosci.1581-14.2015>
 35. Mori Y, Fukuda M, Henley JM. Small GTPase Rab17 regulates the surface expression of kainate receptors but not α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors in hippocampal neurons via dendritic trafficking of Syntaxin-4 protein. *J Biol Chem* 2014;289:20773–20787. <https://doi.org/10.1074/jbc.M114.550632>
 36. Bucci C, De Luca M. Molecular basis of Charcot-Marie-Tooth type 2B disease. *Biochem Soc Trans* 2012;40:1368–1372. <https://doi.org/10.1042/bst20120197>
 37. Bem D, Yoshimura S-I, Nunes-Bastos R, et al. Loss-of-function mutations in RAB18 cause Warburg micro syndrome. *Am J Hum Genet* 2011;88:499–507. <https://doi.org/10.1016/j.ajhg.2011.03.012>
 38. Sveinbjornsdottir S. The clinical symptoms of Parkinson's disease. *J Neurochem* 2016;139(Suppl 1):318–324. <https://doi.org/10.1111/jnc.13691>
 39. Kalia LV, Kalia SK. α -Synuclein and Lewy pathology in Parkinson's disease. *Curr Opin Neurol* 2015;28:375–381. <https://doi.org/10.1097/wco.0000000000000215>
 40. Outeiro TF, Koss DJ, Erskine D, et al. Dementia with Lewy bodies: an update and outlook. *Mol Neurodegener* 2019;14:5. <https://doi.org/10.1186/s13024-019-0306-8>
 41. Brás IC, Dominguez-Mejide A, Gerhardt E, et al. Synucleinopathies: where we are and where we need to go. *J Neurochem* 2020;153:433–454. <https://doi.org/10.1111/jnc.14965>
 42. Smolders S, Van Broeckhoven C. Genetic perspective on the synergistic connection between vesicular transport, lysosomal and mitochondrial pathways associated with Parkinson's disease pathogenesis. *Acta Neuropathol Commun* 2020;8:63. <https://doi.org/10.1186/s40478-020-00935-4>
 43. Tang B, Rabs L. Membrane dynamics, and Parkinson's disease. *J Cell Physiol* 2017;232:1626–1633. <https://doi.org/10.1002/jcp.25713>

44. Shi M-M, Shi C-H, Xu Y-M. Rab GTPases: the key players in the molecular pathway of Parkinson's disease. *Front Cell Neurosci* 2017;11. <https://doi.org/10.3389/fncel.2017.00081>
45. Gao Y, Wilson GR, Stephenson SEM, Bozaoglu K, Farrer MJ, Lockhart PJ. The emerging role of Rab GTPases in the pathogenesis of Parkinson's disease. *Mov Disord* 2018;33:196–207. <https://doi.org/10.1002/mds.27270>
46. Gitler AD, Bevis BJ, Shorter J, et al. The Parkinson's disease protein alpha-synuclein disrupts cellular Rab homeostasis. *Proc Natl Acad Sci U S A* 2008;105:145–150. <https://doi.org/10.1073/pnas.0710685105>
47. Cooper AA, Gitler AD, Cashikar A, et al. α -Synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science* 2006;313:324–328. <https://doi.org/10.1126/science.1129462>
48. Dalfó E, Barrachina M, Rosa JL, Ambrosio S, Ferrer I. Abnormal alpha-synuclein interactions with rab3a and rabphilin in diffuse Lewy body disease. *Neurobiol Dis* 2004;16:92–97. <https://doi.org/10.1016/j.nbd.2004.01.001>
49. Dalfó E, Gómez-Isla T, Rosa JL, et al. Abnormal alpha-synuclein interactions with Rab proteins in alpha-synuclein A30P transgenic mice. *J Neuropathol Exp Neurol* 2004;63:302–313. <https://doi.org/10.1093/jnen/63.4.302>
50. Dalfó E, Ferrer I. Alpha-synuclein binding to rab3a in multiple system atrophy. *Neurosci Lett* 2005;380:170–175. <https://doi.org/10.1016/j.neulet.2005.01.034>
51. Chen RH, Wislet-Gendebien S, Samuel F, et al. α -Synuclein membrane association is regulated by the Rab3a recycling machinery and presynaptic activity. *J Biol Chem* 2013;288:7438–7449. <https://doi.org/10.1074/jbc.M112.439497>
52. Wang TY, Ma Z, Wang C, et al. Manganese-induced alpha-synuclein overexpression impairs synaptic vesicle fusion by disrupting the Rab3 cycle in primary cultured neurons. *Toxicol Lett* 2018;285:34–42. <https://doi.org/10.1016/j.toxlet.2017.12.024>
53. Sung JY, Kim J, Paik SR, Park JH, Ahn YS, Chung KC. Induction of neuronal cell death by Rab5A-dependent endocytosis of alpha-synuclein. *J Biol Chem* 2001;276:27441–27448. <https://doi.org/10.1074/jbc.M101318200>
54. Cheng F, Li X, Li Y, et al. α -Synuclein promotes clathrin-mediated NMDA receptor endocytosis and attenuates NMDA-induced dopaminergic cell death. *J Neurochem* 2011;119:815–825. <https://doi.org/10.1111/j.1471-4159.2011.07460.x>
55. Fang F, Yang W, Florio JB, et al. Synuclein impairs trafficking and signaling of BDNF in a mouse model of Parkinson's disease. *Sci Rep* 2017;7:3868. <https://doi.org/10.1038/s41598-017-04232-4>
56. Freeze B, Acosta D, Pandya S, Zhao Y, Raj A. Regional expression of genes mediating trans-synaptic alpha-synuclein transfer predicts regional atrophy in Parkinson disease. *Neuroimage Clin* 2018;18:456–466. <https://doi.org/10.1016/j.nicl.2018.01.009>
57. Masaracchia C, Hnida M, Gerhardt E, et al. Membrane binding, internalization, and sorting of alpha-synuclein in the cell. *Acta Neuropathol Commun* 2018;6:79. <https://doi.org/10.1186/s40478-018-0578-1>
58. Germann UA, Alam JJ. P38 α MAPK signaling—a robust therapeutic target for Rab5-mediated neurodegenerative disease. *Int J Mol Sci* 2020;21. <https://doi.org/10.3390/ijms21155485>
59. Gonçalves SA, Macedo D, Raquel H, et al. shRNA-based screen identifies endocytic recycling pathway components that act as genetic modifiers of alpha-synuclein aggregation, secretion and toxicity. *PLoS Genet* 2016;12:e1005995. <https://doi.org/10.1371/journal.pgen.1005995>
60. Liu J, Zhang J-P, Shi M, et al. Rab11a and HSP90 regulate recycling of extracellular alpha-synuclein. *J Neurosci* 2009;29:1480–1485. <https://doi.org/10.1523/jneurosci.6202-08.2009>
61. Dinter E, Saridakis T, Nippold M, et al. Rab7 induces clearance of α -synuclein aggregates. *J Neurochem* 2016;138:758–774. <https://doi.org/10.1111/jnc.13712>
62. Ejlerskov P, Rasmussen I, Nielsen TT, et al. Tubulin polymerization-promoting protein (TPPP/p25 α) promotes unconventional secretion of α -synuclein through exophagy by impairing autophagosome-lysosome fusion. *J Biol Chem* 2013;288:17313–17335. <https://doi.org/10.1074/jbc.M112.401174>
63. Yin G, da Fonseca TL, Eisbach SE, et al. α -Synuclein interacts with the switch region of Rab8a in a Ser129 phosphorylation-dependent manner. *Neurobiol Dis* 2014;70:149–161. <https://doi.org/10.1016/j.nbd.2014.06.018>
64. Poehler AM, Xiang W, Spitzer P, et al. Autophagy modulates SNCA/ α -synuclein release, thereby generating a hostile microenvironment. *Autophagy* 2014;10:2171–2192. <https://doi.org/10.4161/auto.36436>
65. Chutna O, Gonçalves S, Villar-Piqué A, et al. The small GTPase Rab11 co-localizes with α -synuclein in intracellular inclusions and modulates its aggregation, secretion and toxicity. *Hum Mol Genet* 2014;23:6732–6745. <https://doi.org/10.1093/hmg/ddu391>
66. Breda C, Nugent ML, Estranero JG, et al. Rab11 modulates α -synuclein-mediated defects in synaptic transmission and behaviour. *Hum Mol Genet* 2015;24:1077–1091. <https://doi.org/10.1093/hmg/ddu521>
67. Wilson GR, Sim JCH, McLean C, et al. Mutations in RAB39B cause X-linked intellectual disability and early-onset Parkinson disease with alpha-synuclein pathology. *Am J Hum Genet* 2014;95:729–735. <https://doi.org/10.1016/j.ajhg.2014.10.015>
68. Steger M, Diez F, Dhekne HS, et al. Systematic proteomic analysis of LRRK2-mediated Rab GTPase phosphorylation establishes a connection to ciliogenesis. *eLife* 2017;6:e31012. <https://doi.org/10.7554/eLife.31012>
69. Islam MS, Nolte H, Jacob W, et al. Human R1441C LRRK2 regulates the synaptic vesicle proteome and phosphoproteome in a drosophila model of Parkinson's disease. *Hum Mol Genet* 2016;25:5365–5382. <https://doi.org/10.1093/hmg/ddw352>
70. Boon JY, Dusonchet J, Trengrove C, Wolozin B. Interaction of LRRK2 with kinase and GTPase signaling cascades. *Front Mol Neurosci* 2014;7:64. <https://doi.org/10.3389/fnmol.2014.00064>
71. Inoshita T, Arano T, Hosaka Y, et al. Vps35 in cooperation with LRRK2 regulates synaptic vesicle endocytosis through the endosomal pathway in drosophila. *Hum Mol Genet* 2017;26:2933–2948. <https://doi.org/10.1093/hmg/ddx179>
72. MacLeod DA, Rhinn H, Kuwahara T, et al. RAB7L1 interacts with LRRK2 to modify intraneuronal protein sorting and Parkinson's disease risk. *Neuron* 2013;77:425–439. <https://doi.org/10.1016/j.neuron.2012.11.033>
73. Beilina A, Rudenko IN, Kaganovich A, et al. Unbiased screen for interactors of leucine-rich repeat kinase 2 supports a common pathway for sporadic and familial Parkinson disease. *Proc Natl Acad Sci U S A* 2014;111:2626–2631. <https://doi.org/10.1073/pnas.1318306111>
74. Pihlström L, Rengmark A, Björnarå KA, et al. Fine mapping and resequencing of the PARK16 locus in Parkinson's disease. *J Hum Genet* 2015;60:357–362. <https://doi.org/10.1038/jhg.2015.34>
75. Kuwahara T, Inoue K, D'Agati VD, et al. LRRK2 and RAB7L1 coordinately regulate axonal morphology and lysosome integrity in diverse cellular contexts. *Sci Rep* 2016;6:29945. <https://doi.org/10.1038/srep29945>
76. Fujimoto T, Kuwahara T, Eguchi T, Sakurai M, Komori T, Iwatsubo T. Parkinson's disease-associated mutant LRRK2 phosphorylates Rab7L1 and modifies trans-Golgi morphology. *Biochem Biophys Res Commun* 2018;495:1708–1715. <https://doi.org/10.1016/j.bbrc.2017.12.024>
77. Liu Z, Bryant N, Kumaran R, et al. LRRK2 phosphorylates membrane-bound Rabs and is activated by GTP-bound Rab7L1 to promote recruitment to the trans-Golgi network. *Hum Mol Genet* 2018;27:385–395. <https://doi.org/10.1093/hmg/ddx410>
78. Eguchi T, Kuwahara T, Sakurai M, et al. LRRK2 and its substrate Rab GTPases are sequentially targeted onto stressed lysosomes and maintain their homeostasis. *Proc Natl Acad Sci U S A* 2018;115:E9115–e9124. <https://doi.org/10.1073/pnas.1812196115>
79. Madero-Pérez J, Fernández B, Ordóñez AJL, et al. RAB7L1-mediated relocalization of LRRK2 to the Golgi complex causes centrosomal deficits via RAB8A. *Front Mol Neurosci* 2018;11:417. <https://doi.org/10.3389/fnmol.2018.00417>

80. Kuwahara T, Funakawa K, Komori T, et al. Roles of lysosomotropic agents on LRRK2 activation and Rab10 phosphorylation. *Neurobiol Dis* 2020;145:105081. <https://doi.org/10.1016/j.nbd.2020.105081>
81. Petridi S, Middleton CA, Ugbo C, Fellgett A, Covill L, Elliott CJH. In vivo visual screen for dopaminergic Rab ↔ LRRK2-G2019S interactions in drosophila discriminates Rab10 from Rab3. *G3 (Bethesda)* 2020;10:1903–1914. <https://doi.org/10.1534/g3.120.401289>
82. Lis P, Burel S, Steger M, et al. Development of phospho-specific Rab protein antibodies to monitor in vivo activity of the LRRK2 Parkinson's disease kinase. *Biochem J* 2018;475:1–22. <https://doi.org/10.1042/bcj20170802>
83. Waschbüsch D, Michels H, Strassheim S, et al. LRRK2 transport is regulated by its novel interacting partner Rab32. *PLoS One* 2014;9:e111632. <https://doi.org/10.1371/journal.pone.0111632>
84. Waschbüsch D, Hübel N, Ossendorf E, et al. Rab32 interacts with SNX6 and affects retromer-dependent Golgi trafficking. *PLoS One* 2019;14:e0208889. <https://doi.org/10.1371/journal.pone.0208889>
85. Seaman MN, Harbour ME, Tattersall D, Read E, Bright N. Membrane recruitment of the cargo-selective retromer subcomplex is catalysed by the small GTPase Rab7 and inhibited by the Rab-GAP TBC1D5. *J Cell Sci* 2009;122:2371–2382. <https://doi.org/10.1242/jcs.048686>
86. Liu TT, Gomez TS, Sackey BK, Billadeau DD, Burd CG. Rab GTPase regulation of retromer-mediated cargo export during endosome maturation. *Mol Biol Cell* 2012;23:2505–2515. <https://doi.org/10.1091/mbc.E11-11-0915>
87. Priya A, Kalaidzidis IV, Kalaidzidis Y, Lambright D, Datta S. Molecular insights into Rab7-mediated endosomal recruitment of core retromer: deciphering the role of Vps26 and Vps35. *Traffic* 2015;16:68–84. <https://doi.org/10.1111/tra.12237>
88. Lai YC, Kondapalli C, Lehneck R, et al. Phosphoproteomic screening identifies Rab GTPases as novel downstream targets of PINK1. *EMBO J* 2015;34:2840–2861. <https://doi.org/10.15252/embj.201591593>
89. Pourjafar-Dehkordi D, Vieweg S, Itzen A, Zacharias M. Phosphorylation of Ser111 in Rab8a modulates Rabin8-dependent activation by perturbation of side chain interaction networks. *Biochemistry* 2019;58:3546–3554. <https://doi.org/10.1021/acs.biochem.9b00516>
90. Vieweg S, Mulholland K, Bräuning B, et al. PINK1-dependent phosphorylation of Serine111 within the SF3 motif of Rab GTPases impairs effector interactions and LRRK2-mediated phosphorylation at Threonine72. *Biochem J* 2020;477:1651–1668. <https://doi.org/10.1042/bcj20190664>
91. Yamano K, Wang C, Sarraf SA, et al. Endosomal Rab cycles regulate Parkin-mediated mitophagy. *eLife* 2018;7. <https://doi.org/10.7554/eLife.31326>
92. Song P, Trajkovic K, Tsunemi T, Krainc D. Parkin modulates endosomal organization and function of the endo-lysosomal pathway. *J Neurosci* 2016;36:2425–2437. <https://doi.org/10.1523/jneurosci.2569-15.2016>
93. Yamano K, Fogel AI, Wang C, van der Bliek AM, Youle RJ. Mitochondrial Rab GAPs govern autophagosome biogenesis during mitophagy. *eLife* 2014;3:e01612. <https://doi.org/10.7554/eLife.01612>
94. Heo JM, Ordureau A, Swarup S, et al. RAB7A phosphorylation by TBK1 promotes mitophagy via the PINK-PARKIN pathway. *Sci Adv* 2018;4:eaav0443. <https://doi.org/10.1126/sciadv.aav0443>
95. Tarpey PS, Smith R, Pleasance E, et al. A systematic, large-scale resequencing screen of X-chromosome coding exons in mental retardation. *Nat Genet* 2009;41:535–543. <https://doi.org/10.1038/ng.367>
96. Giannandrea M, Bianchi V, Mignogna ML, et al. Mutations in the small GTPase gene RAB39B are responsible for X-linked mental retardation associated with autism, epilepsy, and macrocephaly. *Am J Hum Genet* 2010;86:185–195. <https://doi.org/10.1016/j.ajhg.2010.01.011>
97. Gao Y, Martínez-Cerdeño V, Hogan KJ, McLean CA, Lockhart PJ. Clinical and neuropathological features associated with loss of RAB39B. *Mov Disord* 2020;35(4):687–693. <https://doi.org/10.1002/mds.27951>
98. Mata IF, Jang Y, Kim C-H, et al. The RAB39B p.G192R mutation causes X-linked dominant Parkinson's disease. *Mol Neurodegener* 2015;10(50). <https://doi.org/10.1186/s13024-015-0045-4>
99. Lesage S, Bras J, Cormier-Dequaire F, et al. Loss-of-function mutations in RAB39B are associated with typical early-onset Parkinson disease. *Neurol Genet* 2015;1:e9. <https://doi.org/10.1212/nxg.0000000000000009>
100. Güldner M, Schulte C, Hauser AK, Gasser T, Brockmann K. Broad clinical phenotype in parkinsonism associated with a base pair deletion in RAB39B and additional POLG variant. *Parkinsonism Relat Disord* 2016;31:148–150. <https://doi.org/10.1016/j.parkreldis.2016.07.005>
101. Shi CH, Zhang S-Y, Yang Z-H, et al. A novel RAB39B gene mutation in X-linked juvenile parkinsonism with basal ganglia calcification. *Mov Disord* 2016;31:1905–1909. <https://doi.org/10.1002/mds.26828>
102. Ciammola A, Carrera P, Di Fonzo A, et al. X-linked parkinsonism with intellectual disability caused by novel mutations and somatic mosaicism in RAB39B gene. *Parkinsonism Relat Disord* 2017;44:142–146. <https://doi.org/10.1016/j.parkreldis.2017.08.021>
103. Woodbury-Smith M, Deneault E, Yuen RKC, et al. Mutations in RAB39B in individuals with intellectual disability, autism spectrum disorder, and macrocephaly. *Mol Autism* 2017;8:59. <https://doi.org/10.1186/s13229-017-0175-3>
104. Vanmarsenille L, Giannandrea M, Fieremans N, et al. Increased dosage of RAB39B affects neuronal development and could explain the cognitive impairment in male patients with distal Xq28 copy number gains. *Hum Mutat* 2014;35:377–383. <https://doi.org/10.1002/humu.22497>
105. Hodges K, Brewer SS, Labbé C, et al. RAB39B gene mutations are not a common cause of Parkinson's disease or dementia with Lewy bodies. *Neurobiol Aging* 2016;45:107–108. <https://doi.org/10.1016/j.neurobiolaging.2016.03.021>
106. Lin HH, Wu R-M, Lin H-I, Chen M-L, Tai C-H, Lin C-H. Lack of RAB39B mutations in early-onset and familial Parkinson's disease in a Taiwanese cohort. *Neurobiol Aging* 2017;50:169.e163–169.e164. <https://doi.org/10.1016/j.neurobiolaging.2016.10.021>
107. Löchte T, Brüggemann N, Vollstedt E-J, et al. RAB39B mutations are a rare finding in Parkinson disease patients. *Parkinsonism Relat Disord* 2016;23:116–117. <https://doi.org/10.1016/j.parkreldis.2015.12.014>
108. Yuan L, Deng X, Song Z, et al. Genetic analysis of the RAB39B gene in Chinese Han patients with Parkinson's disease. *Neurobiol Aging* 2015;36:2907.e2911–2907.e2902. <https://doi.org/10.1016/j.neurobiolaging.2015.06.019>
109. Koss DJ, Bondarevaite O, Adams S, et al. RAB39B is redistributed in dementia with Lewy bodies and is sequestered within aβ plaques and Lewy bodies. *Brain Pathol* 2020;31(1):120–132. <https://doi.org/10.1111/bpa.12890>
110. Cheng H, Ma Y, Ni X, et al. Isolation and characterization of a human novel RAB (RAB39B) gene. *Cytogenet Genome Res* 2002;97:72–75. <https://doi.org/10.1159/000064047>
111. Pereira-Leal JB, Seabra MC. Evolution of the Rab family of small GTP-binding proteins. *J Mol Biol* 2001;313:889–901. <https://doi.org/10.1006/jmbi.2001.5072>
112. Klöpffer TH, Kienle N, Fasshauer D, Munro S. Untangling the evolution of Rab G proteins: implications of a comprehensive genomic analysis. *BMC Biol* 2012;10:71. <https://doi.org/10.1186/1741-7007-10-71>
113. Naslavsky N, Caplan S. The enigmatic endosome - sorting the ins and outs of endocytic trafficking. *J Cell Sci* 2018;131. <https://doi.org/10.1242/jcs.216499>
114. Lindsay AJ, Jollivet F, Horgan CP, et al. Identification and characterization of multiple novel Rab-myosin Va interactions. *Mol Biol Cell* 2013;24:3420–3434. <https://doi.org/10.1091/mbc.E13-05-0236>
115. Gambarte Tudela J, Buonfigli J, Luján A, et al. Rab39a and Rab39b display different intracellular distribution and function in

- sphingolipids and phospholipids transport. *Int J Mol Sci* 2019;20. <https://doi.org/10.3390/ijms20071688>
116. Erez H, Malkinson G, Prager-Khoutorsky M, De Zeeuw CI, Hoogenraad CC, Spira ME. Formation of microtubule-based traps controls the sorting and concentration of vesicles to restricted sites of regenerating neurons after axotomy. *J Cell Biol* 2007;176:497–507. <https://doi.org/10.1083/jcb.200607098>
 117. Yap CC, Winckler B. Harnessing the power of the endosome to regulate neural development. *Neuron* 2012;74:440–451. <https://doi.org/10.1016/j.neuron.2012.04.015>
 118. Bassani S, Folci A, Zapata J, Passafaro M. AMPAR trafficking in synapse maturation and plasticity. *Cell Mol Life Sci* 2013;70:4411–4430. <https://doi.org/10.1007/s00018-013-1309-1>
 119. Mignogna ML, Giannandrea M, Gurgone A, et al. The intellectual disability protein RAB39B selectively regulates GluA2 trafficking to determine synaptic AMPAR composition. *Nat Commun* 2015;6:6504. <https://doi.org/10.1038/ncomms7504>
 120. Xiao S, McKeever PM, Lau A, Robertson J. Synaptic localization of C9orf72 regulates post-synaptic glutamate receptor 1 levels. *Acta Neuropathol Commun* 2019;7:161. <https://doi.org/10.1186/s40478-019-0812-5>
 121. Weiss JH. Ca permeable AMPA channels in diseases of the nervous system. *Front Mol Neurosci* 2011;4:42–42. <https://doi.org/10.3389/fnmol.2011.00042>
 122. Mohamed NE, Howlett DR, Ma L, et al. Decreased immunoreactivities of neocortical AMPA receptor subunits correlate with motor disability in Lewy body dementias. *J Neural Transm (Vienna)* 2014;121:71–78. <https://doi.org/10.1007/s00702-013-1067-0>
 123. Zaichick SV, McGrath KM, Caraveo G. The role of Ca(2+) signaling in Parkinson's disease. *Dis Model Mech* 2017;10:519–535. <https://doi.org/10.1242/dmm.028738>
 124. Niu M, Zheng N, Wang Z, et al. RAB39B deficiency impairs learning and memory partially through compromising autophagy. *Front Cell Dev Biol* 2020;8. <https://doi.org/10.3389/fcell.2020.598622>
 125. Sellier C, Campanari M-L, Corbier CJ, et al. Loss of C9ORF72 impairs autophagy and synergizes with polyQ Ataxin-2 to induce motor neuron dysfunction and cell death. *EMBO J* 2016;35:1276–1297. <https://doi.org/10.15252/embj.201593350>
 126. Seto S, Sugaya K, Tsujimura K, Nagata T, Horii T, Koide Y. Rab39a interacts with phosphatidylinositol 3-kinase and negatively regulates autophagy induced by lipopolysaccharide stimulation in macrophages. *PLoS One* 2013;8:e83324. <https://doi.org/10.1371/journal.pone.0083324>
 127. Zhang W, Ma L, Yang M, et al. Cerebral organoid and mouse models reveal a RAB39b-PI3K-mTOR pathway-dependent dysregulation of cortical development leading to macrocephaly/autism phenotypes. *Genes Dev* 2020;34:580–597. <https://doi.org/10.1101/gad.332494.119>
 128. Tang BL. RAB39B's role in membrane traffic, autophagy, and associated neuropathology. *J Cell Physiol* 2021;236:1579–1592. <https://doi.org/10.1002/jcp.29962>
 129. Majounie E, Renton AE, Mok K, et al. Frequency of the C9orf72 hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. *Lancet Neurol* 2012;11:323–330. [https://doi.org/10.1016/s1474-4422\(12\)70043-1](https://doi.org/10.1016/s1474-4422(12)70043-1)
 130. Wilke C, Pomper JK, Biskup S, Puskás C, Berg D, Synofzik M. Atypical parkinsonism in C9orf72 expansions: a case report and systematic review of 45 cases from the literature. *J Neurol* 2016;263:558–574. <https://doi.org/10.1007/s00415-016-8021-7>
 131. Zimprich A, Benet-Pagès A, Struhal W, et al. A mutation in VPS35, encoding a subunit of the retromer complex, causes late-onset Parkinson disease. *Am J Hum Genet* 2011;89:168–175. <https://doi.org/10.1016/j.ajhg.2011.06.008>
 132. Vilariño-Güell C, Wider C, Ross OA, et al. VPS35 mutations in Parkinson disease. *Am J Hum Genet* 2011;89:162–167. <https://doi.org/10.1016/j.ajhg.2011.06.001>
 133. Yoshida S, Hasegawa T, Suzuki M, et al. Parkinson's disease-linked DNAJC13 mutation aggravates alpha-synuclein-induced neurotoxicity through perturbation of endosomal trafficking. *Hum Mol Genet* 2018;27:823–836. <https://doi.org/10.1093/hmg/ddy003>
 134. Grozdanov V, Danzer KM. Release and uptake of pathological alpha-synuclein. *Cell Tissue Res* 2018;373:175–182. <https://doi.org/10.1007/s00441-017-2775-9>
 135. Lee HJ, Patel S, Lee SJ. Intravesicular localization and exocytosis of alpha-synuclein and its aggregates. *J Neurosci* 2005;25:6016–6024. <https://doi.org/10.1523/jneurosci.0692-05.2005>
 136. Jang A, Lee H-J, Suk J-E, Jung J-W, Kim K-P, Lee S-J. Non-classical exocytosis of alpha-synuclein is sensitive to folding states and promoted under stress conditions. *J Neurochem* 2010;113:1263–1274. <https://doi.org/10.1111/j.1471-4159.2010.06695.x>
 137. Hasegawa T, Konno M, Baba T, et al. The AAA-ATPase VPS4 regulates extracellular secretion and lysosomal targeting of alpha-synuclein. *PLoS One* 2011;6:e29460. <https://doi.org/10.1371/journal.pone.0029460>
 138. Lee HJ, Suk J-E, Bae E-J, Lee J-H, Paik SR, Lee S-J. Assembly-dependent endocytosis and clearance of extracellular alpha-synuclein. *Int J Biochem Cell Biol* 2008;40:1835–1849. <https://doi.org/10.1016/j.biocel.2008.01.017>
 139. Desplats P, Lee H-J, Bae E-J, et al. Inclusion formation and neuronal cell death through neuron-to-neuron transmission of alpha-synuclein. *Proc Natl Acad Sci U S A* 2009;106:13010–13015. <https://doi.org/10.1073/pnas.09036911106>
 140. Katsuse O, Iseki E, Marui W, Kosaka K. Developmental stages of cortical Lewy bodies and their relation to axonal transport blockage in brains of patients with dementia with Lewy bodies. *J Neurol Sci* 2003;211:29–35.
 141. Kramer ML, Schulz-Schaeffer WJ. Presynaptic alpha-synuclein aggregates, not Lewy bodies, cause neurodegeneration in dementia with Lewy bodies. *J Neurosci* 2007;27:1405–1410. <https://doi.org/10.1523/jneurosci.4564-06.2007>
 142. Volpicelli-Daley LA, Gamble KL, Schultheiss CE, Riddle DM, West AB, Lee VM-Y. Formation of alpha-synuclein Lewy neurite-like aggregates in axons impedes the transport of distinct endosomes. *Mol Biol Cell* 2014;25:4010–4023. <https://doi.org/10.1091/mbc.E14-02-0741>
 143. Volpicelli-Daley LA, Luk KC, Patel TP, et al. Exogenous alpha-synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death. *Neuron* 2011;72:57–71. <https://doi.org/10.1016/j.neuron.2011.08.033>
 144. Outeiro TF, Lindquist S. Yeast cells provide insight into alpha-synuclein biology and pathobiology. *Science* 2003;302:1772–1775. <https://doi.org/10.1126/science.1090439>
 145. Sancenon V, Lee S-A, Patrick C, et al. Suppression of alpha-synuclein toxicity and vesicle trafficking defects by phosphorylation at S129 in yeast depends on genetic context. *Hum Mol Genet* 2012;21:2432–2449. <https://doi.org/10.1093/hmg/dds058>
 146. Agola JO, Jim PA, Ward HH, Basuray S, Wandinger-Ness A. Rab GTPases as regulators of endocytosis, targets of disease and therapeutic opportunities. *Clin Genet* 2011;80:305–318. <https://doi.org/10.1111/j.1399-0004.2011.01724.x>
 147. Qin X, Wang J, Wang X, Liu F, Jiang B, Zhang Y. Targeting Rabs as a novel therapeutic strategy for cancer therapy. *Drug Discov Today* 2017;22:1139–1147. <https://doi.org/10.1016/j.drudis.2017.03.012>
 148. Hong L, Guo Y, BasuRay S, et al. A pan-GTPase inhibitor as a molecular probe. *PLoS One* 2015;10:e0134317. <https://doi.org/10.1371/journal.pone.0134317>
 149. Deraeve C, Guo Z, Bon RS, Blankenfeldt W, et al. Psoromic acid is a selective and covalent Rab-prenylation inhibitor targeting auto-inhibited RabGGTase. *J Am Chem Soc* 2012;134:7384–7391. <https://doi.org/10.1021/ja211305j>
 150. Roelofs AJ, Hulley PA, Meijer A, Ebetino FH, Russell RGG, Shipman CM. Selective inhibition of Rab prenylation by a phosphonocarboxylate analogue of risedronate induces apoptosis, but not S-phase arrest, in human myeloma cells. *Int J Cancer* 2006;119:1254–1261. <https://doi.org/10.1002/ijc.21977>
 151. Cherfils J, Zeghouf M. Regulation of small GTPases by GEFs, GAPs, and GDIs. *Physiol Rev* 2013;93:269–309. <https://doi.org/10.1152/physrev.00003.2012>

152. Evelyn CR, Duan X, Biesiada J, Seibel WL, Meller J, Zheng Y. Rational design of small molecule inhibitors targeting the Ras GEF, SOS1. *Chem Biol* 2014;21:1618–1628. <https://doi.org/10.1016/j.chembiol.2014.09.018>
153. Blangy A, Bouquier N, Gauthier-Rouvière C, et al. Identification of TRIO-GEF1 chemical inhibitors using the yeast exchange assay. *Biol Cell* 2006;98:511–522. <https://doi.org/10.1042/bc20060023>
154. van Adrichem AJ, Fagerholm A, Turunen L, et al. Discovery of MINC1, a GTPase-activating protein small molecule inhibitor, targeting MgcRacGAP. *Comb Chem High Throughput Screen* 2015; 18:3–17. <https://doi.org/10.2174/1386207318666141205112730>
155. Corbier C, Sellier C. C9ORF72 is a GDP/GTP exchange factor for Rab8 and Rab39 and regulates autophagy. *Small GTPases* 2017;8: 181–186. <https://doi.org/10.1080/21541248.2016.1212688>
156. Yoshimura S, Gerondopoulos A, Linford A, Rigden DJ, Barr FA. Family-wide characterization of the DENN domain Rab GDP-GTP exchange factors. *J Cell Biol* 2010;191:367–381. <https://doi.org/10.1083/jcb.201008051>
157. Fukuda M. TBC proteins: GAPs for mammalian small GTPase Rab? *Biosci Rep* 2011;31:159–168. <https://doi.org/10.1042/bsr20100112>
158. Itoh T, Satoh M, Kanno E, Fukuda M. Screening for target Rabs of TBC (Tre-2/Bub2/Cdc16) domain-containing proteins based on their Rab-binding activity. *Genes Cells* 2006;11:1023–1037. <https://doi.org/10.1111/j.1365-2443.2006.00997.x>
159. Kumar AP, Verma CS, Lukman S. Structural dynamics and allostery of Rab proteins: strategies for drug discovery and design. *Brief Bioinform* 2020;22(1):270–287. <https://doi.org/10.1093/bib/bbz161>
160. Palsuledesai CC, Surviladze Z, Waller A, et al. Activation of rho family GTPases by small molecules. *ACS Chem Biol* 2018;13: 1514–1524. <https://doi.org/10.1021/acscchembio.8b00038>
161. Yamauchi J, Torii T, Kusakawa S, et al. The mood stabilizer valproic acid improves defective neurite formation caused by Charcot-Marie-Tooth disease-associated mutant Rab7 through the JNK signaling pathway. *J Neurosci Res* 2010;88:3189–3197. <https://doi.org/10.1002/jnr.22460>
162. Wiegand V, Chang TY, Strauss JF III, Fahrenholz F, Gimpl G. Transport of plasma membrane-derived cholesterol and the function of Niemann-Pick C1 protein. *FASEB J* 2003;17:782–784. <https://doi.org/10.1096/fj.02-0818fje>
163. Yao Y, Cui X, Al-Ramahi I, et al. A striatal-enriched intronic GPCR modulates huntingtin levels and toxicity. *eLife* 2015;4. <https://doi.org/10.7554/eLife.05449>
164. Ginsberg SD, Alldred MJ, Counts SE, et al. Microarray analysis of hippocampal CA1 neurons implicates early endosomal dysfunction during Alzheimer's disease progression. *Biol Psychiatry* 2010;68: 885–893. <https://doi.org/10.1016/j.biopsych.2010.05.030>
165. Ginsberg SD, Mufson EJ, Alldred MJ, et al. Upregulation of select Rab GTPases in cholinergic basal forebrain neurons in mild cognitive impairment and Alzheimer's disease. *J Chem Neuroanat* 2011; 42:102–110. <https://doi.org/10.1016/j.jchemneu.2011.05.012>
166. Ginsberg SD, Mufson EJ, Counts SE, et al. Regional selectivity of rab5 and rab7 protein upregulation in mild cognitive impairment and Alzheimer's disease. *J Alzheimers Dis* 2010;22:631–639. <https://doi.org/10.3233/jad-2010-101080>