

Poster presentation

Open Access

Extensive chaotic dynamics in neural networks in the balanced state

Michael Kreissl*^{1,2,3}, Siegrid Löwel^{4,5} and Fred Wolf^{1,2,3}

Address: ¹Department of Nonlinear Dynamics, Max-Planck-Institute for Dynamics and Self-Organization, D37073 Göttingen, Germany, ²Bernstein Center for Computational Neuroscience, D37073 Göttingen, Germany, ³Faculty of Physics, University of Göttingen, D37073 Göttingen, Germany, ⁴Institute of General Zoology and Animal Physiology, Friedrich-Schiller-University Jena, D07743 Jena, Germany and ⁵Bernstein Group for Computational Neuroscience, D07743 Jena, Germany

Email: Michael Kreissl* - kreissl@nld.ds.mpg.de

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P330 doi:10.1186/1471-2202-10-S1-P330

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P330>

© 2009 Kreissl et al; licensee BioMed Central Ltd.

Neurons embedded in operational cortical networks fire action potentials in highly irregular sequences [1]. The balanced state of cortical networks provides an attractive explanation of this temporally irregular behaviour [2]. In this state, neurons are driven by large input fluctuations, resulting from a dynamically achieved balance of excitation and inhibition. It has been shown that the balanced state can robustly emerge from the collective dynamics of spiking neuron networks [2]. The detailed dynamics of the balanced state, however, is not well understood. Initially, Vreeswijk and Sompolinsky [2] found a kind of "hyper"-chaotic dynamics in networks of binary neurons, characterized by an infinite positive Lyapunov exponent, which is hard to reconcile with classical notions of nonlinear dynamics. More recently, Zillmer et al. [3] and Jahnke et al. [4] reported stable dynamics in networks of leaky integrate and fire neurons. Notably, all of these previous works neglect a major cellular mechanism of dynamic instability, the instability of the membrane voltage near the spike threshold.

To clarify which kind of balanced state dynamics is expected in networks of neuron models describing such dynamic spike generation, we analyzed networks of pulse-coupled theta neurons in sparse, random networks in the balanced state. The theta neuron model is a standard model of type I neuronal excitability [5]. Based on the analytic solution of the single neuron dynamics, we derived a map that was used for event based simulations and obtained the exact single spike Jacobian matrix for the

entire network. This Jacobian was used to calculate the whole set of Lyapunov exponents, following the standard procedure [6]. From the Lyapunov exponents we acquired the entropy production via Pesin's formula and the attractor dimension via the Kaplan-Yorke conjecture.

In general, the studied networks in the balanced state exhibited chaotic dynamics. The largest Lyapunov exponent was positive and finite, giving evidence to conventional chaos. The entropy production and dimension of the chaotic attractor both scaled linearly with the number of neurons in the networks, hence chaos in the balanced state appears to be extensive spatio-temporal. Furthermore, we found a high attractor dimension, corresponding to many effective degrees of freedom that may potentially encode information in the network. Yet, this is accompanied by a rapid loss of information of 0.5 bits per spike. We conclude that extensive chaos should be expected in general for networks of type I neurons in the balanced state. If information loss in cortical networks is as high as in our model, information could hardly be encoded in detailed spiking patterns beyond the immediate stimulus response.

References

1. Softky WR, Koch C: **The highly irregular firing of cortical cells is inconsistent with temporal integration of random EPSPs.** *J Neurosci* 1993, **13**:334-350.
2. Van Vreeswijk C, Sompolinsky H: **Chaotic balanced state in a model of cortical circuits.** *Neural Comput* 1998, **10**:1321-1371.

3. Zillmer R, Livi R, Politi A, Torcini A: **Desynchronization in diluted neural networks.** *Phys Rev E* 2006, **74**:036203.
4. Jahnke S, Memmesheimer RM, Timme M: **Stable irregular dynamics in complex neural networks.** *Phys Rev Lett* 2008, **100**:048102.
5. Gutkin BS, Ermentrout GB: **Dynamics of membrane excitability determine interspike interval variability: a link between spike generation mechanisms and cortical spike train statistics.** *Neural Comput* 1998, **10**:1047-1065.
6. Benettin G, Galgani L, Giorgilli A, Strelcyn JM: **Lyapunov characteristic exponents for smooth dynamical systems and for hamiltonian systems; a method for computing all of them, Part I & Part II.** *Meccanica* 1980, **15**:9-30.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

