- Dunaj-Jurčo, M., Kabešová, M., Kettman, V., Cisařová, I. & Mikloš, D. (1993). *Acta Cryst.* **C49**, 1476–1479.
- Dunaj-Jurčo, M., Ondrejovič, G., Melník, M. & Garaj, J. (1988). *Coord. Chem. Rev.* **83**, 1–28.
- Harrison, W. D. & Hathaway, B. J. (1980). *Acta Cryst.* **B36**, 1069–1074.
- Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.
- Nardelli, M. (1983). *Comput. Chem.* **7**, 95–98.
- Pavelčík, F. (1986). *XP21*. Comenius Univ., Pharmaceutical Faculty, Bratislava, Czechoslovakia.
- Sheldrick, G. M. (1976). *SHELX76*. Program for crystal structure determination. Univ. of Cambridge, England.
- Ugliengo, P., Borzani, G. & Viterbo, D. (1988). *J. Appl. Cryst.* **21**, 75.
- Wicholas, M. & Welford, T. (1974). *Inorg. Chem.* **13**, 316–318.

Acta Cryst. (1993). **C49**, 1482–1483

Structure of Bis(pentamethyldiethylenetriamine)disodium Hexasulfide

SUSANNE BESSER, REGINE HERBST-IRMER AND
DIETMAR STALKE

*Institut für Anorganische Chemie, Universität
Göttingen, Tammannstrasse 4, 3400 Göttingen,
Germany*

ALAN T. BROOKER, RONALD SNAITH AND
DOMENEC S. WRIGHT

*University Chemical Laboratory, Lensfield Road,
Cambridge CB2 1EW, England*

(Received 9 September 1992; accepted 1 February 1993)

Abstract

The new sodium sulfide compound, μ -hexasulfido- $1\kappa^2S^1, S^6:2\kappa^2S^1, S^6$ -bis{[N-(dimethylaminoethyl)-N,N',-N'-trimethyl-1,2-ethanediamine- κ^3N, N', N'']sodium}, $[Na(pmdeta)]_2[S_6]$ (1) [pmdeta = $(Me_2NCH_2CH_2)_2NMe$], was obtained by reacting two equivalents of sodium hydride with elementary sulfur in toluene containing an excess of pmdeata at approximately 238 K. The crystal structure contains a central Na_2S_2 ring in which both S atoms are part of an S_6^{2-} residue. In addition, each Na atom is coordinated by a pmdeata molecule.

Comment

The structure is a contact ion pair where the S_6^{2-} ion adopts a bent zigzag chain and the two Na^+ ions are both bridging the terminal S atoms. Thus a four-membered Na_2S_2 ring is formed. Furthermore, each Na atom is coordinated by a pmdeata molecule. There-

fore, the overall structure is comparable to that of $[Li(tmeda)]_2[S_6]$ (tmeda = $Me_2NCH_2CH_2NMe_2$) (Banister, Barr, Brooker, Clegg, Cunningham, Doyle, Drake, Gill, Manning, Raithby, Snaith, Wade & Wright, 1990), but it is quite different from that of the ionic and acyclic polysulfides BaS_3 , BaS_4 , K_2S_5 and Ca_2S_6 (Wells, 1984). The four-membered Na_2S_2 ring is not planar; it is bent along the $Na1 \cdots Na1^i$ vector. The normal of the $Na1-Na1^i-S3$ plane intersects that of the $Na1-Na1^i-S3^i$ plane with an angle of $19.80(2)^\circ$. The bond lengths within the ring are 2.9073 (9) Å ($Na1-S3$) and 2.8245 (9) Å ($Na1-S3^i$). The bond distances in the S_6^{2-} chain are not of the same length. Whereas the terminal bonds are 2.0344 (7) Å ($S2-S3$) and 2.0487 (7) Å ($S1-S2$), the central $S1-S1^i$ bond is 2.1158 (9) Å long, hence 0.074 Å longer than the others. However, the average S—S bond length is in good agreement with the values for comparable derivatives, such as Cs_2S_6 (Abrahams & Grison, 1953) (average 2.054 Å) and $[Li(tmeda)]_2[S_6]$ (Tatsumi, Inoue, Nakamura, Cramer, VanDorne & Gilje, 1990) (average 1.983 Å).

Fig. 1. Structure of the title compound showing 50% probability displacement ellipsoids. The H atoms are omitted for clarity.

Experimental

Crystal data

$[Na_2(C_9H_{23}N_3)_2][S_6]$
 $M_r = 584.95$
Tetragonal
 $P4_2/c$
 $a = 13.2390(10)$ Å
 $c = 18.400(2)$ Å
 $V = 3225.0(5)$ Å³
 $Z = 4$
 $D_x = 1.205$ Mg m⁻³

Mo $K\alpha$ radiation
 $\lambda = 0.71073$ Å
Cell parameters from 64 reflections
 $\theta = 10-12.5^\circ$
 $\mu = 0.469$ mm⁻¹
 $T = 293.0(10)$ K
 $0.6 \times 0.5 \times 0.4$ mm
Red

Data collection

Stoe Siemens AED four-circle diffractometer
Profile data from $2\theta/\omega$ scans
6917 measured reflections
3567 independent reflections
3273 observed reflections
 $[I > 2\sigma(I)]$
 $R_{int} = 0.0304$

$\theta_{max} = 27.49^\circ$
 $h = -14 \rightarrow 17$
 $k = -12 \rightarrow 12$
 $l = -23 \rightarrow 23$
3 standard reflections
frequency: 90 min
intensity variation: none

Refinement

Refinement on F^2
 Final $R(F) = 0.0269$ for
 $F > 4\sigma(F)$ data
 $wR(F^2) = 0.0616$ for all data
 $S = 1.057$
 3565 reflections
 150 parameters
 Calculated weights
 $w = 1/[\sigma^2(F_o^2) + (0.0202P)^2 + 0.8121P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.194 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.150 \text{ e } \text{\AA}^{-3}$
 Atomic scattering factors
 from *International Tables*
 for *Crystallography* (1992,
 Vol. C, Tables 4.2.6.8 and
 6.1.1.4)
 Absolute configuration ac-
 cording to Flack (1983)

Data collection: *DIF4* (Stoe & Cie, 1988). Cell refinement:
DIF4. Data reduction: *REDU4* (Stoe & Cie, 1988). Program(s)
 used to solve structure: *SHELXS-92* (Sheldrick, 1990). Pro-
 gram(s) used to refine structure: *SHELXL-92* (Sheldrick, 1992).
 Molecular graphics: *SHELXTL-Plus* (Sheldrick, 1991). Soft-
 ware used to prepare material for publication: *SHELXL-92*.

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters ($\text{\AA}^2$)

$$U_{eq} = \frac{1}{3} \sum_i \sum_j U_{ij} a_i^* a_j^* a_i \cdot a_j$$

	x	y	z	U_{eq}
S1	0.07958 (3)	0.50720 (4)	0.98797 (2)	0.0327 (2)
S2	0.11554 (3)	0.63303 (3)	0.92854 (2)	0.0319 (2)
S3	0.10418 (3)	0.60140 (3)	0.82065 (2)	0.0276 (2)
Na1	0.10709 (5)	0.38329 (5)	0.83890 (3)	0.0247 (3)
N1	0.13872 (12)	0.21307 (11)	0.90150 (8)	0.0313 (8)
N2	0.29592 (11)	0.36127 (11)	0.85166 (8)	0.0289 (7)
N3	0.17224 (11)	0.36322 (12)	0.71214 (8)	0.0325 (8)
C11	0.0722 (2)	0.1363 (2)	0.87043 (13)	0.0511 (13)
C12	0.1193 (2)	0.2189 (2)	0.98012 (10)	0.0433 (12)
C13	0.24482 (14)	0.18790 (15)	0.88728 (11)	0.0347 (10)
C21	0.3395 (2)	0.4542 (2)	0.88127 (12)	0.0433 (11)
C22	0.31559 (14)	0.27586 (14)	0.90035 (10)	0.0338 (9)
C23	0.33531 (14)	0.3415 (2)	0.77781 (10)	0.0349 (9)
C31	0.1173 (2)	0.4340 (2)	0.66560 (10)	0.0431 (12)
C32	0.1640 (2)	0.2622 (2)	0.68081 (11)	0.0424 (12)
C33	0.27755 (13)	0.3961 (2)	0.71878 (10)	0.0378 (10)

Table 2. Geometric parameters ($\text{\AA}$, $^\circ$)

S1—S2	2.0487 (7)	S3—Na1	2.9073 (9)
S1—S1 ⁱ	2.1158 (9)	Na1—N3	2.501 (2)
S2—S3	2.0344 (7)	Na1—N2	2.528 (2)
S3—Na1 ⁱ	2.8245 (9)	Na1—N1	2.565 (2)
S2—S1—S1 ⁱ	107.74 (3)	N3—Na1—S3 ⁱ	103.79 (4)
S3—S2—S1	109.65 (3)	N2—Na1—S3 ⁱ	177.09 (4)
S2—S3—Na1 ⁱ	86.70 (2)	N1—Na1—S3 ⁱ	106.14 (4)
S2—S3—Na1	95.21 (2)	N3—Na1—S3	90.14 (4)
Na1 ⁱ —S3—Na1	94.05 (2)	N2—Na1—S3	97.95 (4)
N3—Na1—N2	74.53 (5)	N1—Na1—S3	157.90 (4)
N3—Na1—N1	105.62 (5)	S3 ⁱ —Na1—S3	84.37 (2)
N2—Na1—N1	72.28 (5)		

Symmetry code: (i) $-x, 1 - y, z$.

DS thanks the University of Göttingen and the DAAD (Nato scholarship) for financial support.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and bond distances have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 71149 (13 pp.). Copies may be obtained through The Technical Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England. [CIF reference: HL1023]

References

- Abrahams, S. C. & Grison, E. (1953). *Acta Cryst.* **6**, 206–213.
 Banister, A. J., Barr, D., Brooker, A. T., Clegg, W., Cunningham, M. J., Doyle, M. J., Drake, S. R., Gill, W. R., Manning, K., Raithby, P. R., Snaith, R., Wade, K. & Wright, D. S. (1990). *J. Chem. Soc. Chem. Commun.* pp. 105–107.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
 Sheldrick, G. M. (1991). *SHELXTL-Plus*. Siemens Analytical X-ray Instruments, Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1992). *SHELXL-92*. Program for crystal structure refinement. Univ. of Göttingen, Germany.
 Stoe & Cie (1988). *DIF4*. Version 7.08. Diffractometer control program. Stoe & Cie, Darmstadt, Germany.
 Stoe & Cie (1988). *REDU4*. Data reduction program. Stoe & Cie, Darmstadt, Germany.
 Tatsumi, K., Inoue, Y., Nakamura, A., Cramer, R. E., VanDoorne, W. & Gilje, J. W. (1990). *Angew. Chem.* **102**, 455–457; *Angew. Chem. Int. Ed. Engl.* **29**, 422–424.
 Wells, A. F. (1984). *Structural Inorganic Chemistry*, 5th ed. Oxford Univ. Press.

Acta Cryst. (1993). **C49**, 1483–1485

Structure of μ -[Bis(chloroacetato-*O*)-mercury(II)- $\kappa^2\text{Hg}$]- μ -[1,8-naphthalene-diamino- $\kappa^2\text{NN}'$: $\kappa^2\text{NN}'$]-bis[*cis*-dicarbonyl-(triisopropylphosphine-*P*)ruthenium(I)]-(*Ru*—*Ru*)

S. GARCÍA-GRANDA* AND
 J. F. VAN DER MAELEN URÍA

*Departamento de Química Física y Analítica,
 Facultad de Química, Universidad de Oviedo,
 Julián Clavería s/n, 33006 Oviedo, Spain*

J. A. CABEZA, J. M. FERNÁNDEZ-COLINAS
 AND V. RIERA

*Departamento de Química Orgánica e Inorgánica,
 Facultad de Química, Universidad de Oviedo,
 Julián Clavería s/n, 33006 Oviedo, Spain*

(Received 8 October 1992; accepted 11 February 1993)

Abstract

The molecule shows a twofold symmetry through the Hg atom and the diaminonaphthalene ligand. Ru—Ru and Ru—Hg distances are 2.799 (1) and 2.795 (1) $\text{\AA}$, respectively.

Comment

This work is part of a project concerning the reactivity of the Ru—Ru bonds of binuclear ruthenium(I)